



## POSTER PRESENTATION

## Open Access

# The HIV-1 protective -35SNP effect in Caucasians is CD8 T cell mediated

T Corrah<sup>1\*</sup>, S Brackenridge<sup>2</sup>, N Goonetilleke<sup>2</sup>, H Yang<sup>2</sup>, S Deeks<sup>3</sup>, L Dorrell<sup>2</sup>, M Cohen<sup>4</sup>, A McMichael<sup>2</sup>From AIDS Vaccine 2012  
Boston, MA, USA. 9-12 September 2012

## Background

Previous studies in Caucasians have observed that a single nucleotide polymorphism 35kb upstream of the HLA-C gene (-35SNP) associates with control of HIV-1 viral load set-point and cell surface expression of HLA-C. HIV-1 selectively downregulates HLA-A and HLA-B but not HLA-C, via the action of the Nef protein. Thus it has been speculated that higher cell surface HLA-C expression results in a stronger HLA-C-restricted T cell response which might play a role in the control of HIV-1 replication in individuals with the protective -35C variant. However, HLA-C-restricted CD8 T cell responses are relatively weak and we could find no difference in functional HLA-C-restricted CD8 T cell activity measured by IFN- $\gamma$  ELISPOT assay, according to -35SNP genotype. Therefore, we aimed to examine if there is any correlation between total CD8 T cell function and the -35SNP.

## Methods

The viral suppression assay, which involves directly infecting autologous CD4 T cells with primary HIV-1 strains and co-culturing with autologous CD8 T cells, was used as a surrogate for immune control in vivo. The CD8 T cells from 46 antiretroviral therapy naïve HIV-1 infected Caucasians were assessed using this assay.

## Results

When CD8 T cell antiviral activity was grouped according to -35SNP genotype, the -35CC group possessed significantly higher CD8 T cell antiviral activity than the -35TT group ( $p=0.0151$ ; Mann-Whitney). Protective HLA-B alleles were always in linkage disequilibrium with HLA-C alleles that are in linkage disequilibrium with the -35C allele. Similarly risk HLA-B alleles were in linkage

disequilibrium with HLA-C alleles that are in linkage disequilibrium with the -35T allele.

## Conclusion

In conclusion, the protective -35SNP effect in HIV-1 disease is mediated through CD8 T cells. However, the -35SNP may simply be a marker for protective and risk HLA-B alleles.

## Author details

<sup>1</sup>Addenbrooks Hospital, University of Cambridge, Cambridge, UK. <sup>2</sup>Weatherall Institute of Molecular Medicine, University of Oxford, Oxford, UK. <sup>3</sup>University of California San Francisco, San Francisco, CA, USA. <sup>4</sup>University of North Carolina School of Medicine, NC, USA.

Published: 13 September 2012

doi:10.1186/1742-4690-9-S2-P281

Cite this article as: Corrah et al.: The HIV-1 protective -35SNP effect in Caucasians is CD8 T cell mediated. *Retrovirology* 2012 **9**(Suppl 2):P281.

**Submit your next manuscript to BioMed Central and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at  
[www.biomedcentral.com/submit](http://www.biomedcentral.com/submit)



<sup>1</sup>Addenbrooks Hospital, University of Cambridge, Cambridge, UK  
Full list of author information is available at the end of the article