

Nutritional and socio-economic determinants of cognitive function and educational achievement of Aboriginal schoolchildren in rural Malaysia

Hesham M. Al-Mekhlafi^{1,2*}, Mohammed A. Mahdy^{1,2}, Atiya A. Sallam³, W. A. Ariffin⁴,
Abdulsalam M. Al-Mekhlafi¹, Adel A. Amran⁵ and Johari Surin¹

¹Department of Parasitology, Faculty of Medicine, University of Malaya, Kuala Lumpur, Malaysia

²Department of Parasitology, Faculty of Medicine, Sana'a University, Sana'a, Yemen

³Department of Social and Preventive Medicine, Faculty of Medicine, University of Malaya, Kuala Lumpur, Malaysia

⁴Department of Pediatrics, Faculty of Medicine, University of Malaya, Kuala Lumpur, Malaysia

⁵Department of Physiology, Faculty of Medicine, Universiti Kebangsaan Malaysia, Jalan Raja Muda Abdul Aziz, 50300 Kuala Lumpur, Malaysia

(Received 19 April 2010 – Revised 1 December 2010 – Accepted 21 February 2011 – First published online 15 April 2011)

Abstract

A community-based cross-sectional study was carried out among Aboriginal schoolchildren aged 7–12 years living in remote areas in Pos Betau, Pahang, Malaysia to investigate the potential determinants influencing the cognitive function and educational achievement of these children. Cognitive function was measured by intelligence quotient (IQ), while examination scores of selected school subjects were used in assessing educational achievement. Blood samples were collected to assess serum Fe status. All children were screened for soil-transmitted helminthes. Demographic and socio-economic data were collected using pre-tested questionnaires. Almost two-thirds (67.6%) of the subjects had poor IQ and most of them (72.6%) had insufficient educational achievement. Output of the stepwise multiple regression model showed that poor IQ was significantly associated with low household income which contributed the most to the regression variance (r^2 0.059; $P=0.020$). Low maternal education was also identified as a significant predictor of low IQ scores (r^2 0.042; $P=0.043$). With educational achievement, Fe-deficiency anaemia (IDA) was the only variable to show significant association (r^2 0.025; $P=0.015$). In conclusion, the cognitive function and educational achievement of Aboriginal schoolchildren are poor and influenced by household income, maternal education and IDA. Thus, effective and integrated measures to improve the nutritional and socio-economic status of rural children would have a pronounced positive effect on their education.

Key words: Cognitive function: Educational achievement: Nutrition: Socio-economic status: Aboriginal schoolchildren

Numerous factors are known to affect cognitive function (the mental processes by which knowledge is acquired including perception, reasoning, creativity, problem-solving and possible intuition measured by intelligence quotient (IQ)) and educational achievement of children (the progress of individuals in school measured by the results of examinations). Poor cognitive and educational performances among children can be attributed to a complex web of causation which involves insufficient infrastructure, health problems, poor socio-economic status and environmental causes. Health problems that may affect children's cognitive function and educational achievement include childhood infections^(1,2), malnutrition and micronutrient deficiencies including Fe-deficiency anaemia (IDA)^(3,4), intestinal parasitic infections, mainly soil-transmitted helminthes (STH)⁽⁵⁾, visual and hearing impairment⁽⁶⁾, epilepsy⁽⁷⁾ and many other

problems. Moreover, stimulation-poor environments lead to poor development of the child's brain, which hinder the child's cognitive development^(8,9). Needless to say, poor socio-economic status is a strong inhibitor limiting educational achievement. Such a situation does not motivate children to study and an unsatisfactory home or class environment also does not encourage learning.

Orang Asli (a Malay term transliterated as 'original people') are the indigenous minority inhabitants of Peninsular Malaysia. They comprise only 0.6% of the total population in Malaysia (26.6 million in 2006). Despite intensive efforts by government and private sectors to improve the quality of life of Aboriginal communities throughout 53 years of independence (since 1957), little success has been achieved, and these communities are still plagued with poverty and a wide range of health problems. Very little is known about

Abbreviations: IDA, Fe-deficiency anaemia; IQ, intelligence quotient; STH, soil-transmitted helminthes; TS, total scores of the three school subjects.

* **Corresponding author:** Dr Hesham Al-Mekhlafi, fax +60 3 79764754, email halmekhlafi@yahoo.com

the cognitive functions and educational achievement of rural Aboriginal schoolchildren. Beside the expected shortage in educational facilities, the main problems that impede the educational process for Aboriginal people are poverty, poor health and nutritional status, and a high drop-out rate^(10,11). A national strategy to minimise the educational gap between rural and urban areas has been implemented and more than two million rural schoolchildren, including Aboriginal children, will benefit from this agenda. Hence, the present study was carried out to investigate the potential determinants that influence the cognitive function and educational achievement of Aboriginal schoolchildren in Peninsular Malaysia.

Materials and methods

Study area and population

The present study was carried out between August 2006 and January 2007. Selection of the study area was discussed with the Department of Aboriginal People Affairs and the following criteria were used to select the school: enrolment of more than 250 pupils, easy access and willingness of the teachers to cooperate. Sekolah Kebangsaan Batau (the Batau National School), a primary school for Aboriginal children in Pos Batau, Kuala Lipis, Pahang, Malaysia, met these criteria. The school had an enrolment of 405 pupils in the target age range of 7–12 years. Of these, sixty-nine were absent at the time of the visit, twenty-nine refused to participate and fifteen were excluded for having infections with fever.

This area is considered to be a remote area. Consisting of eighteen villages, it is located in a valley region about 50 km from the town of Kuala Lipis and 200 km northeast of Kuala Lumpur. There is a lack of sanitation with a high prevalence of intestinal parasitic infections, malnutrition and other health problems. There is a clinic in the area for health services equipped with an ambulance to send critical cases to the nearest hospital (50 km). Most of the residents work as farmers, labourers, rubber tappers and some do odd jobs such as selling forest products. Aboriginal individuals live in houses made of wood or bamboo. However, most of the houses have electricity during night-time only and a supply of piped water as the main source for drinking water while water for domestic needs (bathing, washing clothes and utensils and feeding animals) is collected from the rivers located adjacent to the villages.

Throughout many visits to the school and the villages, community meetings were held with the headmaster, heads of the villages, parents and their school-age children before the commencement of the present study in order to give a clear explanation of the objectives and the involvement of the children in the present study. Informed verbal consent was obtained from the participants; verbal consent was witnessed and formally recorded. The present study was conducted according to the guidelines laid down in the Declaration of Helsinki and all procedures involving human subjects were approved by the Medical Ethics Committee of the University of Malaya Medical Centre, University of Malaya, Malaysia.

Study design and sample size

The main design of the present study was a randomised placebo-controlled trial aimed at investigating the effects of vitamin A supplementation on the cognitive function and educational achievement of Aboriginal schoolchildren; the cross-sectional study described in the present paper served as a baseline assessment. A sample size of 214 children, 107 per intervention arm, was estimated to give the study at least 80% power at 5% significance to detect a significant difference in mean scores of IQ and total scores of the three school subjects (TS) between the vitamin A-supplemented group and the placebo group. Although this sample size included an extra 20% to avoid the effects of drop-out, we invited all children in the target age group to participate (universal sampling).

Demographic and socio-economic data

The demographic and socio-economic data of the subjects were collected using a pre-tested questionnaire constructed in English and then translated into Malay, the local language. Socio-economic status was assessed by a set of questions about the parents' educational and employment status, total household income and type and condition of houses. During the visits to the villages, the children and their parents were interviewed in their home settings by trained assistants to elicit information on the subjects' medical history, socio-economic status and personal hygiene practices.

Parasitology

Fresh faecal samples were collected into wide-mouth screw-cap 100 ml clean containers. The samples were examined by the Kato–Katz technique⁽¹²⁾ for the presence of STH, *Ascaris lumbricoides*, *Trichuris trichiura* and hookworm eggs. Egg counts, as a measure of worm burden, were also carried out using this technique and the results were recorded as eggs per g of stool (epg). The intensity of infection was graded as heavy, moderate or light according to criteria proposed by the WHO⁽¹³⁾. Infections of moderate to heavy intensity are mainly responsible for the morbidity and health consequences due to STH⁽¹⁴⁾. Therefore, scores for the intensity of infections were given to each STH species (light = 1, mild = 2 and heavy = 3) and infections with worm score ≥ 5 were included in the analysis.

Nutritional status assessment

All subjects underwent physical examination including anthropometric measurements for weight and height. The children were weighed without shoes using a calibrated scale (Seca 709; Seca, Hamburg, Germany); height was measured by standing the child against a vertical wall, and marking off on the wall with the aid of a clipboard. Weight-for-age, height-for-age and weight-for-height Z-scores were used as indicators for underweight, stunting (chronic malnutrition) and wasting (acute malnutrition), respectively. The Z-scores

were calculated using reference data from the National Center for Health Statistics (NCHS) and the WHO in EPI-Info for windows 2002 software (Centers for Disease Control and Prevention, Atlanta, GA, USA). Children who had a *Z*-score below -2 standard deviations of the NCHS reference population median values were considered to be significantly malnourished.

Haematological and biochemical analysis

About 2–3 ml of venous blood were collected from each subject into a plain tube for biochemical analysis. Hb concentrations were measured directly after blood withdrawal using the HemoCue hemoglobinometer (HemoCue, AB, Angelholm, Sweden). Serum ferritin levels were analysed by means of the ADVIA Centaur Analyzer (Siemens Medical Solutions Diagnostics, Deerfield, IL, USA), and children with concentrations of less than 10 $\mu\text{g/l}$ were considered to have deficient Fe stores. Meanwhile, serum Fe and total Fe-binding capacity (TIBC) were determined colorimetrically and then the percentage transferrin saturation was calculated from the ratio of serum Fe concentration:TIBC⁽¹⁵⁾. For quality control, 20% of the samples were randomly selected and examined in duplicate. Children were identified to have IDA if they were anaemic and had low serum ferritin and/or low serum Fe, percentage transferrin saturation and high TIBC^(16,17).

Cognitive function assessment

IQ, also called cognitive function, was assessed using a Test of Nonverbal Intelligence, 3rd edition (TONI-3) which is a language-free, cognitive ability measurement tool for measuring abstract and figural problem solving, a major aspect of intelligence, in subjects ranging in age from 6 years 0 months to 89 years 11 months. The TONI kit has been used in assessing the cognitive function of rural children in Malaysia⁽¹¹⁾. The kit was administered to schoolchildren individually in a comfortable room which was well lit and free from noise. Tables provided in the manual were used to convert the raw scores into the corresponding IQ deviation quotients, which were then categorised into one of seven categories ranging from very poor to very superior, as follows: >130 (very superior); 121–130 (superior); 111–120 (above average); 90–110 (average); 80–89 (below average); 70–79 (poor); and <70 (very poor).

Educational achievement assessment

The assessment of educational achievement was based on the numerical marks of the end-of-semester examinations for three selected subjects, mathematics, English language and Malay language. The marks were collected from the records provided by the headmaster's office. These subjects represent the essential academic components which the student must pass in order to enter higher levels of education, and the Malay language was the medium of instruction for all subjects. Previous studies have used mathematics, English and national languages to assess the educational achievement of urban

and rural schoolchildren in Malaysia and abroad^(4,11,18). For individual selected school subjects and overall results, the highest possible scores were 100 and 300, respectively. The test scores considered as sufficient achievement based on each subject (mathematics, English language and Malay language) and the total scores of the three subjects (TS) were ≥ 50 or ≥ 150 , respectively.

Statistics

Statistical analysis of data was done using SPSS (version 13, September 2004; SPSS, Inc., Chicago, IL, USA). The distribution of IQ and TS measurements was examined using the Shapiro–Wilk test and found to be normal. For descriptive analysis, proportion was used to present the prevalence of categorical variables while mean or median was used to present the quantitative data. The independent *t* test was used to compare the mean IQ and total educational achievement scores between groups. A significance level of 0.05 was used for all statistical tests. However, in order to retain all possible significant associations⁽¹⁹⁾, variables that showed association with a *P* value ≤ 0.20 were used to develop a multiple linear regression model 'STEPWISE' to identify the determinants of cognitive function and educational achievement.

Results

A total of 292 schoolchildren agreed voluntarily to participate in the present study and delivered stool samples for examination. Of these children, only 241 children came for an interview to fill in the questionnaires and underwent a physical examination including anthropometry and cognitive function assessment. Therefore, the statistical analyses for the associations between the variables were based on this sample size.

General characteristics of subjects

The socio-economic and health characteristics of the subjects are shown in Table 1. They comprised 292 children (145 males; 147 females) aged between 7 and 12 years with a median age of 10 years (interquartile range 8–11 years). In general, poverty prevails in this area and about one-third of the fathers had formal education of at least 6 years while only 22.4% of the mothers had similar formal education. During the visits to the villages, we observed the poor personal hygiene of the children; they swim in the rivers very close to defecation sites, play bare-footed, consume unwashed fruits picked from the ground, eat without washing their hands and indulge in soil eating (geophagy).

Health and nutritional status

Almost all children (98.6%) were infected either by one or more of the STH species. The overall prevalence of ascariasis, trichuriasis and hookworm infections was 67.8, 95.5 and 13.4%, respectively. Almost one-third (29.8%) of the children had heavy trichuriasis while 22.3% had heavy ascariasis. All hookworm infections in this population were light infections.

Table 1. General characteristics of Aboriginal schoolchildren participants in the study (*n* 292)

Characteristics	Frequency (%)
Age group	
≤ 10 years	69.2
> 10 years	30.8
Sex	
Female	50.3
Socio-economic status	
Fathers' education level (at least 6 years)	34.3
Mothers' education level (at least 6 years)	22.4
Low household income (<RM450)	81.2
Working fathers	51.4
Working mothers	58.1
Large family size (eight or more members)	35.0
Piped water supply	87.4
Nutritional status	
Significant underweight (weight-for-age $Z < -2$)	37.3
Significant stunting (height-for-age $Z < -2$)	43.6
Significant wasting (weight-for-height $Z < -2$)	5.6
Fe-deficiency anaemia*	34.0
Intestinal parasitic infections	
Heavy ascariasis (mean epg > 50 000)	22.3
Heavy trichuriasis (mean epg > 10 000)	29.8
Hookworm infections	13.4
Mixed STH infections	62.8

RM, Malaysian ringgit; epg, eggs per gram of stool; STH, soil-transmitted helminthes.
 * Fe-deficiency anaemia: Hb concentration < 120 g/l, low serum ferritin (<10 g/l) and/or low serum Fe (<10.6 mmol/l), high total Fe-binding capacity (>75 μmol/l) and low transferrin saturation (<16%).

The prevalence of mild and significant underweight was 52.3 and 37.3%, respectively, while the prevalence of mild stunting and wasting was 43.6 and 43.1%, respectively, and of significant stunting and wasting was 43.6 and 5.6%, respectively. Overall, 117 (48.5%) children were anaemic (Hb < 120 g/l) and the prevalence of IDA was 34%, which accounted for 70.1% of the anaemia cases among these subjects. Data on the prevalence of intestinal parasitic infections and IDA among these subjects have been published previously^(20,21).

Cognitive function and educational achievement

The mean IQ scores of these children was 76.4 (SD 5.4), which is considered as poor. The majority (67.6%) of the subjects were in the category of poor while 24.1 and 8.1% of them were in the category of below average and very poor, respectively. However, none of the subjects was distributed in the grades of above average, superior or very superior.

The mean educational score of each subject was low and most of the children (72.6%) had insufficient TS (Table 2). Moreover, 56.8% of the children had sufficient achievement for Malay language (mean score 50.9), whereas only 32.4 and 24.9% of them had sufficient scores for English language and mathematics, respectively.

Determinants of low cognitive function and low educational achievement

The differences in mean scores of IQ and TS in relation to some socio-economic, health, nutritional and demographic factors were examined and the results are shown in Table 3. Regarding the IQ (indicator of low cognitive function), children living in poverty (income < 450 Malaysian ringgit (RM)) had significantly lower IQ scores ($t = 3.041$; $P = 0.007$) than children from households with higher incomes. Similarly, the children of mothers with low educational levels had significantly lower IQ scores ($t = 3.256$; $P = 0.001$) than the children of mothers with at least 6 years of formal education. Children suffering IDA also had significantly lower IQ scores than non-anaemic children ($t = 2.140$; $P = 0.033$).

With TS (indicator of low educational achievement), children infected with STH of moderate to heavy intensities (worm scores ≥ 5) had significantly lower TS scores ($t = 1.990$; $P = 0.030$) than non-infected children or those with light infections. Similarly, children of families with a low household income ($t = 2.251$; $P = 0.025$) and also those who were suffering from IDA ($t = 2.460$; $P = 0.007$) had significantly lower TS scores than children of families with a monthly household income > RM450 and those without IDA.

Stepwise multiple linear regression analyses were performed to identify the significant determinants of children's IQ and TS with control for the effects of possible confounders. The output of the models showed that IQ score was found to be significantly influenced by low household income which contributed the most to the regression variance ($r^2 = 0.059$; $P = 0.020$), followed by low mothers' educational level ($r^2 = 0.042$; $P = 0.043$), while the non-significant variables ($P > 0.05$) were removed in a stepwise manner. In another model, the determinants of TS were identified after adding IQ score to the independent variables and controlling the effect of age; IDA was retained as the significant determinant of educational achievement ($r^2 = 0.025$; $P = 0.015$).

Table 2. Educational achievement test scores of school subjects among Aboriginal schoolchildren (*n* 241)

(Mean values, standard deviations and ranges)

Subject	Score			Percentage with test score ≥ 50 , or ≥ 150 for TS
	Mean	SD	Range	
ML	50.9	8.7	30–75	56.8
EL	45.9	7.5	24–61	32.4
MT	43.9	7.2	22–63	24.9
TS*	140.6	14.9	107–184	27.4

ML, Malay language; EL, English language; MT, mathematics; TS, total score.

* TS = ML + EL + MT.

Table 3. Univariate and multivariate analyses for scores of intelligence quotient (IQ) and total scores of educational achievement (TS) by socio-economic, health and nutritional factors among Aboriginal schoolchildren (*n* 241)‡

(Mean values and standard deviations)

Factors	IQ			TS		
	Mean	SD	<i>t</i>	Mean	SD	<i>t</i>
Sex			0.603			-1.139
Male	76.2	5.4		141.7	14.3	
Female	76.6	5.5		139.6	15.6	
Age group			0.361			1.248
≤ 10 years	76.3	5.4		142.5	15.5	
> 10 years	76.6	5.5		138.6	13.0	
Household income			3.041*†			2.251*
≥ RM450	78.5	5.3		145.0	15.1	
< RM450	72.9	5.0		139.6	14.2	
Fathers' education			1.260			0.339
≥ 6 years formal education	77.5	5.7		140.9	15.5	
No formal education	76.5	5.4		140.2	14.8	
Mothers' education			3.256*†			0.277
≥ 6 years formal education	78.7	5.9		142.2	14.2	
No formal education	72.1	4.8		140.5	13.8	
Mothers' employment status			1.691			1.125
Working	75.9	5.2		139.7	14.5	
Not working	77.1	5.6		141.9	15.6	
Family size			1.047			1.670
Eight or more members (large)	75.8	5.2		138.1	11.0	
Less than eight members	76.7	5.5		141.7	12.9	
STH infections (worm score ≥ 5)			-0.340			1.990*
Moderate to heavy	75.3	4.9		137.7	12.2	
Negative to light	76.5	6.5		142.8	13.8	
Fe status			2.140*			2.460*†
IDA	71.8	4.2		135.4	10.5	
Non-anaemic	76.9	5.6		142.3	13.3	
Weight status			0.820			0.193
Underweight	76.8	5.2		140.9	13.1	
Non-underweight	76.2	5.5		140.5	12.3	
Height status			0.144			0.771
Stunted	76.4	5.4		139.8	13.5	
Non-stunted	76.5	5.4		141.3	12.6	

RM, Malaysian ringgit; STH, soil-transmitted helminthes; IDA, Fe-deficiency anaemia.

* Significant association ($P < 0.05$; independent *t* test).† Confirmed as a significant determinant by multiple linear regression ($P < 0.05$).

‡ Variables included in the multiple linear regression model were IDA, STH infections (worm score ≥ 5), household income, family size and IQ score.

Discussion

The present study showed that the IQ grade of the subjects was poor and more than two-thirds of them had insufficient educational achievement. These findings are very similar to the results obtained by a previous study among Aboriginal schoolchildren in Perak, Malaysia⁽¹¹⁾. On the other hand, the subjects of the present study showed lower cognitive function, represented by poor IQ, compared with children in other rural areas in Sarawak, East Malaysia⁽²²⁾. Similarly, we found that the educational achievement of these children was much lower than rural children in Sarawak and those in urban areas^(12,22). Studies from other countries have reported poor IQ and educational levels among rural schoolchildren^(4,23). These findings were not surprising as the rural areas usually lack well-equipped schools, expert teachers and sufficient infrastructure, including electricity. However, we observed that the school is well equipped with different facilities including a computer laboratory with Internet access, a library, and

visual and science laboratories. It is generally accepted that poor socio-economic profile, parasitic infections, malnutrition, non-delivery of educational assistance, contrast in the pedagogy and the culture in Aboriginal communities could propagate the weaknesses in cognitive function and educational achievement.

Regarding the potential determinants of cognitive function and educational achievement, our findings were in accordance with previous studies from other countries that poverty, low educational level of parents⁽⁴⁾, malnutrition including micronutrient deficiencies^(24,25) and intestinal parasitic infections^(23,26) are major risk factors of insufficient cognitive and educational performance. We found higher IQ scores among children whose mothers had higher educational levels and whose families had higher monthly incomes. These findings are consistent with a previous study on urban primary schoolchildren in Selangor, Malaysia⁽¹⁰⁾. It is well known that educated parents play a crucial role in the educational process of their children. Poverty also limits educational achievement

in different ways, with children of poor families being forced to work, which discourages their school attendance and performance.

Beside these factors, we found that children suffering from IDA were more prone to have lower cognitive function and educational achievement than non-anaemic children. Throughout previous reports, IDA has been conclusively seen to delay psychomotor development and impair the cognitive function of preschool and school-age children in Thailand⁽⁴⁾, Indonesia⁽²⁷⁾ and Zaire⁽²⁴⁾. Moreover, a previous study found that schoolchildren from a non-institutionalised US population with Fe deficiency with or without anaemia had twice the risk of scoring insufficient mathematics scores than children with normal Fe status⁽²⁵⁾. Furthermore, Fe supplements given to anaemic children and adults have been shown to produce significant improvements in their cognitive function and educational achievement^(3,28–30). Double-blind clinical trials aimed at assessing the effects of Fe deficiency on attention and learning processes in preschool and school-aged children found that IDA produces alterations in cognitive processes related to visual attention and concept acquisition and that these alterations are reversed with Fe treatment^(27,31). In contrast, previous studies reported that the poor scores in school subjects and cognitive functions tests of schoolchildren with IDA were not reversed by Fe supplementation^(32,33).

Concern has been expressed about the potential impact of IDA on the cognitive function and psychomotor development of children throughout the neonatal, postnatal, infancy and childhood periods. A long-term study aimed at investigating the association of maternal Fe status by measuring umbilical cord serum ferritin concentrations and applying test scores of psychomotor development and functioning at 5 years of age showed that the children who had the lowest ferritin concentrations had significantly lower language ability and tractability than children who had higher ferritin concentrations⁽³⁴⁾. In the same vein, a previous study found that children who had moderate anaemia as infants achieved lower IQ scores upon entry to school than did children who were non-anaemic during infancy⁽³⁵⁾. These studies revealed that the effects of maternal Fe deficiency are not restricted to prematurity and impaired fetal growth, but also extend to abnormal psychomotor development and impaired cognitive function of these children in later life. Several mechanisms may explain the potential causal association between Fe deficiency and impaired cognitive and educational performance. Fe deficiency results in decreased body Fe stores, including decreased Fe in the central nervous system, even before erythrocyte production is affected⁽³⁶⁾. In experimental animals, Fe has been shown to play a key role in brain function. Several areas of the brain contain Fe, sometimes in large quantities⁽³⁷⁾, and Fe-deficient animals show alterations both in neurotransmitters and behaviour that do not usually respond to Fe replenishment⁽³⁸⁾.

Meanwhile, the present study showed a lower educational achievement among children who were infected with STH of moderate to heavy intensities compared with those who were not infected or who harboured light STH infections.

The evidence that worms may affect cognitive function is largely circumstantial. More specifically, the clinical consequences of moderate and heavy infections with *T. trichiura* include stunting and IDA^(16,39), both of which are associated with impaired cognitive function and learning ability^(25,27). However, no significant association was reported between IDA and these infections among the subjects of the present study⁽²¹⁾. A previous double-blind placebo-controlled trial among Jamaican schoolchildren reported that moderate to high loads of *T. trichiura* infection have an adverse effect on certain cognitive functions but that this is reversible by therapy⁽²³⁾. The mechanism by which parasitic infections impaired cognitive function and educational achievement could be directly through mediators in the systemic circulation including cellular and humoral immune responses to infections⁽²³⁾ or could be due to the fact that children infected with helminthes are lethargic and weak, and have frequent attacks of diarrhoea or dysentery and abdominal pain which may disturb the child's attention in the classroom.

Although the present study is the first, to our knowledge, to investigate the potential determinants influencing the cognitive function and educational achievement of Aboriginal schoolchildren in Peninsular Malaysia, some limitations related to study design (cross-sectional) should be considered when interpreting the study findings. Therefore, the present study demonstrated the association of cognitive and educational performance with socio-economic status and IDA but did not explain causality. Other potentially important determinants such as school-level factors (a viable curriculum, teachers' motivation, and a conducive learning environment) and family-level factors (home environment and parents' motivation) were not addressed in the present study.

In conclusion, the present study provides a community-based figure for the cognitive and educational performance of Aboriginal schoolchildren, which was found to be poor and influenced by household income, maternal education and IDA. It is recommended that the School Health Service in Malaysia should also monitor the health and growth of primary schoolchildren. Effective and integrated measures to improve the socio-economic, nutritional and health status of Aboriginal and rural children would have a pronounced positive effect on their education.

Acknowledgements

The authors would like to acknowledge the headmaster, teachers and schoolchildren of Sekolah Kebangsaan Betau, the staff of the Department of Orang Asli Affairs, Ministry of Rural Development, and the staff of Clinic Pos Betau for their fruitful help. The present study was funded by the University of Malaya, Kuala Lumpur, Malaysia (research grant code PS178/2007B).

The contribution of the authors to the present study was: H. M. A., J. S., A. A. S. and W. A. A. were involved in the study design, analysis of data and writing of the manuscript; H. M. A., M. A. M., A. M. A. and A. A. A. collected the data. All authors read and approved the final manuscript.

There are no conflicts of interest.

References

- Lorntz B, Soares AM, Moore SR, *et al.* (2006) Early childhood diarrhea predicts impaired school performance. *Pediatr Infect Dis J* **25**, 513–520.
- Niehaus MD, Moore SR, Patrick PD, *et al.* (2002) Early childhood diarrhea is associated with diminished cognitive function 4 to 7 years later in children in a northeast Brazilian shantytown. *Am J Trop Med Hyg* **66**, 590–593.
- Stoltzfus R, Kvalsvig J, Chwaya H, *et al.* (2001) Effects of iron supplementation and anthelmintic treatment on motor and language development of preschool children in Zanzibar: double blind, placebo controlled study. *BMJ* **323**, 1389–1393.
- Sunghong R, Mo-suwan L & Chongsuvivatwong V (2002) Effects of haemoglobin and serum ferritin on cognitive function in school children. *Asia Pac J Clin Nutr* **7**, 245–255.
- Nokes C & Bundy DA (1994) Does helminth infection affect mental processing and academic achievement? *Parasitol Today* **10**, 14–18.
- Roberts JE, Burchinal MR & Zeisel SA (2002) Otitis media in early childhood in relation to children's school-age language and academic skills. *Pediatrics* **110**, 696–706.
- Bourgeois BF (2004) Determining the effects of antiepileptic drugs on cognitive function in pediatric patients with epilepsy. *J Child Neurol* **19**, Suppl. 1, S15–S24.
- Thorsdottir I (2005) Supplement and stimulation for stunted children. *Lancet* **366**, 1756–1758.
- Walker SP, Chang SM, Powell CA, *et al.* (2005) Effects of early childhood psychosocial stimulation and nutritional supplementation on cognition and education in growth-stunted Jamaican children: prospective cohort study. *Lancet* **366**, 1804–1807.
- Zaini MZ, Lim CT, Low WY, *et al.* (2005) Effects of nutritional status on academic performance of Malaysian primary school children. *Asia Pac J Public Health* **17**, 81–87.
- Zaleha MI, Iskandar ZA, Khalid AK, *et al.* (2000) Effect of iodized oil supplementation on thyroid hormone levels and mental performance among Orang Asli schoolchildren and pregnant mothers in an endemic goiter area in Peninsular Malaysia. *Asia Pac J Clin Nutr* **9**, 274–281.
- Martin LK & Beaver PC (1968) Evaluation of Kato thick smears technique for quantitative diagnosis of helminth infections. *Am J Trop Med Hyg* **17**, 382–391.
- World Health Organization (2002) *Prevention and Control of Intestinal Parasitic Infections. Prevention and Control of Schistosomiasis and Soil-transmitted Helminthiasis*. WHO Technical Report Series no. 912. Geneva: WHO.
- Albonico M, Montresor A, Crompton DWT, *et al.* (2006) Intervention for the control of soil-transmitted helminthiasis in the community. *Adv Parasitol* **61**, 311–348.
- Blanck MH, Pfeiffer CM, Caudill SP, *et al.* (2003) Serum iron and iron-binding capacity: a round-robin interlaboratory comparison study. *Clin Chem* **49**, 1672–1675.
- Nor Aini U, Al-Mekhlafi HM, Azlin M, *et al.* (2007) Serum iron status in Orang Asli children living in endemic areas of soil-transmitted helminthes. *Asia Pac J Clin Nutr* **16**, 724–730.
- World Health Organization (2001) *Iron Deficiency Anaemia: Assessment, Prevention, and Control*. Geneva: WHO.
- Zalilah MS, Jenny TB & Nan EJ (2000) Nutrition and educational achievement of urban primary schoolchildren in Malaysia. *Asia Pac J Clin Nutr* **9**, 264–273.
- Bendel RB & Afifi AA (1977) Comparison of stopping rules in forward regression. *J Am Stat Assoc* **72**, 46–53.
- Al-Mekhlafi HM, Atiya AS, Lim YAL, *et al.* (2007) An unceasing problem: soil-transmitted helminthiasis in rural Malaysian communities. *Southeast Asian J Trop Med Public Health* **38**, 998–1007.
- Al-Mekhlafi HM, Surin J, Atiya AS, *et al.* (2008) Anaemia and iron deficiency anaemia among Aboriginal schoolchildren in rural Peninsular Malaysia: an update on a continuing problem. *Trans R Soc Trop Med Hyg* **102**, 1046–1052.
- Zaleha MI, Hayati AR, Rizam AR, *et al.* (2003) Micronutrients and its correlation with mental performance among schoolchildren in Bario, Sarawak: a preliminary study. *Med J Malaysia* **58**, 309–319.
- Nokes C, Grantham-McGregor SM, Sawyer AW, *et al.* (1992) Moderate to heavy infections of *Trichuris trichiura* affect cognitive function in Jamaican school children. *Parasitology* **104**, 539–547.
- Boivin MJ & Giordani B (1993) Improvements in cognitive performance for schoolchildren in Zaire, Africa, following an iron supplement and treatment for intestinal parasites. *J Pediatr Psychol* **18**, 249–264.
- Halterman JS, Kaczorowski JM, Aligne CA, *et al.* (2001) Iron deficiency and cognitive achievement among school-aged children and adolescents in the United States. *Pediatrics* **107**, 1381–1386.
- Ezeamama AE, Friedman JF, Acosta LP, *et al.* (2005) Helminth infection and cognitive impairment among Filipino children. *Am J Trop Med Hyg* **72**, 540–548.
- Soewondo S, Husaini MA & Pollitt E (1989) Effects of iron deficiency on attention and learning processes in pre-school children: Bandung, Indonesia. *Am J Clin Nutr* **50**, 667–674.
- Aukett MA, Parks YA, Scott PH, *et al.* (1986) Treatment with iron increases weight gain and psychomotor development. *Arch Dis Child* **16**, 849–857.
- Soemantri AG, Pollitt E & Kim I (1985) Iron deficiency anemia and educational achievement. *Am J Clin Nutr* **42**, 1221–1228.
- Murray-Kolb LE & Beard JL (2007) Iron treatment normalizes cognitive functioning in young women. *Am J Clin Nutr* **85**, 778–787.
- Seshadri S & Gopaldas T (1989) Impact of iron supplementation on cognitive functions in preschool and school-aged children: the Indian experience. *Am J Clin Nutr* **50**, 675–686.
- Harahap H, Jahari AB, Husaini MA, *et al.* (2000) Effects of an energy and micronutrient supplement on iron deficiency anaemia, physical activity, and motor and mental development in undernourished children in Indonesia. *Eur J Clin Nutr* **54**, Suppl. 2, S114–S119.
- Pollitt E, Hathirat P, Kotchabhakdi NJ, *et al.* (1989) Iron deficiency and educational achievement in Thailand. *Am J Clin Nutr* **50**, 687–697.
- Tamura T, Goldenberg RL, Hou J, *et al.* (2002) Cord serum ferritin concentrations and mental and psychomotor development of children at five years of age. *J Pediatr* **140**, 165–170.
- Lozoff B, Jimenez E & Wolf AW (1991) Long-term developmental outcome of infants with iron deficiency. *N Engl J Med* **325**, 687–694.
- Pollitt E & Leibel RL (1976) Iron deficiency and behavior. *J Pediatr* **88**, 372–381.
- Dwork AJ, Lawler G, Zybert PA, *et al.* (1990) An autoradiographic study of the uptake and distribution of iron by the brain of the young rat. *Brain Res* **518**, 31–39.
- Beard JL (2001) Iron biology in immune function, muscle metabolism and neuronal functioning. *J Nutr* **131**, Suppl. 2, S568–S579.
- Cooper ES, Bundy DA, MacDonald TT, *et al.* (1990) Growth suppression in the *Trichuris* dysentery syndrome. *Eur J Clin Nutr* **44**, 138–147.