



University of Dundee

Clinical assessment to screen for the detection of oral cavity cancer and potentially malignant disorders in apparently healthy adults

Walsh, Tanya; Liu, Joseph L. Y.; Brocklehurst, Paul; Lingen, Mark; Kerr, Alexander R.; Ogden, Graham; Warnakulasuriya, Saman; Scully, Crispian

Published in:
Cochrane Database of Systematic Reviews

DOI:
[10.1002/14651858.CD010173](https://doi.org/10.1002/14651858.CD010173)

Publication date:
2012

Document Version
Publisher's PDF, also known as Version of record

[Link to publication in Discovery Research Portal](#)

Citation for published version (APA):

Walsh, T., Liu, J. L. Y., Brocklehurst, P., Lingen, M., Kerr, A. R., Ogden, G., ... Scully, C. (2012). Clinical assessment to screen for the detection of oral cavity cancer and potentially malignant disorders in apparently healthy adults. *Cochrane Database of Systematic Reviews*, (11), [CD010173]. [10.1002/14651858.CD010173](https://doi.org/10.1002/14651858.CD010173)

General rights

Copyright and moral rights for the publications made accessible in Discovery Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from Discovery Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain.
- You may freely distribute the URL identifying the publication in the public portal.

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Clinical assessment to screen for the detection of oral cavity cancer and potentially malignant disorders in apparently healthy adults (Protocol)

Walsh T, Liu JLY, Brocklehurst P, Lingen M, Kerr AR, Ogden G, Warnakulasuriya S, Scully C



**THE COCHRANE
COLLABORATION®**

This is a reprint of a Cochrane protocol, prepared and maintained by The Cochrane Collaboration and published in *The Cochrane Library* 2012, Issue 11

<http://www.thecochranelibrary.com>



Clinical assessment to screen for the detection of oral cavity cancer and potentially malignant disorders in apparently healthy adults (Protocol)

Copyright © 2012 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.

TABLE OF CONTENTS

HEADER	1
ABSTRACT	1
BACKGROUND	2
OBJECTIVES	4
METHODS	5
ACKNOWLEDGEMENTS	7
REFERENCES	8
ADDITIONAL TABLES	10
APPENDICES	15
HISTORY	16
CONTRIBUTIONS OF AUTHORS	17
DECLARATIONS OF INTEREST	17
SOURCES OF SUPPORT	17

[Diagnostic Test Accuracy Protocol]

Clinical assessment to screen for the detection of oral cavity cancer and potentially malignant disorders in apparently healthy adults

Tanya Walsh¹, Joseph LY Liu², Paul Brocklehurst¹, Mark Lingem³, Alexander R Kerr⁴, Graham Ogden⁵, Saman Warnakulasuriya⁶, Crispian Scully⁷

¹School of Dentistry, The University of Manchester, Manchester, UK. ²University of Dundee, Dental Health Services Research Unit, Scottish Dental Clinical Effectiveness Programme, NHS Education for Scotland, Dundee, UK. ³Pritzker School of Medicine, Division of Biological Sciences, Department of Pathology, University of Chicago, Chicago, Illinois, USA. ⁴Department of Oral and Maxillofacial Pathology, Radiology and Medicine, New York University College of Dentistry, New York, USA. ⁵Unit of Oral Surgery & Medicine, School of Dentistry, University of Dundee, Dundee, UK. ⁶Department of Oral Medicine, King's College London, London, UK. ⁷Eastman Dental Institute for Oral Health Care Sciences, University College London, London, UK

Contact address: Tanya Walsh, School of Dentistry, The University of Manchester, Coupland III Building, Oxford Road, Manchester, M13 9PL, UK. tanya.walsh@manchester.ac.uk.

Editorial group: Cochrane Oral Health Group.

Publication status and date: New, published in Issue 11, 2012.

Citation: Walsh T, Liu JLY, Brocklehurst P, Lingem M, Kerr AR, Ogden G, Warnakulasuriya S, Scully C. Clinical assessment to screen for the detection of oral cavity cancer and potentially malignant disorders in apparently healthy adults. *Cochrane Database of Systematic Reviews* 2012, Issue 11. Art. No.: CD010173. DOI: 10.1002/14651858.CD010173.

Copyright © 2012 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.

ABSTRACT

This is the protocol for a review and there is no abstract. The objectives are as follows:

The objective of this review is to estimate the accuracy of the conventional oral examination (COE) used singly or in combination with another index test (Additional [Table 1](#)) as a screening test for the detection of oral cancer and potentially malignant disorders (PMD) of the lip and oral cavity of apparently healthy adults.

The secondary objective of this review is to estimate the accuracy of the different tests with COE, when compared with each other.

We will use meta-regression to explore possible sources of heterogeneity, ways in which the observed diagnostic test accuracy varies according to particular characteristics. Covariates to be included in these analyses will include.

- Characteristics of the study sample: prevalence of carcinoma or PMD disease in the study (> 50% prevalence), inclusion of HPV + adults, tobacco users / high alcohol consumption.
- Target condition (oral squamous cell carcinoma alone or oral squamous cell carcinoma and PMD).
- Attributes of the screening programme: prospective organised or opportunistic, type of reference standard (examination and clinical evaluation by physician with specialist knowledge or extended follow-up), operator (dental or general practice professionals or other healthcare workers).

BACKGROUND

Target condition being diagnosed

The target conditions of interest are oral squamous cell carcinoma (OSCC), the most common form of oral cavity cancer (Scully 2000a), and potentially malignant disorders (PMD), of the lip and oral cavity in apparently healthy adults. At a meeting of international oral cancer and precancer experts held in 2005, the concept of precancer, along with issues surrounding classification and definition, aetiology, diagnosis and management was extensively discussed. Through consensus, the term 'potentially malignant disorders' was selected to convey the fact that not all precancerous lesions and conditions will transform to cancer, but rather that there is the potential for malignant transformation (van der Waal 2009; Warnakulasuriya 2007).

The natural history of oral cancer is not fully understood, given variations in disease processes and dysplastic changes in PMD (Napier 2008; Scully 2009). Most oral carcinomas are preceded by PMD of which erythroplakia, non-homogeneous leukoplakia, erosive lichen planus, oral submucous fibrosis and actinic keratosis are perhaps the most important (Warnakulasuriya 2007). The concept of a two-step process of cancer development of the oral mucosa (i.e. precursor to established lesion) is established. Oral leukoplakia is the best-known precursor lesion and between < 1% and 18% develop into oral cancer. The original 1978 World Health Organization (WHO) definition of oral leukoplakia has been revised to read "The term leukoplakia should be used to recognize white plaques of questionable risk having excluded (other) known diseases or disorders that carry no increased risk for cancer" (Warnakulasuriya 2007). The presence of epithelial dysplasia can help predict malignant development but the diagnosis is essentially subjective, with not all lesions exhibiting dysplasia, some becoming malignant and some regressing. Carcinoma can also develop from lesions in which epithelial dysplasia was not previously diagnosed. Numerous attempts have been made to relate biological characteristics to the malignant potential of leukoplakias, but finding a definitive characteristic remains elusive (Reibel 2003). Most authorities regard leukoplakia as a dynamic rather than a static process, in terms of its progression and the development of malignancy (Napier 2008). For example, Jaber et al (Jaber 2003) followed up 630 patients who attended oral medicine clinics in London and Bristol between 1972 and 1996. The majority of oral leukoplakia (43.8%) had features of mild dysplasia, 30% moderate and 24.7% severe oral epithelial dysplasia. Whilst this is strongly suggestive that dysplasia is typically associated with oral leukoplakia, it can also be seen in patients with reactive epithelial changes such as candidosis, viral infections, lichen planus, and denture-induced hyperplasia. The study concluded that predicting the severity of histopathological change from clinical examination alone remains difficult. Estimates of malignant transformation rates (MTR) vary enormously, from site to site within the mouth, from population to population and from study to study (Napier 2008). The MTR

of hospital-based surveys are consistently higher than community-based studies because of sampling bias. Petti (Petti 2003) calculated a global MTR of oral leukoplakia of 1.36% per year (95% confidence interval (CI) 0.69% to 2.03%) based on the prevalence of leukoplakia, but this far exceeds the numbers of actual cases of malignancy. Virtually all studies emphasize the chronicity of oral PMD, with an increasing tendency to malignant change in the first 5 years. For example, the incidence of OSCC arising from leukoplakia in Californians was greatest in the second year of follow-up (11 out of 45; 24%) (Silverman 1984). The proportion of PMD that will develop OSCC is uncertain but low; best estimates suggest a rate of less than 2% per annum (Napier 2008).

The early detection and excision of some PMD can prevent malignancy, or if malignancy is detected, there is some evidence that appropriate treatment can reduce disease severity and improve survival rates (Brocklehurst 2010b; van der Waal 2009; Warnakulasuriya 2007). Leukoplakias can be treated by a number of methods. According to Lodi et al's systematic review (Lodi 2006), the effectiveness of surgical interventions, including laser therapy and cryotherapy, has not been studied by means of a randomised controlled trial (RCT) with a no treatment/placebo arm. Vitamin A and retinoids have been tested by five RCTs, two studies investigated beta carotene or carotenoids, the other drugs tested were bleomycin (one study), mixed tea (one study) and ketorolac (one study). None of the treatments tested showed a benefit when compared with the placebo. Lodi et al concluded that there was no evidence of effective treatment in preventing the malignant transformation of leukoplakia. Where resolution of a lesion is observed, relapses and adverse effects are common.

Technologies to treat and manage oral cancer have progressed substantially, as shown by systematic reviews of RCT of interventions, e.g. Bessell 2011; Furness 2011; Glennly 2010. Once progressed to frank malignancy, traditional treatment is surgery and radiotherapy. More recently, systemic chemotherapy has been included as part of the treatment regimen before or during radiotherapy. Surgery for the treatment of oral cancer is followed by exacting reconstructive surgery to restore form and function. Debilitating side effects can occur as a result of radiotherapy and chemotherapy, adversely affecting an individual's quality of life. The 5-year survival following diagnosis has remained at around 50% for the past 30 years in most countries (Parkin 2001; Warnakulasuriya 2009). This is in marked contrast to the improved survival rates in many other cancers, such as those of the breast and the colon (Cancer Research UK), but may be explained at least in some part by the fact that oral cancer is more often diagnosed at a late stage of the disease, when prognosis is poorer and the risks of significant morbidity and mortality are substantially higher (Rogers 2009; Rusthoven 2010).

Index test(s)

The standard screening of apparently healthy adults for oral cancer and PMD is a systematic and thorough visual inspection and examination of the oral mucosa and palpation of the neck under normal (incandescent) light. In most instances this is carried out by a frontline clinician such as a general dentist. This conventional visual and tactile oral examination (COE) used as a screen can be conducted with the minimum of effort and distress to the individual (Additional Table 1). Screening can be carried out opportunistically, for instance when an individual presents to their dentist for a check-up, or as part of an organised screening programme. The COE is usually followed by referral for further investigation if this is deemed necessary. The form of further investigation is variable nationally or internationally, with different investigational pathways. For instance this could take the form of examination by an oral medicine specialist or specialist oral surgeon at a specialist clinic or hospital facility.

Reviews of primary studies of diagnostic test accuracy in this area have identified a number of index tests which could be used as adjuncts to the COE to improve earlier detection of oral cavity cancer and PMD (Fedele 2009; Leston 2010; Lingen 2008; Patton 2008; Rethman 2010). These include:

- vital staining (Toluidine Blue, Tolonium chloride)
- light-based detection (such as ViziLite and ViziLite Plus, Microlux/DL, VELscope, Orascoptic DK, Identafi 3000)
- blood and saliva analysis.

Vital staining and oral cytology are long available adjuncts to a conventional oral examination (Leston 2010; Lingen 2008). Other tests such as light-based detection systems have become commercially available only more recently. Blood analysis and saliva analysis are more novel tests at an early stage of evaluation.

Of the index tests listed above, vital staining, light-based detection and blood and saliva analysis could be used as screening adjuncts to the COE (Additional Table 1). In this review, we will restrict vital staining index tests to those applied in a rinse form. A companion review (Diagnostic tests for oral cancer and potentially malignant disorders in patients presenting with clinically evident lesions) will restrict vital staining index tests to those applied to a lesion that has been visualized. Where access to general dental practitioners or general practitioners is limited, either as a result of geographical location or barriers to uptake of healthcare provision, screening using the index tests listed above could, in principle, be undertaken by trained healthcare workers.

It is worth noting there are regional differences in regulations on the use of some of the above tests. For example, Toluidine Blue having been consistently rejected as a stand alone technique, is not cleared to be a stand alone screening technique in the United States. It is included in the ViziLite Plus system. However, the Toluidine Blue only component is approved by the US Food and Drug Administration (FDA) as a marking device.

Of the index tests listed above, all have the potential to be used as adjuncts to the COE (Additional Table 1) by healthcare workers or clinicians undertaking screening in the general population. Adding

any one of the proposed index tests to the COE, the tests could have a triage role in detecting lesions of uncertain significance with referral where appropriate. For instance, traumatic keratoses are common, and referring each patient with a white patch to a specialist to undergo a scalpel biopsy is excessive, and incurs increased financial cost and patient worry. A non-invasive index test or combination of tests adjunctive to the COE that provides a frontline clinician with a high degree of accuracy would not only reduce the number of patients with benign disease being referred, but could avoid the need for invasive biopsy in patients testing negative.

Rationale

Oral cancer is a significant global health problem with increasing incidence and mortality rates (Ferlay 2010; Warnakulasuriya 2009). Cancer of the lip or oral cavity is a relatively common cancer worldwide, with an estimated 263,000 new cases and 127,000 deaths in 2008, and an increasing incidence in recent years (Ferlay 2010). There is wide geographic variation in disease incidence and mortality, with almost double the incidence in developing countries as in developed countries, and a threefold increase in mortality. Tobacco use, alcohol consumption, betel quid chewing and low socio-economic status have traditionally been thought to be the most important risk factors of oral cancer (Conway 2008; Faggiano 1997; La Vecchia 1997; Macfarlane 1995; Ogden 2005). The human papillomavirus (HPV) is also an important risk factor in oral squamous cell carcinoma and there is some evidence that HPV positive cancers associate with better survival (Ang 2010). Men have had a higher incidence of oral cancer than women (Ferlay 2010), but this disparity can be explained by men having a higher exposure to the above risk factors (Freedman 2007). The gender difference has narrowed in recent decades from a ratio of 5 males to 1 female diagnosed with oral cancers in the 1960s to less than 2 to 1 in 2008 (Ferlay 2010). Although traditionally the risk of oral cancer increases with age, the incidence among younger adults has been increasing in the European Union and the United States (Warnakulasuriya 2009). The 5-year survival rate depends on multiple factors, including patient and tumour characteristics, treatment received and stage at diagnosis. Oral cancer incidence and mortality can be reduced using three approaches: (i) primary prevention, (ii) secondary prevention, screening and early detection, and (iii) improved treatment (Scully 2000b).

Successful early detection of oral cancer or PMD is highly dependent on whether 'at risk' individuals present for screening examination. Early diagnosis relies on the awareness and motivation of the clinician or patient in identifying a suspicious lesion or symptom while it is still at an early stage. Whilst many organisations advocate cancer-related checks, including the American Cancer Society for individuals of all risk groups (American Cancer Society 1992) and the US Preventive Health Services Task Force for high risk individuals (US Preventive Services Task Force), there is much global

variation in the provision and promotion of routine oral cancer examinations. Currently, no national population-based screening programmes for oral cancer have been implemented in the developed countries, although opportunistic screening has been advocated (Brocklehurst 2010a). Consequently, individuals will often present for examination at a later stage of the disease, when the risks of significant morbidity and mortality are substantially higher. A province-wide programme is being evaluated in British Columbia, Canada (Rosin 2006) but the evaluation is ongoing and no final results have been reported to date. Brocklehurst et al's systematic review identified only one RCT using visual examination with a follow-up period of 9 years which was carried out in India. The authors of the review concluded that the current evidence is insufficient to recommend population-based screening and suggested that opportunistic screening of high risk groups may potentially improve outcomes (Brocklehurst 2010a).

There is some debate in the literature on anticipated differences in diagnostic accuracy of prospective population-based invitational screening programmes and a more opportunistic approach (when patients attend their general (dental) practitioner for routine examination or for treatment). In Downer et al's systematic review of test performance in screening for oral cancer and PMD, only prospective investigations of population screening with specified reference standards were included. The pooled sensitivities and specificities were 0.85 (95% CI 0.730 to 0.919) and 0.97 (95% CI 0.930 to 0.982) respectively (Downer 2004). An opportunistic approach that focuses on high risk groups is also possible (McGurk 2010; Sankaranarayanan 1997). A simulation study which used neural network and machine learning techniques suggested opportunistic screening aimed at high risk groups may be both effective and cost effective (Speight 2006). However, the results of the only completed randomised controlled trial undertaken on oral cancer screening as an intervention demonstrated that the yield from both whole population and opportunistic approaches is likely to be important (Brocklehurst 2010a). Many individuals with risk factors may not attend the dentist and are therefore not amenable to an opportunistic approach (Netuveli 2006; Yusof 2006). Furthermore, the evidence suggests that diagnostic/screening tools based on machine learning such as Speight et al's (Speight 2006) can provide useful but preliminary evidence, as they tend to be limited by "the fitting of models that are implausible and the tendency ... to understate misclassification errors" (Liu 2006).

Reviews assessing the test accuracy of a conventional oral examination as a population screening tool (e.g. Downer 2004; Moles 2002) have highlighted methodological flaws in the primary diagnostic test accuracy studies, although explicit methodological quality assessment of these studies using a validated and widely used checklist was not undertaken.

In this review we aim to identify screening tests for oral cancer and PMD to evaluate the diagnostic accuracy of the COE and the accuracy of the other index tests (Additional Table 1) used as adjuncts to the oral examination in apparently healthy adults.

The index tests proposed for evaluation in this review are suitable for use in the community or as part of a dental examination in a general dental practitioners' office. The review will include both prospective investigations of organised screening programmes and prospective opportunistic screening. It is important that this review consider both, as opportunistic screening of patients attending in a general practice setting are self selecting and may not be representative of the population of interest. In either scenario, screening may be carried out by dental professionals or healthcare workers. The purpose of the screening is to identify the presence or absence of PMDs which require referral to secondary care for definitive diagnosis and possibly treatment. The proposed index tests cannot confirm whether a PMD is cancerous before deciding on referral to secondary care; biopsy with histopathology is currently the only confirmatory method of oral cancer diagnosis.

The Cochrane Oral Health Group has undertaken a number of intervention reviews in the field of treatment of oral and oropharyngeal cancers (Bessell 2011; Furness 2011; Glenny 2010) and screening programmes for the early detection and prevention of oral cancer (Brocklehurst 2010a). This screening test accuracy review will complement the intervention reviews.

OBJECTIVES

The objective of this review is to estimate the accuracy of the conventional oral examination (COE) used singly or in combination with another index test (Additional Table 1) as a screening test for the detection of oral cancer and potentially malignant disorders (PMD) of the lip and oral cavity of apparently healthy adults.

Secondary objectives

The secondary objective of this review is to estimate the accuracy of the different tests with COE, when compared with each other.

Investigation of sources of heterogeneity

We will use meta-regression to explore possible sources of heterogeneity, ways in which the observed diagnostic test accuracy varies according to particular characteristics. Covariates to be included in these analyses will include.

- Characteristics of the study sample: prevalence of carcinoma or PMD disease in the study (> 50% prevalence), inclusion of HPV + adults, tobacco users / high alcohol consumption.
- Target condition (oral squamous cell carcinoma alone or oral squamous cell carcinoma and PMD).
- Attributes of the screening programme: prospective organised or opportunistic, type of reference standard (examination and clinical evaluation by physician with specialist

knowledge or extended follow-up), operator (dental or general practice professionals or other healthcare workers).

METHODS

Criteria for considering studies for this review

Types of studies

Studies of clinical cohorts of apparently healthy adults which report the diagnostic accuracy of the conventional oral examination (COE) used singly or in combination with an index test listed in Additional Table 1, for oral cancer and potentially malignant disorders (PMD) with respect to one of the reference standards. These will include cross-sectional studies (or consecutive series) and randomised controlled trials (RCTs) of test accuracy. We will exclude case series and case control studies which can lead to inflated estimates of prevalence and test accuracy (Whiting 2004). We will exclude studies reported in abstract form alone, uncontrolled reports and randomised controlled trials of the effectiveness of screening programmes (intervention studies). Where randomised or paired comparative designs are available these will be included in the review and analysed separately. Studies should report data for true positives, false positives, true negatives and false negatives for each test. Only studies reporting results at an individual level (as opposed to a lesion level) will be included.

Participants

Apparently healthy adults (aged 16 years or over) attending an organised screening programme or screened during attendance at a dental or general practice examination.

Index tests

The COE used as a screen, alone or in combination with any other screening tests previously listed (Additional Table 1). The COE (comparator test) is the initial point of the screen, which all adults will receive. The index test will be used as an adjunct following the COE irrespective of whether oral cancer or PMD is suspected by the COE alone (i.e. a positive test result is a positive result from either the COE or the index test or both).

Comparator tests

The comparator test is the COE used as a stand alone screen as previously described. This is the standard screen commonly used in clinical practice, although problems with its use have been identified (Lingen 2008).

Target conditions

Following the consensus views of the expert working group of the WHO Collaborating Centre for Oral Cancer and Precancer, the target conditions of the lip or oral cavity of interest are noted as: Carcinoma

- Squamous cell carcinoma

Potentially malignant disorders

- Leukoplakia
- Erythroplakia
- Lichen planus
- Lupus erythematosus
- Submucous fibrosis
- Actinic keratosis
- Hereditary disorders such as dyskeratosis congenita or epidermolysis bullosa.

Reference standards

The reference standard is examination and clinical evaluation by a physician with specialist knowledge, such as an oral and maxillofacial pathologist or oral medicine specialist, working to the current diagnostic guidelines of their locality. We will include the diagnostic protocol / guidelines used in each study in the 'Characteristics of included studies' table. Studies with confirmatory biopsy of individuals who screened negative by the index test may exist, but are questionable ethically (Downer 2004). Confirmation of individuals screened negative by an index test may be done by extended follow-up of a minimum of 5 years.

Search methods for identification of studies

Electronic searches

The following databases will be searched using a highly sensitive search strategy:

- Cochrane Oral Health Group's Trials Register (to present)
- Cochrane Register of Diagnostic Test Accuracy Studies (to present)
- MEDLINE (1948 to present)
- EMBASE (1980 to present)
- MEDION (2003 to present).

The MEDLINE search strategy outlined in Appendix 1 will be modified for the listed databases. The search will not be limited by language or publication status. Non-English articles will be translated, unless a translator cannot be found through The Cochrane Collaboration.

The search strategy has been constructed in accordance with this protocol and that of a companion Cochrane Diagnostic Test Accuracy review 'Diagnostic tests for oral cancer in patients presenting

with clinically suspicious lesions' being undertaken concurrently by the same review team.

Searching other resources

We will also search relevant conference proceedings. We will locate further studies through citation searches and reference lists of key articles, and by contacting authors of identified articles to request information of any unpublished or ongoing studies.

Data collection and analysis

Selection of studies

Titles and abstracts of all articles identified from the searches will be independently assessed by two review authors (Tanya Walsh (TW) and Joseph Liu (JL)). For articles that appear to meet the inclusion criteria, or where a clear decision cannot be made from scanning the title and abstract alone, full reports will be obtained. Two review authors (TW and JL) will independently assess each report for inclusion. Where disagreements occur, the review authors will attempt to resolve these by discussion. If needed, a third review author (Paul Brocklehurst (PB)) will be asked to help resolve the discrepancies in consultation with the other two review authors.

Data extraction and management

Two review authors (TW and JL) will independently extract data using a piloted data collection form. Discrepancies will be resolved through discussion. If an agreement cannot be reached, a third review author will be consulted (PB). Study authors will be contacted to obtain relevant missing data if these are not available in the printed report.

The following data will be recorded from each study.

- Sample characteristics (age, sex, socio-economic status, risk factors (e.g. HPV status, prevalence of tobacco use and alcohol consumption), number of participants / lesions, lesion site)
- Setting (country, disease prevalence, type of screening)
- The type of index test(s) used (category, name, positivity threshold)
- Study information (design, reference standard, case definition, training and calibration of personnel)
- Study results (true positive, true negative, false positive, false negative, any equivocal results, withdrawal).

Assessment of methodological quality

The revised QUADAS tool, QUADAS-2 (Whiting 2011) will be used to assess the quality of the primary diagnostic studies over four key domains: patient selection, index test, reference standard and flow and timing of participants through the study.

In the first phase of the tool, the review question will be stated in terms of patient sample, index test, reference standard and target condition. This information is detailed in the [Criteria for considering studies for this review](#) section of this protocol. In phase two, the QUADAS-2 tool will be tailored to use with this review (Additional [Table 2](#)). Review specific guidance will be used to facilitate documentation of the pertinent descriptive information contained in the primary studies. Customised instructions to aid judgement of the signalling questions will be given (following [Patton 2008](#)). Two core signalling questions were removed: 'Was a case-control design avoided?' (this study design was excluded from the review); 'Did all patients receive a reference standard?' (this was a criterion for inclusion). Three additional signalling items relating to commercial funding, training and calibration and multiple index tests have been added to the core signalling questions. In phase three, a flow diagram will be drawn. In the final phase, an overall judgement of risk of bias and applicability is to be undertaken. A risk of bias judgement ('high', 'low' or 'unclear') will be reached for each domain. If the answers to all signalling questions within a domain are judged as 'yes' indicating low risk of bias, then the domain will be judged to be at low risk of bias. If any signalling question within a domain is judged as 'no' indicating high risk of bias then this indicates that potential bias exists. This will be followed by a judgement for concerns regarding applicability for the patient selection, index test and reference standard domains. We will pilot the use of the QUADAS-2 tool independently on five study reports. Where disagreements occur between the two review authors the review specific descriptions will be clarified until consistency is obtained.

Results of the quality assessment for all included studies will be summarised in a narrative report. A summary tabular presentation of the results for each domain will be also provided separately for risk of bias and concerns regarding applicability, along with a graphical display summarising this information.

Statistical analysis and data synthesis

Data for the true positive, true negative, false positive and false negative values for each test in each study will be tabulated. For each index test, estimates of diagnostic accuracy as sensitivity and specificity along with their 95% confidence intervals will be displayed as coupled forest plots, and plotted in receiver operating characteristic (ROC) space.

Meta-analysis will be used to combine the results of studies for each index test. Random-effects models will be used. If however there are too few studies to reliably estimate between study variability then fixed-effect models will be used. The statistical software SAS will be used throughout (SAS Institute Inc., Cary, USA).

Classification of responses of the various tests are given in Additional [Table 1](#). Where a common threshold of response is reported, the analysis will estimate the expected values of sensitivity and specificity (Bivariate approach [Reitsma 2005](#)). Where there is a variation in thresholds, the expected hierarchical summary ROC

curve for the tests across different thresholds will be estimated (Macaskill 2010; Rutter 2001). Hierarchical SROC curves will be fitted using the Proc NLMixed procedure in SAS. The proposed analysis is subject to change based on information reported in the primary studies. For example, if there is little variation in the positive thresholds of the blood and salivary index tests, it will not be appropriate to attempt to fit a summary ROC curve (Macaskill 2010). Where the primary studies have published more than one threshold result, accuracy estimates will be reported for all the thresholds. Data from all thresholds will be extracted and used in the analysis.

The statistical analysis plan can be specified as follows.

Primary analyses: The primary analyses will compare the COE used singly or in combination with any other test (Additional Table 1) with the reference standard.

- COE
- COE with vital rinse
- COE with light detection
- COE with blood / salivary analysis.

This will either estimate the average sensitivity and specificity of a test or describe the variation in sensitivity and specificity at different thresholds by estimating a hierarchical summary ROC curve, depending on nature of the index tests. Parameter estimates will include sensitivity, specificity and their correlation or hierarchical summary ROC curve.

Secondary analysis: The comparative accuracy of the index tests with the reference standard will be the focus of the secondary analyses. A preliminary analysis will graphically display the sensitivities and specificities of the index tests. This will be followed by a series of indirect pairwise analyses and structured as follows.

- COE versus COE with vital rinse
- COE versus COE with light detection
- COE versus COE with blood / salivary analysis
- COE with vital rinse versus COE with light detection
- COE with vital rinse versus COE with blood / salivary analysis

analysis

- COE with light detection versus COE with blood / salivary analysis.

All studies will be included in each pairwise comparison. Where studies of direct comparisons exist (i.e. paired data from all patients or randomising individuals to different tests) the results of these studies will be analysed and reported separately.

The methodology used is akin to the investigation of heterogeneity (as below) i.e. adding a covariate for test type into the Bivariate or

Hierarchical SROC analysis.

Investigations of heterogeneity

Meta-regression analyses will be carried out to explore possible sources of heterogeneity. Covariates to be included in these analyses will include characteristics of the study sample (prevalence of carcinoma or PMD in the study (> 50% prevalence), inclusion of HPV + adults, tobacco users / high alcohol consumption); target condition (oral squamous cell carcinoma alone or oral squamous cell carcinoma and PMD); aspects of study design (prospective organised or opportunistic); type of reference standard (examination and clinical evaluation by physician with specialist knowledge or extended follow-up) and operator (dental or general practice professionals or other healthcare workers). Different thresholds of binary categorisations will also be considered.

The log likelihood of models including the covariate will be compared to those models without the covariate. Formal model comparisons of either the Hierarchical SROC or Bivariate models will be undertaken using the Likelihood Ratio statistic to statistically compare the effects of adding or removing covariates. If statistical evidence of heterogeneity is found further investigations will be undertaken.

Sensitivity analyses

Sensitivity analyses will be conducted. This will entail restricting the analysis to studies where the reference standard is examination and clinical evaluation by physician with specialist knowledge or where differential verification is avoided.

Assessment of reporting bias

Tests for reporting bias will not be conducted because current tests are misleading when applied to systematic reviews of diagnostic test accuracy (Leeflang 2008; Tang 2000).

ACKNOWLEDGEMENTS

We would like to thank NHS Education Scotland and the Scottish Dental Clinical Effectiveness Programme for their support on this review; Anne Littlewood for her advice on searching the literature; Derek Richards for his thoughtful feedback on the protocol; and Luisa Fernandez Mauleffinch and Philip Riley for their assistance in facilitating this review.

REFERENCES

Additional references

American Cancer Society 1992

American Cancer Society. Update January 1992: the American Cancer Society guidelines for the cancer-related checkup. *CA: a Cancer Journal for Clinicians* 1992;**42**(1): 44–5.

Ang 2010

Ang KK, Harris J, Wheeler R, Weber R, Rosenthal DI, Nguyen-Tan PF, et al. Human papillomavirus and survival of patients with oropharyngeal cancer. *New England Journal of Medicine* 2010;**363**(1):24–35.

Bessell 2011

Bessell A, Glenny A-M, Furness S, Clarkson JE, Oliver R, Conway DI, et al. Interventions for the treatment of oral and oropharyngeal cancers: surgical treatment. *Cochrane Database of Systematic Reviews* 2011, Issue 9. [DOI: 10.1002/14651858.CD006205.pub3]

Brinkmann 2011

Brinkmann O, Kastratovic DA, Dimitrijevic MV, Konstantinovic VS, Jelovac DB, Antic J, et al. Oral squamous cell carcinoma detection by salivary biomarkers in a Serbian population. *Oral Oncology* 2011;**47**(1):51–5.

Brocklehurst 2010a

Brocklehurst P, Kujan O, Glenny AM, Oliver R, Sloan P, Ogden G, et al. Screening programmes for the early detection and prevention of oral cancer. *Cochrane Database of Systematic Reviews* 2010, Issue 11. [DOI: 10.1002/14651858.CD004150.pub3]

Brocklehurst 2010b

Brocklehurst PR, Baker SR, Speight PM. Oral cancer screening: what have we learnt and what is there still to achieve?. *Future Oncology* 2010;**6**(2):299–304.

Buchen 2011

Buchen L. Cancer: Missing the mark. Why is it so hard to find a test to predict cancer?. *Nature* 2011;**471**(7339): 428–32.

Cancer Research UK

Cancer Research UK. <http://info.cancerresearchuk.org/cancerstats/types/oral/> (accessed 8th April 2011).

Conway 2008

Conway DI, Petticrew M, Marlborough H, Berthiller J, Hashibe M, Macpherson LM. Socioeconomic inequalities and oral cancer risk: a systematic review and meta-analysis of case-control studies. *International Journal of Cancer* 2008;**122**(12):2811–9.

Downer 2004

Downer MC, Moles DR, Palmer S, Speight PM. A systematic review of test performance in screening for oral cancer and precancer. *Oral Oncology* 2004;**40**(3):264–73.

Faggiano 1997

Faggiano F, Partanen T, Kogevinas M, Boffetta P. Socioeconomic differences in cancer incidence and

mortality. In: Kogevinas M, Pearce N, Susser M, Boffetta P editor(s). *Social Inequalities and Cancer*. Lyon: IARC Scientific Publications No 138. International Agency for Research in Cancer, 1997.

Fedele 2009

Fedele S. Diagnostic aids in the screening of oral cancer. *Head & Neck Oncology* 2009;**1**:5.

Ferlay 2010

Ferlay J, Shin HR, Bray F, Forman D, Mathers C, Parkin DM. Estimates of worldwide burden of cancer in 2008: GLOBOCAN 2008. *International Journal of Cancer* 2010;**127**(12):2893–917.

Freedman 2007

Freedman ND, Abnet CC, Leitzmann MF, Hollenbeck AR, Schatzkin A. Prospective investigation of the cigarette smoking-head and neck association by sex. *Cancer* 2007;**110**(7):1593–601.

Furness 2011

Furness S, Glenny AM, Worthington HV, Pavitt S, Oliver R, Clarkson JE, et al. Interventions for the treatment of oral cavity and oropharyngeal cancer: chemotherapy. *Cochrane Database of Systematic Reviews* 2011, Issue 4. [DOI: 10.1002/14651858.CD006386.pub3]

Garg 2005

Garg AX, Adhikari NK, McDonald H, Rosas-Arellano MP, Devereaux PJ, Beyene J, et al. Effects of computerized clinical decision support systems on practitioner performance and patient outcomes: a systematic review. *JAMA* 2005;**293**(10):1223–38.

Glenny 2010

Glenny AM, Furness S, Worthington HV, Conway DI, Oliver R, Clarkson JE, et al. Interventions for the treatment of oral cavity and oropharyngeal cancer: radiotherapy. *Cochrane Database of Systematic Reviews* 2010, Issue 12. [DOI: 10.1002/14651858.CD006387.pub2]

Jaber 2003

Jaber MA, Porter SR, Speight P, Eveson JW, Scully C. Oral epithelial dysplasia: clinical characteristics of western European residents. *Oral Oncology* 2003;**39**(6):589–96.

La Vecchia 1997

La Vecchia C, Tavani A, Franceschi S, Levi F, Corrao G, Negri E. Epidemiology and prevention of oral cancer. *Oral Oncology* 1997;**33**(5):302–12.

Landis 1977

Landis JR, Koch GG. The measurement of observer agreement for categorical data. *Biometrics* 1977;**33**(1): 159–74.

Lee 2009

Lee JM, Garon E, Wong DT. Salivary diagnostics. *Orthodontics and Craniofacial Research* 2009;**12**(3):206–11.

Leeflang 2008

Leeflang MM, Deeks JJ, Gatsonis C, Bossuyt PM, Cochrane Diagnostic Test Accuracy Working Group. Systematic

- reviews of diagnostic test accuracy. *Annals of Internal Medicine* 2008;**149**(12):889–97.
- Leeftang 2008a**
Leeftang MM, Moons KG, Reitsma JB, Zwinderman AH. Bias in sensitivity and specificity caused by data-driven selection of optimal cutoff values: mechanisms, magnitude, and solutions. *Clinical Chemistry* 2008;**54**(4):729–37.
- Leston 2010**
Seoane Leston J, Diz Dios P. Diagnostic clinical aids in oral cancer. *Oral Oncology* 2010;**46**(6):418–22.
- Li 2006**
Li Y, Elashoff D, Oh M, Sinha U, St John MA, Zhou X, et al. Serum circulating human mRNA profiling and its utility for oral cancer detection. *Journal of Clinical Oncology* 2006;**24**(11):1754–60.
- Lingen 2008**
Lingen MW, Kalmar JR, Karrison T, Speight PM. Critical evaluation of diagnostic aids for the detection of oral cancer. *Oral Oncology* 2008;**44**(1):10–22.
- Liu 2006**
Liu JL, Wyatt JC, Deeks JJ, Clamp S, Keen J, Verde P, et al. Systematic reviews of clinical decision tools for acute abdominal pain. *Health Technology Assessment* 2006;**10**(47):1–167.
- Lodi 2006**
Lodi G, Sardella A, Bez C, Demarosi F, Carrassi A. Interventions for treating oral leukoplakia. *Cochrane Database of Systematic Reviews* 2006, Issue 4. [DOI: 10.1002/14651858.CD001829.pub3]
- Macaskill 2010**
Macaskill P, Gatsonis C, Deeks JJ, Harbord RM, Takwoingi Y. Chapter 10: Analysing and presenting results. In: Deeks JJ, Bossuyt PM, Gatsonis C (editors). *Cochrane Handbook for Systematic Reviews of Diagnostic Test Accuracy* version 1.0 (updated December 2010). The Cochrane Collaboration, 2010. Available from <http://srda.cochrane.org/>.
- Macfarlane 1995**
Macfarlane GJ, Zheng T, Marshall JR, Boffetta P, Niu S, Brasure, et al. Alcohol, tobacco, diet and the risk of oral cancer: a pooled analysis of three case-control studies. *European Journal of Cancer. Part B, Oral Oncology* 1995;**31B**(3):181–7.
- McGurk 2010**
McGurk M, Scott SE. The reality of identifying early oral cancer in the general dental practice. *British Dental Journal* 2010;**208**(8):347–51.
- Moles 2002**
Moles DR, Downer MC, Speight PM. Meta-analysis of measures of performance reported in oral cancer and precancer screening studies. *British Dental Journal* 2002;**192**(6):340–4.
- Napier 2008**
Napier SS, Speight PM. Natural history of potentially malignant oral lesions and conditions: an overview of the literature. *Journal of Oral Pathology and Medicine* 2008;**37**(1):1–10.
- Netuveli 2006**
Netuveli G, Sheiham A, Watt RG. Does the 'inverse screening law' apply to oral cancer screening and regular dental check-ups?. *Journal of Medical Screening* 2006;**13**(1):47–50.
- Ogden 2005**
Ogden GR. Alcohol and oral cancer. *Alcohol* 2005;**35**(3):169–73.
- Parkin 2001**
Parkin DM. Global cancer statistics in the year 2000. *Lancet Oncology* 2001;**2**(9):533–43.
- Patton 2008**
Patton LL, Epstein JB, Kerr AR. Adjunctive techniques for oral cancer examination and lesion diagnosis: a systematic review of the literature. *Journal of American Dental Association* 2008;**139**(7):896–905.
- Petti 2003**
Petti S. Pooled estimate of world leukoplakia prevalence: a systematic review. *Oral Oncology* 2003;**39**(8):770–80.
- Reibel 2003**
Reibel J. Prognosis of oral pre-malignant lesions: significance of clinical, histopathological, and molecular biological characteristics. *Critical Reviews in Oral Biology and Medicine* 2003;**14**(1):47–62.
- Reitsma 2005**
Reitsma JB, Glas AS, Rutjes AW, Scholten RJ, Bossuyt PM, Zwinderman AH. Bivariate analysis of sensitivity and specificity produces informative summary measures in diagnostic reviews. *Journal of Clinical Epidemiology* 2005;**58**(10):982–90.
- Rethman 2010**
Rethman MP, Carpenter W, Cohen EE, Epstein J, Evans CA, Flaitz CM, et al. Evidence-based clinical recommendations regarding screening for oral squamous cell carcinomas. *Journal of the American Dental Association* 2010;**141**(5):509–20.
- Rogers 2009**
Rogers SN, Brown JS, Woolgara JA, Lowe D, Magennis P, Shaw RJ, et al. Survival following primary surgery for oral cancer. *Oral Oncology* 2009;**45**(3):201–11.
- Rosin 2006**
National Institute of Dental and Craniofacial Research. Bringing the promise of molecular medicine to oral cancer screening. <http://www.nidcr.nih.gov/Research/ResearchResults/InterviewsOHR/TIS032006.htm> (accessed 18th April 2011).
- Rusthoven 2010**
Rusthoven KE, Raben D, Song JI, Kane M, Altoos TA, Chen C. Survival and patterns of relapse in patients with oral tongue cancer. *Journal of Oral and Maxillofacial Surgery* 2010;**68**(3):584–9.

- Rutter 2001**
Rutter CM, Gatsonis CA. A hierarchical regression approach to meta-analysis of diagnostic test accuracy evaluations. *Statistics in Medicine* 2001;**20**(19):2865–84.
- Sankaranarayanan 1997**
Sankaranarayanan R. Health care auxiliaries in the detection and prevention of oral cancer. *Oral Oncology* 1997;**33**(3):149–54.
- Scully 2000a**
Scully C, Porter S. ABC of oral health. Oral cancer. *BMJ* 2000;**321**(7253):97–100.
- Scully 2000b**
Scully C, Shotts R. ABC of oral health. Mouth ulcers and other causes of orofacial soreness and pain. *BMJ* 2000;**321**(7254):162–5.
- Scully 2009**
Scully C, Bagan J. Oral squamous cell carcinoma overview. *Oral Oncology* 2009;**45**(4-5):301–8.
- Silverman 1984**
Silverman S Jr, Gorsky M, Lozada F. Oral leukoplakia and malignant transformation. A follow-up study of 257 patients. *Cancer* 1984;**53**(3):563–8.
- Speight 2006**
Speight PM, Palmer S, Moles DR, Downer MC, Smith DH, Henriksson M, et al. The cost-effectiveness of screening for oral cancer in primary care. *Health Technology Assessment* 2006;**10**(14):1–144.
- Tang 2000**
Tang JL, Liu JL. Misleading funnel plot for detection of bias in meta-analysis. *Journal of Clinical Epidemiology* 2000;**53**(5):477–84.
- US Preventive Services Task Force**
US Preventive Services Task Force. <http://www.uspreventiveservicestaskforce.org/> (accessed 8th April 2011).
- van der Waal 2009**
van der Waal I. Potentially malignant disorders of the oral and oropharyngeal mucosa: terminology, classification and present concepts of management. *Oral Oncology* 2009;**45**(4-5):317–23.
- Warnakulasuriya 2007**
Warnakulasuriya S, Johnson NW, van der Waal I. Nomenclature and classification of potentially malignant disorders of the oral mucosa. *Journal of Oral Pathology and Medicine* 2007;**36**(10):575–80.
- Warnakulasuriya 2009**
Warnakulasuriya S. Global epidemiology of oral and oropharyngeal cancer. *Oral Oncology* 2009;**45**(4-5):309–16.
- Whiting 2004**
Whiting P, Rutjes AW, Reitsma JB, Glas AS, Bossuyt PM, Kleijnen J. Sources of variation and bias in studies of diagnostic accuracy: a systematic review. *Annals of Internal Medicine* 2004;**140**(3):189–202.
- Whiting 2011**
Whiting PF, Rutjes AW, Westwood ME, Mallett S, Deeks JJ, Reitsma JB, et al. QUADAS-2: a revised tool for the quality assessment of diagnostic accuracy studies. *Annals of Internal Medicine* 2011;**155**(8):529–36.
- Wyatt 1995**
Wyatt JC, Altman DG. Commentary: Prognostic models: clinically useful or quickly forgotten?. *BMJ* 1995;**311**:1539.
- Yusof 2006**
Yusof ZY, Netuveli G, Ramli AS, Sheiham A. Is opportunistic oral cancer screening by dentists feasible? An analysis of the patterns of dental attendance of a nationally representative sample over 10 years. *Oral Health and Preventive Dentistry* 2006;**4**(3):165–71.
* Indicates the major publication for the study

ADDITIONAL TABLES

Table 1. Screening tests for PMDs and oral cavity cancer

Test	Characteristics	Classification of response	Other information
Conventional oral examination (COE).	A standard visual and tactile examination of the oral mucosa under normal (incandescent) light	The presence of an oral mucosal abnormality is classified as a positive test result; the absence of any oral mucosal abnormalities is classified as a negative test result	Traditionally been used as an oral cancer screen, but its utility is debated (Lingen 2008). Advantages: quick and easy once trained, minimally invasive. Disadvantages: oral mucosal abnormalities are not necessarily clinically or biologically ma-

Table 1. Screening tests for PMDs and oral cavity cancer (Continued)

			lignant; only a small percentage of leukoplakias are progressive or become malignant; COE cannot distinguish between those that are or are not; some precancerous lesions may exist within oral mucosa that appears clinically normal by COE alone (Lingen 2008).
Vital rinsing (e.g. Toluidine Blue, Tolonium chloride).	Vital rinsing refers to the use of dyes such as Toluidine Blue or Tolonium chloride to stain oral mucosa tissues for PMD or malignancy (Leston 2010; Lingen 2008; Patton 2008). The procedure is as follows. <ul style="list-style-type: none"> • Pre-rinse with acetic acid • Rinse with water • Apply Toluidine Blue • Post-rinse with acetic acid • Rinse with water • Observe mucosa to check for staining. 	The result of the test is classified as positive if tissue is stained and negative if no tissue is stained, or equivocal if no definitive result can be obtained	Advantages: ability to define areas that could be malignant or abnormal but cannot be seen; assess the extent of the PMD for excision. Disadvantages: benign inflammatory lesions subject to stain; failure of some cancerous lesions to stain; variation in test performance depending on how thorough the test procedures are followed; contraindicated in those who are known to be allergic to iodine
Light-based detection (e.g. ViziLite and ViziLite plus, Microlux/DL, VELscope, Identafi 3000)	Light-based systems to identify premalignant and malignant lesions and to highlight their presence through tissue reflectance (Leston 2010; Lingen 2008; Patton 2008). E.g. using ViziLite Plus or Microlux/DL, the procedure is as follows (Lingen 2008). <ul style="list-style-type: none"> • Pre-rinse with acetic acid • Use blue-light light source to visually assess the oral cavity. ViziLite Plus also provides a toluidine chloride solution (TBlue) to aid in the marking of the lesion for biopsy once the light source is removed	The result of the test is classed as negative if the appearance of the epithelium is lightly bluish white and positive if the appearance of the epithelium is distinctly white (acetowhite)	Advantages: simple to use; non-invasive; do not require consumable reagents; provide real time results; can be performed by a wide range of operators after a short training period. Disadvantages: the necessity of a dark environment; high initial set up (for VELscope) or recurrent costs (for ViziLite in low income countries); lack of permanent record unless photographed; inability to objectively measure visualisation results
Blood and saliva analysis.	These novel technologies are at an early stage of development and evaluation. Analysis of blood or saliva samples which tests for the presence of	Cut-off probabilities vary widely and are dependent on the individual biomarker or combination of biomarkers examined.	Advantages: non-invasive (saliva tests) or minimally invasive (blood tests). Disadvantages: there is a tendency for the estimated diag-

Table 1. Screening tests for PMDs and oral cavity cancer (Continued)

	<p>bio-markers of PMD and oral cancer (Brinkmann 2011; Lee 2009; Li 2006).</p>	<p>Molecular markers for diagnosis include changes in cellular DNA, altered mRNA transcripts, altered protein levels</p>	<p>nostic accuracy of new health technologies to decline over time as evidence from independent evaluations accumulate (Wyatt 1995). This bias, which can be substantial, has been demonstrated in other domains, e.g. acute abdominal pain (Liu 2006) and clinical decision support systems (Garg 2005). Promising biomarker tests in several clinical areas were eventually been shown to be disappointing (Buchen 2011). It remains to be seen whether this is the case with oral cancer and PMDs</p>
--	--------------------------------------------------------------------------------	--------------------------------------------------------------------------------------------------------------------------	----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

PMDs = potentially malignant disorders.

Table 2. Indicators for the assessment of methodological quality

Domain	Patient selection	Index test	Reference standard	Flow and timing
Description.	<p>Describe methods of patient selection. Describe included patients (characteristics, prior testing, presentation, intended use of index test and setting).</p>	<p>Describe the index test (s) and how it was conducted and interpreted. Describe the sequence of tests, any training or calibration of assessors (levels of agreement should be reported. Where this is measured by the kappa statistic, acceptable values range from 0.61 (moderate agreement) to 1.00 (almost perfect agreement) (Landis 1977)), any procedures taken to ensure blinding of examiners, post-hoc or a priori threshold specification, any conflict of interest or commercial funding</p>	<p>Describe the reference standard and how it was conducted and interpreted. Any measures taken to ensure assessors were blinded to the results of the index tests should be documented, along with the sequence of reference and index tests</p>	<p>Describe the characteristics and proportion of patients who did not receive the index test (s) and/or reference standard, who received a reference standard other than examination and clinical evaluation by a specialist physician, or who were excluded from the 2 x 2 table (refer to flow diagram). Describe the time interval and any interventions between index test (s) and reference standard. The length of time between the index test and reference standard should be short in the majority of cases. If the period elapsed between</p>

Table 2. Indicators for the assessment of methodological quality (Continued)

				initial screening and reference standard (examination and clinical evaluation) is greater than 6 weeks then this will be considered an unacceptable delay
Signalling questions (Yes/No/Unclear).	<p>Was a consecutive or random sample of patients enrolled? Classify as Yes if consecutive patients or a random sample of individuals were recruited Classify as No if non-consecutive patients or a non-random sample of individuals were recruited Classify as Unclear if patient selection was not clearly described.</p>	<p>Was calibration of examiners undertaken and results reported? Classify as Yes if the examiners participated in dedicated training and calibration was reported to an acceptable standard Classify as No if the examiners did not participate in dedicated training or was not assessed, or training was undertaken but calibration was not to an acceptable standard Classify as Unclear if the information on training and calibration was not stated</p>	<p>Is the reference standard likely to correctly classify the target condition? The reference standard is an examination and clinical evaluation by a physician with specialist knowledge which if stated as such should be acceptable. Ideally this should be undertaken independently by more than one specialist. Alternatively an acceptable reference standard is extended follow-up Classify as Yes if the test result is independently confirmed by at least two specialists Classify as No if the test result is not independently confirmed by at least two specialists or there was lack of agreement between specialists Classify as Unclear if the study does not state who confirmed the results.</p>	<p>Was there an appropriate time interval between the index test(s) and reference standard? Classify as Yes if the delay between the index test(s) and reference standard is considered acceptable for the majority of participants Classify as No if the delay between the index test(s) and reference standard is considered unacceptable for the majority of participants Classify as Unclear if the delay between the index test(s) and reference standard is not explicitly stated</p>
	<p>Did the study avoid inappropriate exclusions? Classify as Yes if the sample consisted of apparently healthy individuals. Classify as No if only individuals with existing PMDs were recruited. Classify as Unclear if exclusions were not clearly</p>	<p>Were the index test results interpreted without knowledge of the results of the reference standard? Classify as Yes if interpreters of index test results clearly do not know results of reference standard Classify as No if inter-</p>	<p>Were the reference standard results interpreted without knowledge of the results of the index test? Classify as Yes if specialists clearly do not know index test results when performing the examination and clinical evaluation or evaluating fol-</p>	<p>Did all patients receive the same reference standard? Classify as Yes if the same reference standard was used in all participants Classify as No if the same reference standard was not used in all participants Classify as Unclear if it</p>

Table 2. Indicators for the assessment of methodological quality (Continued)

	described.	<p>preters of index test results clearly know results of reference standard</p> <p>Classify as Unclear if study did not provide any information on whether interpreters of index tests were blinded to reference standard</p>	<p>low-up data</p> <p>Classify as No if specialists clearly know index test results when performing the examination and clinical evaluation or evaluating follow-up data</p> <p>Classify as Unclear if study did not provide any information on whether specialists were blinded to the index test results</p>	<p>is unclear whether different reference standards were used</p>
		<p>Where multiple index tests were used, were the results of the second index test interpreted without knowledge of the results of the first index test?</p> <p>Classify as Yes if index test results were interpreted without knowledge.</p> <p>Classify as No if the index test results were interpreted with knowledge.</p> <p>Classify as Unclear if it is unclear whether the results of the second index test were interpreted without knowledge of the results of the first index test</p>		<p>Were all patients included in the analysis?</p> <p>Classify as Yes if all patients were included in the analysis.</p> <p>Classify as No if only some patients were included in the analysis.</p> <p>Classify as Unclear if it is unclear whether all patients were included in the analysis.</p>
		<p>If a threshold was used, was it pre-specified?</p> <p>Classify as Yes if the threshold was pre-specified.</p> <p>Classify as No if the threshold was not pre-specified.</p> <p>Classify as Unclear if it is unclear whether the threshold was pre-specified.</p>		

Table 2. Indicators for the assessment of methodological quality (Continued)

		Were any conflicts of interest stated? Classify as Yes if the study declared no conflict of interest. Classify as No if the study declared a conflict of interest. Classify as Unclear if there was no information on conflict of interest.		
Risk of bias: High/Low/Unclear.	Could the selection of individuals have introduced bias?	Could the conduct or interpretation of the index test have introduced bias?	Could the reference standard, its conduct, or its interpretation have introduced bias?	Could the patient flow have introduced bias?
Concerns regarding applicability: High/Low/Unclear.	Are there concerns that the included individuals do not match the review question?	Are there concerns that the index test, its conduct, or interpretation differ from the review question?	Are there concerns that the target condition as defined by the reference standard does not match the review question?	

PMDs = potentially malignant disorders.

APPENDICES

Appendix I. MEDLINE via OVID search strategy

1. exp Mouth/
2. Cheek/
3. or/1-2
4. exp Carcinoma, squamous cell/di
5. exp Precancerous conditions/di
6. (tumor\$ or tumour\$ or cancer\$ or carcinoma\$ or carcinogen\$ or neoplas\$ or malignan\$ or metasta\$ or dysplas\$ or lesion\$ or ulcer\$).tw,ot.
7. (pre-cancer\$ or precancer\$ or premalignan\$ or precursor\$ or "lichen planus" or leukoplakia or "submucous fibrosis" or (actinic adj2 keratosis) or candidiasis or erythroplakia or erythroplas\$ or erythroleukoplakia or hyperplas\$ or hyperkeratos\$).tw,ot.
8. or/4-7
9. 3 and 8
10. exp Mouth neoplasms/di
11. Lichen Planus, Oral/di
12. Oral submucous fibrosis/di
13. Oral candidiasis/di

14. ((oral\$ or mouth\$ or bucca\$ or “oral cavit\$” or (oral adj mucosa\$) or (mouth adj mucosa\$) or lip or lips or tongue\$ or gingiv\$ or palat\$ or cheek\$ or “intra oral\$” or intraoral\$ or gum or gums or labial\$) adj3 (tumor\$ or tumour\$ or cancer\$ or carcinoma\$ or carcinogen\$ or neoplas\$ or malignan\$ or metasta\$ or dysplas\$ or lesion\$ or ulcer\$ or pre-cancer\$ or precancer\$ or premalignan\$ or precursor\$ or “lichen planus” or leukoplakia or “submucous fibrosis” or (actinic adj2 keratosis) or candidiasis or erythroplakia or erythroplas\$ or erythroleukoplakia or hyperplas\$ or hyperkerato\$)).tw,ot.
15. or/10-14
16. 9 or 15
17. Cytodiagnosis/
18. Cytological techniques/
19. Cytophotometry/
20. (brush adj3 biops\$).tw,ot.
21. (“oral cdx” or oralcdx).tw,ot.
22. (“modified liquid based cytology” or (exfoliat\$ adj3 cytolog\$)).tw,ot.
23. (brush\$ and (cytodiagnosis or cytopathology)).tw,ot.
24. Tolonium chloride/du
25. Coloring agents/du
26. (“tolonium chloride” or “tolu?dine blue” or “tolu?dine b” or tblue or t-blue).tw,ot.
27. (tolu?dine adj6 (dye\$ or rins\$ or stain\$ or wash\$)).tw,ot.
28. exp Luminescence/du
29. Fluorescence/
30. Spectrometry, fluorescence/
31. exp Luminescent Agents/du
32. Light/du
33. Tomography, Optical Coherence/
34. (visual\$ adj5 (“light emitting diode” or “blue spectrum” or LED or luminous\$)).tw,ot.
35. (visual?ation adj3 adjunct\$).tw,ot.
36. (vizilite or microlux\$ or orascope or velscope).tw,ot.
37. lumenoscop\$.tw,ot.
38. ((tumor\$ or tumour\$ or cancer\$ or carcinoma\$ or neoplas\$ or carcinogen\$ or malignan\$ or metasta\$ or lesion\$ or ulcer\$) adj5 (fluorescen\$ or autofluorescen\$ or luminescen\$ or chemiluminescen\$)).tw,ot.
39. (tissue adj3 reflect\$).tw,ot.
40. Spectrophotometry/
41. Acetic acid/du
42. (acetic acid adj3 (wash\$ or rins\$)).tw,ot.
43. acetowhite.tw,ot.
44. Saliva/an, ch
45. Tumor Markers, Biological/an
46. ((“tumo?r marker\$” or “neoplas\$ marker\$”) adj3 (blood or saliva)).tw,ot.
47. ((analy\$ or screen\$ or test\$ or examin\$) adj3 (blood or saliva)).tw,ot.
48. Diagnosis, Oral/
49. Mass screening/
50. Physical examination/
51. ((oral\$ or mouth\$) adj5 (exam\$ or histolog\$ or check\$ or inspect\$)).tw,ot.
52. (visual\$ adj3 (exam\$ or inspect\$ or screen\$)).tw,ot.
53. or/17-52
54. 16 and 53

HISTORY

Protocol first published: Issue 11, 2012

CONTRIBUTIONS OF AUTHORS

Tanya Walsh (TW) and Joseph Liu (JL) wrote this protocol with contributions from Paul Brocklehurst (PB), Graham Ogden (GO), Crispian Scully (CS), Ross Kerr (RK), Mark Lingem (ML) and Saman Warnakulasuriya (SW).

DECLARATIONS OF INTEREST

None declared at this point in time.

SOURCES OF SUPPORT

Internal sources

- MAHSC, UK.

The Cochrane Oral Health Group is supported by the Manchester Academic Health Sciences Centre (MAHSC) and the NIHR Manchester Biomedical Research Centre.

- NIHR-MBRC, UK.

Health Sciences Research, School of Dentistry is supported by the Manchester Academic Health Sciences Centre (MAHSC) and the NIHR Manchester Biomedical Research Centre.

- SCDEP, UK.

This review has been supported by the Scottish Dental Clinical Effectiveness Programme.

External sources

- Cochrane Oral Health Group Global Alliance, UK.

All reviews in the Cochrane Oral Health Group are supported by Global Alliance member organisations (British Orthodontic Society, UK; British Society of Paediatric Dentistry, UK; National Center for Dental Hygiene Research & Practice, USA and New York University College of Dentistry, USA) providing funding for the editorial process (www.ohg.cochrane.org).

- National Institute for Health Research (NIHR), UK.

All reviews in the Cochrane Oral Health Group are supported by NIHR Systematic Reviews Programme infrastructure funding.