



Short-term effects of CO₂ leakage on the soil bacterial community in a simulated gas leakage scenario

Jing Ma¹, Wangyuan Zhang², Shaoliang Zhang², Qianlin Zhu¹, Qiyang Feng¹ and Fu Chen¹

¹ Low Carbon Energy Institute, China University of Mining and Technology, Xuzhou, China

² School of Environment Science and Spatial Informatics, China University of Mining and Technology, Xuzhou, China

ABSTRACT

The technology of carbon dioxide (CO₂) capture and storage (CCS) has provided a new option for mitigating global anthropogenic emissions with unique advantages. However, the potential risk of gas leakage from CO₂ sequestration and utilization processes has attracted considerable attention. Moreover, leakage might threaten soil ecosystems and thus cannot be ignored. In this study, a simulation experiment of leakage from CO₂ geological storage was designed to investigate the short-term effects of different CO₂ leakage concentration (from 400 g m⁻² day⁻¹ to 2,000 g m⁻² day⁻¹) on soil bacterial communities. A shunt device and adjustable flow meter were used to control the amount of CO₂ injected into the soil. Comparisons were made between soil physicochemical properties, soil enzyme activities, and microbial community diversity before and after injecting different CO₂ concentrations. Increasing CO₂ concentration decreased the soil pH, and the largest variation ranged from 8.15 to 7.29 ($p < 0.05$). Nitrate nitrogen content varied from 1.01 to 4.03 mg/Kg, while Olsen-phosphorus and total phosphorus demonstrated less regular downtrends. The fluorescein diacetate (FDA) hydrolytic enzyme activity was inhibited by the increasing CO₂ flux, with the average content varying from 22.69 to 11.25 mg/(Kg h) ($p < 0.05$). However, the increasing activity amplitude of the polyphenol oxidase enzyme approached 230%, while the urease activity presented a similar rising trend. Alpha diversity results showed that the Shannon index decreased from 7.66 ± 0.13 to 5.23 ± 0.35 as the soil CO₂ concentration increased. The dominant phylum in the soil samples was *Proteobacteria*, whose proportion rose rapidly from 28.85% to 67.93%. In addition, the proportion of *Acidobacteria* decreased from 19.64% to 9.29% ($p < 0.01$). Moreover, the abundances of genera *Methylophilus*, *Methylobacillus*, and *Methylovorus* increased, while *GP4*, *GP6* and *GP7* decreased. Canonical correlation analysis results suggested that there was a correlation between the abundance variation of *Proteobacteria*, *Acidobacteria*, and the increasing nitrate nitrogen, urease and polyphenol oxidase enzyme activities, as well as the decreasing FDA hydrolytic enzyme activity, Olsen-phosphorus and total phosphorus contents. These results might be useful for evaluating the risk of potential CO₂ leakages on soil ecosystems.

Submitted 21 April 2017
Accepted 22 October 2017
Published 14 November 2017

Corresponding author
Fu Chen, chenfu@cumt.edu.cn

Academic editor
Budiman Minasny

Additional Information and
Declarations can be found on
page 21

DOI 10.7717/peerj.4024

© Copyright
2017 Ma et al.

Distributed under
Creative Commons CC-BY 4.0

OPEN ACCESS

Subjects Environmental Sciences, Microbiology, Soil Science

Keywords CO₂ geological storage, Potential CO₂ leakage, *Acidobacteria*, *Methylophilus*, Soil enzyme activity

INTRODUCTION

Under the background of the rapid development of the world economy and the consumption of fossil fuels, global warming has become one of the most urgent challenges for human development and survival. Carbon dioxide (CO₂) capture and storage (CCS) technology was developed to separate and collect the CO₂ from the point source and inject to deep reservoir or salt water layer for long-term sequestration ([Zhou et al., 2013](#); [Ko et al., 2016](#); [Li et al., 2016](#)). The CCS method has a huge potential for long-term CO₂ emission reduction, and also has the advantage of low cost. Although CCS technology has made a great contribution to alleviating the global greenhouse effect, there remains a risk of leakage from geological storage. A strong, rapid leakage of CO₂ gas could cause huge losses of life and property. Studies have shown that once CO₂ leakage occurred, the concentration of CO₂ in the air may increase to 3–5%. As CO₂ is an asphyxiant gas, this concentration could cause headache, dizziness, nausea, and even death following long-term exposure ([West et al., 2005](#); [Patil, Colls & Steven, 2010](#); [Tian et al., 2013](#); [Andrzej & Katarzyna, 2013](#)). Furthermore, slow leakage of CO₂ into soil can lead to variations of soil gas composition, moisture, pH, and subsequently, microbial communities ([Beaubien et al., 2008](#)). This would eventually result in the change of the soil ecological environment. Therefore, it is crucial to be prepared to deal with the possibility and evaluate its risk on the soil environment ([Sáenz de Miera et al., 2016](#)). In addition, information regarding the impacts of accidental leakage on near surface soil communities has gradually enriched. Considering that bacteria are ubiquitous and possess tremendous metabolic and physiological versatility, they are an essential part in the soil ecosystem and are critical to virtually all biogeochemical processes ([Horner-Devine, Carney & Bohannan, 2004](#)). Moreover, bacteria are the most abundant and diverse group of soil microorganisms ([Acosta-Martínez et al., 2010](#)). Therefore, it is especially important to understand how the microbial community responds to CO₂ gas leakage ([Prosser, 2007](#); [Acosta-Martínez et al., 2008](#); [Van Der Heijden, Bardgett & Van Straalen, 2008](#); [Sáenz de Miera et al., 2014](#)).

In recent years, the rapid development of culture-independent methods has gradually led to increased research focus on microbial community diversity and composition. This has provided new horizons for assessing the reactions of bacterial community to external disturbances, such as elevated CO₂ concentration ([Dumbrell et al., 2010](#); [Lemos et al., 2011](#); [Maček et al., 2011](#); [Mandić-Mulec et al., 2012](#); [Sáenz de Miera et al., 2016](#); [Schloss et al., 2009](#); [Tait et al., 2015](#); [Taylor et al., 2014](#)). The increasing CO₂ concentration could bring about a variation of soil biochemical conditions, which would cause the shifting diversity or functionality of indigenous bacteria ([Fernández-Montiel et al., 2015](#)). However, the elevated CO₂ could directly affect species richness, or it may trigger changes in the composition without any discernible effect on diversity ([Morales & Holben, 2014](#)). The degradation of organic matter, as well as the biogeochemical cycling of nutrients may also be impacted ([Tait et al., 2015](#)).

Many researchers have investigated the impacts of CO₂ leakages from geological sources in natural sites ([Frerichs et al., 2013](#); [Gabilondo & Bécáres, 2014](#); [Krüger et al., 2011](#); [McFarland, Waldrop & Haw, 2013](#); [Oppermann et al., 2010](#); [Sáenz de Miera et al., 2014](#)).

Considering the long periods of CO₂ emissions from natural seepages, the ecosystem might have already adapted through species substitution (Krüger *et al.*, 2011). Furthermore, these natural sites have presented existing special features, such as soil type, soil moisture and temperature, which might have not appeared in the potential scenario of a CCS leakage. Therefore, the results from the natural sites may not be extrapolated to the conditions occurring in other areas after possible leakage from the CCS process (Ziogou *et al.*, 2013; Fernández-Montiel *et al.*, 2015). Therefore, the current study was carried out in an experimental site in an attempt to assess the short-term effects on bacterial communities rather than the long-term effects. The aim was to analyze the first responses of a non-adapted ecosystem to simulated CO₂ leakage. Also, the consequences of the underground low gas emissions through the quantitative evaluation of soil microbial communities were investigated. The impacts on the soil bacterial community from the simulated below-ground CO₂ emissions were examined. In addition, a limited understanding of the importance of abiotic factors in regulating biodiversity and the structure of some important soil bacterial communities were also investigated (West *et al.*, 2009; Chen *et al.*, 2017). Moreover, soil enzymes are essential to the soil ecosystem for the material cycle and energy conversion. All biochemical reactions in the soil are catalyzed by soil enzymes, which also reflects the strength and direction of the biochemical processes in soil (Sardans, Penuelas & Estiarte, 2008; Duan *et al.*, 2015). Therefore, the soil enzyme activity can be used as an important indicator to evaluate the soil fertility. Usually many factors can affect soil enzyme activities, such as soil physicochemical properties, soil biota, agricultural vegetation and the human factor (Naidu & Sarathambal, 2012; Sarathambal *et al.*, 2016). However, few studies have examined the effect of high soil CO₂ levels on microbial enzyme activities (Beulig *et al.*, 2016).

In this study, a small simulation experiment, which includes an automatic control auxiliary device and continuous CO₂ injection, was established to simulate the leakage from a geological storage site. This simple simulation device was cheap and practical, fully meeting the demand for investigating the short-term effects on the bacterial community, which was different from our previous field simulation experimental platform (Chen *et al.*, 2016). The relationship between environmental factors and the soil bacterial community was also studied. The ecosystem functions could be changed overall by the community composition changes that enhance the presence of several taxa. Such knowledge will be useful to understand the potential effects of CO₂ leakages from geological storage sites on soil ecosystems in the future.

MATERIALS AND METHODS

Study site and experimental design

This experiment was established to simulate CO₂ leakage during CCS. This study was conducted at the Wenchang campus of China University of Mining and Technology (117.206194°E, 34.229568°N), Xuzhou, Jiangsu Province, China. The experiment contained five treatments, each with three replicates. The treatments consist of the following CO₂ gas flux levels: low (L, 400 g m⁻² day⁻¹), medium (M, 1,000 g m⁻² day⁻¹), high (H,




Figure 1 Schematic of the CO₂ leakage simulation experimental platform.

Full-size  DOI: [10.7717/peerj.4024/fig-1](https://doi.org/10.7717/peerj.4024/fig-1)

1,500 g m⁻² day⁻¹), extreme (E, 2,000 g m⁻² day⁻¹) and a control with no CO₂ injection (C). Containers used in this experiment were 30 cm (height) × 30 cm (diameter) and filled with a 25 cm topsoil layer. A layer of porous plate was placed at a distance of 5 cm from the container bottom. A CO₂ output control device was formed by connecting the cylinders, pressure reducing valve, diverter, PVC hose and other devices. The PVC hose was inserted into the center of the bottom of the container and fixed at the 2–3 cm mark in the bottom section above the outlet position (Fig. 1). During the experiment, the 5-cm space was filled with CO₂ gas, and the gas spread evenly into the soil.

By adjusting the pressing deducing valve and the adjustable flow meter, the uninterrupted CO₂ gas leaking into the soil in the four treatments were maintained treatment levels. The experiment began on March 19, 2016 (T0, before ventilation) and several samplings days took place subsequently (April 1st, April 16th, May 1st, June 3rd (denoted as T1, T2, T3, and T4, respectively)). The ventilation was then stopped and final sampling took place on July 4th (T5) to investigate the restoration of the soil ecological environment after the CO₂ stress was relieved.

Sample collection and soil property analysis

Approximately 100 g soil samples were collected from five points from the middle position of each container at about 10 cm deep and then were mixed to bulk samples from each container. Next, these mixed samples were put into a sterile sealing bag, and immediately sent to the laboratory. About 5 g soil was stored at –20 °C for subsequent analysis of microbial diversity. Other parts were placed directly in indoor shaded conditions for drying. Sand grinding was manually removed through a 2-mm screen. These samples were then passed through the 0.149-mm sieve for the testing of physical and chemical properties.

Soil pH and electrical conductivity (EC) values were measured using a pH meter and conductivity meter, respectively (PHC-3C, DDS-307A, Shanghai Leici, China). Organic matter (OM) content was measured with colorimetric methods using hydration heat

during the oxidation of potassium dichromate (*Chen et al., 2016*). Total nitrogen content was measured by dual wavelength spectrometry (*Norman, Edberg & Stucki, 1985*). The total phosphorus content was determined by the acid digestion molybdenum and antimony colorimetric method. The available phosphorus was measured by the hydrochloric acid ammonium chloride method.

Soil urease activity was measured by the phenol colorimetry of sodium hypochlorite (*Guan, 1986*). Polyphenol oxidase activity was determined by pyrogallol colorimetry (*Guan, 1986*). Protease activity was measured by the ninhydrin colorimetric method (*Guan, 1986*). The triphenyltetrazolium chloride (TTC) method was used to measure soil dehydrogenase activity (*Guan, 1986*). The fluorescein diacetate (FDA) hydrolysis activities were determined by the fluorescein colorimetric method (*Guan, 1986*). Three repetition groups were set up in each test. The negative control groups were provided that had either no substrate or no soil.

DNA extraction, PCR amplification and Illumina MiSeq sequencing

According to the instruction manual, 20 soil samples of DNA were extracted from 0.5 g fresh soil samples with the E.Z.N.ATM Mag-Bind Soil DNA Kit (Omega Biotek, Guangzhou, China). The 20 soil samples were marked as C0–C3, L0–L3, M0–M3, H0–H3, E0–E3 (collected at T0, T1, T2 and T3, respectively). The V3–V4 region of the bacterial 16S rRNA genes were amplified using the primer set 341F (CCCTACACGACGCTCTTCCGATCTG-CCTACGGGNGGCWGCAG) and 805R (GACTGGAGTTCCTTGGCACCCGA GAATTCCA-GACTACHVGGGTATCTAATCC). The 50 μ L PCR reaction system included 5 μ L of 10 \times PCR buffer, 0.5 μ L of each primer (50 μ M), 0.5 μ L of dNTP (10 mM each), 0.5 μ L Platinum Taq (5U/ μ L), 42 μ L sterilized ultrapure water and 1 μ L of template DNA (10 ng/ μ L). Amplification conditions were as follows: 3 min at 94 $^{\circ}$ C, 5 cycles of 94 $^{\circ}$ C for 30 s, 45 $^{\circ}$ C for 20 s, and 65 $^{\circ}$ C for 30 s, 20 cycles of 94 $^{\circ}$ C for 20 s, 55 $^{\circ}$ C for 20 s, and 72 $^{\circ}$ C for 30 s and final 5 min incubation at 72 $^{\circ}$ C to complete the extension process. According to the instruction manual, PCR products were pooled and purified with the Agencourt AMPure XP (Beckman, Indianapolis, IN, USA). Quantification of the purified PCR products was carried out using the Qubit2.0 test kit (Life Technologies, Carlsbad, CA, USA). Finally, 20 pmol DNA was analyzed sequentially using the Illumina MiSeq platform (Sangon Biotech, Shanghai, China).

Bioinformatics analysis

The sample sequencing data were distinguished with the barcode sequence, and the sequence of each sample underwent quality control. Then the non-specific amplification sequences and chimeric were removed with the usearch method (<http://www.drive5.com/usearch/>). Operational taxonomic units (OTUs) with a 97% similarity cutoff were also clustered using the usearch method. Mothur software was used to analyze the alpha diversity (<http://www.mothur.org/>). The Chao (<http://www.mothur.org/wiki/Chao>) and Shannon indices (<http://www.mothur.org/wiki/Shannon>) were conducted to reveal the richness and diversity. A RDP classifier was used to classify the species (<http://rdp.cme.msu.edu/misc/resources.jsp>). According to the taxonomic results, a

species' abundance diagram and rich infrared images were constructed with Origin8 and R software. The beta diversity analysis was performed using the weighted UniFrac metric algorithmic with R software. Canonical correlation analysis (CCA) was used to study the relationship between environmental stressors and microbial communities with Canoco 4.5 for windows software. SPSS 20 software was used to analyze the data of soil properties and enzyme activities and spearman correlations were calculated. Significance for all statistical analyses was accepted at $\alpha = 0.01$ or $\alpha = 0.05$. One-way ANOVA was performed and Duncan's test were applied in the case of multiple comparisons.

RESULTS

Effects of CO₂ leakage on the soil characteristics

The soil characteristics are summarized in Fig. 2. Figure 2A shows that soil pH changed significantly under CO₂ stress ($p < 0.05$). The pH value of C treatment showed a slight fluctuation. The value was 8.15 at T0, then it basically maintained at about 8.04 from T1 to T5. Considering the environmental impact of the test itself, the pH value of C treatment can be considered stable. The L, M, H and E treatments were significantly different from the C treatment ($p < 0.05$). The pH decreased significantly with the increasing CO₂ concentration. The spearman correlation coefficient was between -0.9 and -1 from T1 to T5. Eventually, at T5, the soil pH of the L and M treatments dropped to about 7.34, whereas the soil pH under the H and E treatments reached about 7.29. Figure 2B demonstrates that the EC first increased and then decreased overtime. The maximum EC value was detected at T2. Furthermore, the H treatment showed the maximum variation amplitude. Also, similar values at T4 and T5 in C and E treatments were observed. The soil organic matter showed a tendency to decrease abruptly and then increase (Fig. 2C). During the period of T0–T1, the content of organic matter decreased substantially, and then increased from T1 to T4. Once again, the organic matter content decreased in the recovery period (T5), and the value became lower than the initial content. Overall, the rates of change were greater in the L, M, H, and E treatments than that in the C treatment.

The nitrate nitrogen content of all treatments increased over time with some fluctuations (Fig. 2D). The average value of the L, M, H, and E treatments was about 1.01 mg/Kg at T0, whereas the value reached 4.03 mg/Kg at T5. The nitrate nitrogen value of the L, M, H, and E treatments increased significantly from T1 to T2. At T3, nitrate nitrogen values of the M, H and E treatments decreased slightly and then continued rising to T4 ($p < 0.05$). However, in the recovery period, the value declined again. The change trend of the total phosphorus was shown in Fig. 2E, which accumulated at T1, and then decreased at T2. In other words, after half a month of ventilation, the basic data increased and the increase amplitude was between 2.5% and 10% (T1). The value then suddenly fell approximately 25% ~ 33% at T2 (Fig. 2E). Also, from Fig. 2E it can be speculated that there was a tendency to decrease at T2–T5 when CO₂ is applied. Furthermore, the overall trend of the soil Olsen-phosphorus was complex (Fig. 2F). The Olsen-phosphorus in the C treatment fluctuated, but the L, M, H, and E treatments also fluctuated but at a significantly




Figure 2 Soil physicochemistry change under different CO₂ flux, (A) pH value; (B) EC value; (C) total organic matter; (D) nitrate-N; (E) total phosphorus; (F) Olsen-phosphorus. (C, Control; L, 400 g m⁻² d⁻¹; M, 1,000 g m⁻² d⁻¹; H, 1,500 g m⁻² d⁻¹; E, 2,000 g m⁻² d⁻¹).

Full-size DOI: [10.7717/peerj.4024/fig-2](https://doi.org/10.7717/peerj.4024/fig-2)

higher concentration. For example, the results of L treatment from T0 to T5 were 13.75 mg/Kg, 8.32 mg/Kg, 15.18 mg/Kg, 7.19 mg/Kg, 13.55 mg/Kg, and 7.44 mg/Kg, respectively. The M, H and E treatments had similar fluctuations to the L treatment presenting a different response.

Effects of CO₂ leakage on the soil enzyme activity

The redox enzyme plays an important role in the process of material transformation in soil. The transformation of organic matter, the formation of humus and its components are all related to the oxidation of the soil enzyme, such as dehydrogenase and polyphenol oxidase. Through the analysis of soil enzyme activity, the transformation of some nutrients and the status of soil fertility might be understood and predicted.

Dehydrogenase is an enzyme related with respiratory metabolism, and plays the role of hydrogen intermediate transfer body in the dehydrogenation reaction. [Figure 3A](#) shows that the dehydrogenase change under the L, M, H, and E treatments appeared as a decrease—rise—decrease tendency with different response times. At T1, excluding the C treatment, dehydrogenase activity of all treatments decreased, and then increased. The rising state of the L treatment ended after T2, while the rising phenomenon of the dehydrogenase activity of M, H, and E treatments lasted until T3 ([Fig. 3A](#)) ($p < 0.05$). At the recovery period, they showed a slight rise once again.

The FDA hydrolytic enzyme is widely used in soil quality assessment. The FDA hydrolysis activity presented a downward trend under the effects of CO₂ leakage ([Fig. 3B](#)) ($p < 0.05$). The average value of the L, M, H, and E treatments was 22.69 mg/ (Kg h) at T0, which then reduced to 11.25 mg/ (Kg h) at T5. The response of the L, M, H, and E treatments was uniform, excluding the M treatment, which appeared slightly larger than the other treatments at each time period ([Fig. 3B](#)).

Polyphenol oxidase could promote the oxidation of one or two or three phenols, and could also characterize the activity of some soil microorganisms. The activity of polyphenol oxidase showed an upward trend under CO₂ stress ([Fig. 3C](#)) ($p < 0.05$). The mean value of the L, M, H, and E treatments at T4 increased by 230% compared with the average value at T0. During the ventilation experiment, the activities of polyphenol oxidase showed different increasing trends. Compared with that at T4, there was little change of polyphenol oxidase activity at recovery phase.

The protease activity in the ventilation time showed an irregular variation trend in [Fig. 3D](#) ($p < 0.05$). For example, the activity of the L treatment firstly decreased at T1, increased at T2 and T3, and then decreased at T4. The activities of the M, H and E treatments increased at T1, declined at T2, increased at T3, and decreased at T4 ([Fig. 3D](#)). In the recovery period, the M treatment appeared to rise sharply, which could not be ruled out by the experimental error.

Soil urease is the key enzyme in the conversion of nitrogen in soil. The enzymatic reaction product of urease is one of the plant nitrogen sources, and its activity can be used to indicate the status of soil nitrogen. [Figure 3E](#) indicates that the urease activity also showed a complex change with a decrease - rise - decrease tendency ($p < 0.05$). The response time of each of the CO₂-addition treatments was basically consistent, whereas the response level was different. For times after T2, the urease activities of the M and E treatments were significantly higher than those of the L and H treatments. There was an increase in the recovery period ([Fig. 3E](#)).




Figure 3 Soil enzyme activity change under different CO₂ flux, (A) dehydrogenase activity; (B) FDA hydrolysis activity; (C) polyphenol oxidase activity; (D) protease activity; (E) urease activity. (C, Control; L, 400 g m⁻² d⁻¹; M, 1,000 g m⁻² d⁻¹; H, 1,500 g m⁻² d⁻¹; E, 2,000 g m⁻² d⁻¹).

Full-size [DOI: 10.7717/peerj.4024/fig-3](https://doi.org/10.7717/peerj.4024/fig-3)

Table 1 Alpha diversity indices of soil microorganisms (C, Control; L, 400 g m⁻² day⁻¹; M, 1,000 g m⁻² day⁻¹; H, 1,500 g m⁻² day⁻¹; E, 2,000 g m⁻² day⁻¹, where 0–3 represents samples collected at T0–T3, respectively).

Sample ID	Chao 1 index	Shannon index
C0	15,604.19 ± 1,704.233ef	7.65 ± 0.237d
L0	15,601.76 ± 425.667ef	7.66 ± 0.128d
M0	16,543.62 ± 1,022.004f	7.63 ± 0.264d
H0	17,185.51 ± 677.133f	7.454 ± 0.322d
E0	15,136.54 ± 645.322def	7.44 ± 0.113d
C1	14,815.69 ± 2,714.734def	7.38 ± 0.186d
C2	14,666.78 ± 1,144.127def	7.33 ± 0.070d
C3	14,567.04 ± 1,403.888def	7.30 ± 0.036d
L1	12,457.13 ± 1,062.828ab	7.11 ± 0.457d
L2	12,044.64 ± 1,172.405ab	7.03 ± 0.426d
L3	11,238.16 ± 1,177.577a	6.38 ± 0.293c
M1	11,163.37 ± 1,289.129a	6.38 ± 0.846c
M2	10,900.97 ± 1,346.372a	6.03 ± 0.486bc
M3	10,660.81 ± 1,103.014a	6.09 ± 0.544bc
H1	15,105.19 ± 652.463def	6.20 ± 0.041bc
H2	14,614.52 ± 572.322def	6.15 ± 0.040bc
H3	14,509.60 ± 998.913def	6.13 ± 0.523bc
E1	13,398.76 ± 627.144bcd	5.31 ± 0.464a
E2	14,295.70 ± 917.406cde	5.68 ± 0.301ab
E3	13,238.25 ± 216.176bcd	5.23 ± 0.347a

Effects of CO₂ leakage on the taxonomic composition of microbial consortia

The richness and diversity of microbial communities can be reflected by the Alpha diversity analysis (Table 1). After comparing the Chao 1 index of each treatment, it can be found that soil bacterial richness in the L and M treatments decreased with the increasing CO₂ concentration, while the H and E treatments showed a slight increase. In addition, the Shannon index was used to measure the heterogeneity of the community. The data in Table 1 also showed that Shannon index declined with the rising CO₂ concentration, which indicated that the microbial community diversity in the soil decreased under CO₂ stress.

Overall, the bacterial categories were relatively abundant in the 20 soil samples (Fig. 4A). *Proteobacteria*, *Acidobacteria*, *Bacteroidetes*, *Verrucomicrobia*, *Actinobacteria*, *Gemmatimonadetes*, *Firmicutes*, *Planctomycetes*, *Cyanobacteria*, *Chloroflexi*, *Candidatus Saccharibacteria* (TM7), and *Nitrospirae* comprised more than 90% of the total sequences in each soil sample. These high proportion phyla (>0.5%) were used to analyze the effects of stress on the soil bacterial structure. The most abundant phylum was *Proteobacteria* which accounted for 28.85%–67.93% (Fig. 4A). Figure 4A shows that the abundant proportion of *Proteobacteria* increased with the increasing CO₂ concentration. The average proportion was about 30.35% in the C treatment at T0, whereas it increased to 65.28% in the E treatment at T1. The abundance of phylum *Candidatus Saccharibacteria* (TM7) also




Figure 4 Taxonomic composition of soil samples on Phylum (A) and genus level (B). (C, Control; L, 400 g m⁻² day⁻¹; M, 1,000 g m⁻² day⁻¹; H, 1,500 g m⁻² day⁻¹; E, 2,000 g m⁻² day⁻¹, where 0–3 represents samples collected at T0–T3, respectively).

Full-size DOI: [10.7717/peerj.4024/fig-4](https://doi.org/10.7717/peerj.4024/fig-4)

presented a gradually increasing trend with the rising CO₂ flux. Furthermore, *Acidobacteria* was the second most abundant phylum. In the C treatment, it demonstrated an average abundance of 19.64% at T0. In the E treatment at T2, it was 9.29%. The higher the CO₂ stress concentration, the lower the abundance of phylum *Acidobacteria*. Moreover, the abundances of other phyla such as *Verrucomicrobia*, *Actinobacteria*, *Gemmatimonadetes*, *Firmicutes*, *Planctomycetes*, *Cyanobacteria*, *Chloroflexi*, and *Nitrospirae* decreased with the

increasing CO₂ flux (Fig. 4A). Furthermore, the proportion of *Bacteroidetes* showed no significant variability between treatments.

Figure 4B shows the abundance variation at the genus level. The abundances of genera *Methylophilus*, *Methylobacillus* and *Methylovorus* increased rapidly after the ventilation. The average proportions were 0.04%, 0.076% and 0.006% in the C treatment at T0. These numbers increased to 20.68%, 12.30% and 10.16% in the E treatment at T2. The higher the CO₂ concentration was, the greater was the abundance of genera. In addition, *Saccharibacteria*, *Variovorax* and *Methylotenera* presented the same trend. In contrast, the abundances of *Gp6*, *Gp4*, *Gemmatimonas*, *Gp7* and *Streptophyta* at T0 were 8.23%, 6.12%, 5.81%, 1.87% and 2.05%. These proportions reduced to 3.97%, 2.67%, 2.57%, 0.66% and 0.03% in the E group at T2. *Kofleria* and *Aciditerrimonas* showed the same trend that the higher the concentration of the stress concentration, the smaller the abundance.

Comparison of bacterial consortia compositions

Based on the weighted UniFrac algorithms, principal co-ordinates analysis (PcoA) was used to analyze the relation among the 20 samples (Fig. 5A). According to the PcoA, samples with higher similarity gather together, while samples with lower similarity are positioned further apart. Figure 5A shows that every sample of its own group was grouped closely, excluding the samples from the M treatment. At T0, from the C treatment to E treatment, their arrangement presented a pattern along the horizontal axis (Fig. 5A). This analysis also indicated that there were significant differences among these six bacterial community groups with different CO₂ leakage conditions. A sample distance heatmap was also one of the beta diversity indicators based on the weighted UniFrac algorithms, which presented the distance between the samples. In the heatmap, a color block represents the distance value (Fig. 5B). A deeper red color indicates a higher similarity between the samples, whereas a deeper blue color indicates a greater distance. The samples were also clustered in the heatmap, so the distance between samples can be seen through the cluster tree. As shown in Fig. 5B, the similarity was higher between samples in each group. The C and L treatments had higher similarity, whereas the M, H and E treatments had closer sample distances. Sample distances increased with the increasing CO₂ concentration, which is supported by the cluster tree.

Canonical correlation analysis

According to the above experimental results, it can be inferred that soil CO₂ concentration affected the soil physicochemical properties and microbial community structure, thus affecting the soil enzyme activity. Moreover, the soil community structure could be indirectly affected by the change of physicochemical properties and enzyme activity. Subsequently, CCA was used to analyze the correspondence between the environmental factors and microbial community composition under different CO₂ ventilation concentration.

As shown in Fig. 6, samples of the initial group at T0 were gathered with samples of the C treatment at T2. The sample locations of the L, M, H and E treatments were significantly changed. The angle size between the arrows of environmental factors represented the




Figure 5 Principal co-ordinates analysis (A) and heatmap of Beta diversity analysis (B) using weighted UniFrac. (C, Control; L, $400 \text{ g m}^{-2} \text{ day}^{-1}$; M, $1,000 \text{ g m}^{-2} \text{ day}^{-1}$; H, $1,500 \text{ g m}^{-2} \text{ day}^{-1}$; E, $2,000 \text{ g m}^{-2} \text{ day}^{-1}$, where 0–3 represents samples collected at T0–T3, respectively).

Full-size [DOI: 10.7717/peerj.4024/fig-5](https://doi.org/10.7717/peerj.4024/fig-5)




Figure 6 Canonical correlation analysis ordination plot indicating the relationship between the bacterial community and soil properties (A) or some frequent OTUs (B). (C, Control; L, $400 \text{ g m}^{-2} \text{ day}^{-1}$; M, $1,000 \text{ g m}^{-2} \text{ day}^{-1}$; H, $1,500 \text{ g m}^{-2} \text{ day}^{-1}$; E, $2,000 \text{ g m}^{-2} \text{ day}^{-1}$; 0 stands for the samples collected at T0; 1, 2 and 3 stand for the samples collected at T1, T2 and T3, respectively).

Full-size DOI: [10.7717/peerj.4024/fig-6](https://doi.org/10.7717/peerj.4024/fig-6)

correlation between environmental factors (Fig. 6A). The EC value, urease and polyphenol oxidase were significantly correlated with nitrate nitrogen. In addition, total phosphorus, FDA hydrolysis, Olsen phosphorus, and protease were grouped together, whereas pH, organic matter, and dehydrogenase had a greater distance. Moreover, the goodness of fit statistic for environmental variables indicated that ranking was highly correlated with pH, total phosphorus, urease, FDA hydrolysis, nitrate nitrogen, and polyphenol oxidase ($p < 0.01$) (see long arrow in Fig. 6A). The correlation between environmental factors and species information can be expressed by the angle between the environmental factor arrow line and the linking line which connected the species point and center point. Therefore, on the left part of Fig. 6B, the correlations between nitrate nitrogen, EC value and OTU3 (*Methylobacillus*), OTU9 (*Methylophilus*), OTU11 (*Methylophilus*), OTU12 (*Methylophilus*) were relatively large. Urease and polyphenol oxidase also had a closer relationship with OTU2 (*Methylophilus*) and OTU8 (*Methylovorus*).

DISCUSSION

Soil characteristic impacts under CO₂ leakage stress

The biological and biochemical processes in the soil are the important foundations of terrestrial ecosystems. Leakage of CO₂ during the geological sequestration process could elevate the soil CO₂ concentration (Österreicher Cunha et al., 2015; Šibanc et al., 2014). As a result, the soil physicochemical properties, soil enzyme activity and microbial community structure might be affected, thus influencing the terrestrial ecosystems (Awasthi et al., 2014). It is therefore essential to study the impact of CO₂ leakage.

In this study, the tested soil was weak alkaline (pH = 8.15) at the beginning of the experiment. As the ventilation concentration increased, the soil pH value decreased to 7.29. Moreover, in the early part of the experiment, the pH of each treatment decreased rapidly and gradually tended to be gentle. These results implied a strong negative correlation between CO₂ concentration and the pH value (r between -0.9 and -1). Sáenz de Miera et al. (2016) reported a similar result and concluded that the CO₂ injection introduced a pH value decrease of the CO₂ injection sites relative to a zero CO₂-addition control. Fernández-Montiel, Pedescoll & Bécáres (2016) showed that the soil pH changed from slightly acidic at the control site to acidic in higher CO₂ flux sites. Furthermore, Chen et al. (2016) showed a similar negative concentration between the pH value and the CO₂ flux. However, McFarland, Waldrop & Haw (2013) found that soil pH did not show a significant correlation with the CO₂ flux. In contrast, ventilation also had an obvious effect on the EC value. The EC value of soil leaching solution can reflect the water-soluble salt content in soil. The injection of CO₂ can affect the solubility of calcium carbonate, magnesium carbonate, and calcium sulfate, thus affecting the water-soluble salt content in soil leaching solution. The EC values of the C treatment with no CO₂ flux were relatively stable throughout the experimental period, whereas, the EC value of CO₂-addition treatments were affected by the injection time and concentration of CO₂ gas. The overall trend showed that the greater the concentration of ventilation, the greater the EC values. As the increasing ventilation time, the water soluble CO₂ in soil reached saturation, then the solubility of calcium carbonate,

magnesium carbonate and calcium sulfate in soil might be no longer affected by the gas ventilation. Therefore, the EC values of experimental treatments began to decline after time T2. *McFarland, Waldrop & Haw (2013)* indicated that the soil EC dropped sharply with an increasing CO₂ concentration. In China, *Zhang et al. (2016)* investigated the effect of CO₂ leakage on the soil properties and found that soil pH decreased with an increasing CO₂ leakage concentration, while the EC did not change.

Soil organic matter plays an important role in the terrestrial ecosystem, and it represents an important index to estimate the soil carbon storage and to evaluate the soil fertility and quality. In the current study, there was no significant correlation between organic matter and ventilation time or concentration. At first, the soil microorganisms could not adapt to the gas flux, so the organic matter content decreased sharply after aeration. However, the soil organic content increased from T1 to T4. This result indicates that microorganisms might adapt to the changing environment. However, when the gas flux was terminated, the soil organic matter content decreased sharply during the recovery period. It was speculated that the organic matter was degraded by surviving bacteria. In future research, we might apply ¹³C labeling to determine the carbon flow. Moreover, the organic matter content was positively correlated with the CO₂ flux, which supports the findings of *Sáenz de Miera et al. (2014)* and *Chen et al. (2016)*. These results also implied that injection of CO₂ flux decreased soil pH value, and influenced the microbial activities, then might result in the change of soil organic matter content.

An increased CO₂ concentration in soil can influence nitrogen content through affecting nitrogen fixation, mineralization, nitrification, denitrification, anaerobic ammonium oxidation, ammonia volatilization and other biochemical processes (*Yue et al., 2007; Chen et al., 2014*). On the whole, the nitrate nitrogen content increased with the increasing ventilation time and concentration from T0 to T4. A similar result has been represented by *McFarland, Waldrop & Haw (2013)*. However, a contrary result was shown by *Chen et al. (2016)* that the nitrate nitrogen content decreased sharply along the elevated CO₂ flux. In the non-planting soils, the organic matter might originate from microbial metabolic activities. The increasing CO₂ concentration may lead to the decreasing quantity and activity of microorganisms, thus affecting the organic matter content. After the CO₂ gas had been injected into the soil, an acidic and anaerobic environment might be formed. Then, the concentration of carbonate ions was potentially influenced and the change resulted in the concentration variation of ammonium ion (*Blagodatskaya et al., 2010; Phillips et al., 2012*). Moreover, the anaerobic environment may affect the anaerobic ammonium oxidation process. In the same way, the increase of soil CO₂ concentration might promote nitrification, which could increase the nitrate nitrogen concentration in the soil (*Ross et al., 2000; West et al., 2015*).

Phosphorus is one of the indispensable nutrient elements for plant growth and development. It is easily fixed in the soil, while the utilization rate is low. Although the change of Olsen-phosphorus and total phosphorus did not show an obvious correlation with the CO₂ concentration, their contents were lower under CO₂-addition treatments than under control treatment. Phosphorus is a necessary element for microbial metabolism. Some soil microorganisms may produce acidic substances through metabolism, then

dissolve some insoluble phosphates and apply them to their own metabolic processes. However, the phosphorus content in soil was related to soil organic matter content. As shown in Fig. 2C, the soil organic content increased with the increasing ventilation time from T1 to T4, whereas the change of Olsen-phosphorus and total phosphorus showed a decreasing trend (Figs. 2E and 2F). This result indicated that the soil phosphorus content might be negatively correlated with soil organic matter. It might be speculated that the CO₂ ventilation influenced the soil organic matter and bacterial activity, thus influencing the change of total phosphorus and Olsen-phosphorus contents. Sáenz de Miera *et al.*, (2014) reported that the total phosphorus was not well correlated with the CO₂ flux, the value was greater in the high and extreme site when compared with the control site, but they did not discuss the result. Fernández-Montiel, Pedescoll & Bécares (2016) reported that the Olsen phosphorus significantly increased as the CO₂ flux increased.

Soil enzyme activity impacts of high CO₂ concentration

Soil enzyme activities demonstrated different characteristics under CO₂ stress. The FDA hydrolysis showed a significant decrease, which indicates that the increasing CO₂ concentration restrained its activity. In addition, the inhibitory effects of CO₂ concentration on the activities of protease and dehydrogenase were not significant. The dehydrogenase activity was related to microbial respiratory metabolism. It was then speculated that the complex change of dehydrogenase activity might be related to the inhibition of soil microbial respiratory metabolism under increasing CO₂ stress. However, the increasing CO₂ concentration showed different degrees of stimulation to promote the activities of polyphenol oxidase and urease. The activity of polyphenol oxidase increased considerably, while the increase of urease activity fluctuated. Sinsabaugh (2010) mentioned that the chemical oxidation during the experiment assay can cause higher polyphenol oxidase values. However, McFarland, Waldrop & Haw (2013) demonstrated that there was no significant change in polyphenol oxidase activity under an increasing CO₂ flux. In another study, the researchers found that the elevated atmospheric CO₂ concentration enhanced the protease activity, while polyphenol oxidase and urease activities remained relatively unchanged (Shi *et al.*, 2016). Their study implied that elevated CO₂ concentration improved the availability of soil organic content, and it increased the soil microbial activity. Then, it induced the changes of soil enzyme activities. It can be speculated that the elevated soil CO₂ concentration decreased soil pH, organic matter content, total phosphorus and Olsen phosphorus content, resulting in the decrease of FDA hydrolysis activity. Moreover, it might be inferred that the increasing urease activity was associated with the elevated nitrate nitrogen content in the soil.

Phylogenetic impacts of elevated soil CO₂ concentration

Soil samples were collected for microbial analysis after one month of ventilation. The alpha and beta diversity analysis results showed that soil microbial community diversity decreased under CO₂ stress. However, the abundances of adapted dominant species increased with the increasing ventilation time and soil CO₂ concentration, while the microbial structure tended to be single. By means of species substitution, the soil microbial community

might adapt to the regional environment caused by high CO₂ concentration. It can be concluded that a local acid hypoxic soil environment might be formed under CO₂ stress. This represents a great survival challenge to a variety of bacteria and could cause many aerobic bacteria to die, thus leading to the decrease of soil bacterial richness and diversity. [Dunbar et al. \(2014\)](#) examined the impact of 11 years of elevated CO₂ on soil bacteria and contrasting results to the current study. They concluded that bacterial community in the 0–5 cm depth mineral soil demonstrated only minor changes after 11 years of elevated CO₂.

As [Fig. 4](#) shows, the distribution of phyla abundance differed under different CO₂ stresses, implying a possible impact of soil CO₂ flux. Results showed that dominant phylum of soil samples was *Proteobacteria*, which was promoted under the increasing soil CO₂ concentration. Meanwhile, [Šibanc et al. \(2014\)](#) reported a contrasting result which stated that the abundance of *Proteobacteria* decreased with increasing CO₂ concentration in spite of its dominant position. [Sáenz de Miera et al. \(2014\)](#) showed that a change in the abundance of the *Proteobacteria* phylum was less substantial under an increasing CO₂ concentration at a naturally occurring CO₂ gas vent. Moreover, it has rarely been mentioned in relevant literature that the abundances of genera *Methylophilus*, *Methylobacillus*, *Methylovorus* and *Methylothenera* belonging to the family *Methylophilaceae* (class *Betaproteobacteria*) sharply increased. Their growth substrates are generally with methyl groups. *Methylophilus* has been recognized as the aerobic methanol-oxidizing bacterium capable of degrading a variety of nitrogen-containing contaminants, such as methylamine, tetramethylammonium, and formamide ([Kolb, 2009](#)). Genera *Methylobacillus*, *Methylophilus* and *Methylovorus* have been assigned as the terrestrial obligate and restricted facultative *Methylobacteria*. The genus *Methylobacillus* was also presented to utilize methanol or methylamine as a sole source of carbon and energy ([Gogleva et al., 2011](#)). *Methylobacillus* sp. has been detected in riparian wetlands of Jiuduansha which could produce several oxidases that oxidize diverse compounds ([Strom, Dinarieva & Netrusov, 2001](#); [Hu et al., 2014](#)). Moreover, [Khan et al. \(2015\)](#) proposed that the *Methylophilus* could be used as a candidate for explosive detoxification in a nitrogen-deficient environment. Furthermore, genera *Methylobacillus*, *Methylothenera* and *Methylovorus* have been identified as methanotrophs which was related with methane metabolism under elevated CO₂ through the use of the whole genome metagenomic approach ([Bhattacharyya et al., 2016](#)). [Fernández-Montiel, Pedescoll & Bécares \(2016\)](#) also found the presence of *Methylocella palustris* in extreme flux which could produce CO₂ from CH₄ in an oxic environment. In contrast, *Variovorax* also showed a rising trend with an increasing CO₂ concentration. [Bablu & Sumathi \(2015\)](#) reported that the whole cells of *Variovorax* sp. BS1 had the ability to biodegrade dimethyl phthalate (DMP). Above all, it might be speculated that increasing soil CO₂ concentration might promote the increase of some methyl organic compounds in this experiment, which leads to a more suitable environment for these genera belonging to *Proteobacteria*. Furthermore, there is speculation that the injected CO₂ was utilized to produce methane which finally was consumed by these *Methanotrophs*.

The *Acidobacteria* played a significant role in the soil ecological processes, and this diverse phylum was distributed widely in various natural environments ([Tringe et al., 2005](#); [Janssen, 2006](#); [Liu et al., 2016](#)). Based on 16S rDNA gene analysis, this phylum generally occupied

10–50% of the total soil bacterial communities (*Fracchia et al., 2006; Penn et al., 2006; Lee & Cho, 2011*). *Liu et al. (2016)* demonstrated that the abundances of subgroups Gp4, Gp6 and Gp7 were significantly positive in relation to the soil pH value. Our study showed a similar result that the abundance of *Acidobacteria* decreased with the decreasing soil pH which implied that the relative abundance of *Acidobacteria* was significantly and positively correlated with soil pH. Moreover, there was a decline in the subgroups Gp4, Gp6 and Gp7 of *Acidobacteria* abundances. The shift of *Acidobacteria* should be related to the changes in soil pH values due to the soil acidification after CO₂ injection. Moreover, the phenomenon that the subgroups 4, 6, 7, 10, 11, 16, 17, 18, 22, and 25 were positively correlated with soil pH was presented in the broad-scale survey of *Acidobacteria* communities across 87 soils throughout North and South America by *Jones et al. (2009)*. The occurrence of this phenomenon might be related to the different subgroups of *Acidobacteria*, or perhaps even the same subgroup might show different responses to the soil environmental factors. *Sáenz de Miera et al. (2014)* presented that the soil pH decreased and the abundance of *Acidobacteria* _Gp4 and *Acidobacteria* _Gp6 sharply declined as the CO₂ flux increased. However, in another study *Sáenz de Miera et al. (2016)* showed that the relative abundance of *Acidobacteria* increased, and the abundances of classes Gp4 and Gp6 increased with a CO₂ injection in Cubillos del Sil soil. On the contrary, the abundances of classes Gp4 and Gp6 did not change considerably under high CO₂ injection in Hontomín soil. These phenomena indicated that it was impossible to determine whether the pH itself affected the shaping up of *Acidobacteria*, or whether the pH was indirectly related with the observed bacterial consortium changes through other environmental factors. Such environmental factors often co-vary together with the soil pH changes (*Rousk et al., 2010*). Previous studies have suggested that other soil parameters, such as soil C/N ratio, ammonium, phosphorus concentration, mineral element contents, and soil enzyme activity were also related with the *Acidobacteria* community composition (*Naether et al., 2012; Navarrete et al., 2013; Zhang et al., 2014*).

Sáenz de Miera et al. (2014) have reported that as the flux increased, the relative abundance of the *Chloroflexi* phylum increased while *Acidobacteria*, *Verrucomicrobia* and *Gemmatimonadetes* decreased. This result was slightly different from our result, which concluded that *Chloroflexi* phylum also decreased under CO₂ stress. Usually, leakage of CO₂ might result in an anaerobic environment, and the relative abundance of anaerobic bacteria such as *Bacteroidetes*, *Chloroflexi* and *Firmicutes* would show an increasing trend (*Šibanc et al., 2014*). However, *Bacteroidetes* did not show a significant relationship with the CO₂ injection, whereas *Chloroflexi* and *Firmicutes* decreased with an increasing CO₂ concentration. *Planctomycetacia* presented a notable increase after 14 days of CO₂ exposure from a controlled CO₂ sub-seabed leak in Ardmucknish Bay (*Tait et al., 2015*). In contrast, *Planctomycetes* phylum displayed a declining trend in our experiment.

Regarding the temporal variation shown in [Fig. 5A](#), the PCoA results indicated that the bacterial communities changed throughout this experiment. [Figure 5A](#) also displayed that the C treatment at T0 was far away from the C treatment at Ti ($i = 1, 2, 3$). Whereas, in the [Fig. 5B](#), there were just two major groups: one containing L and C samples and the other containing M, H, and E samples. As shown in [Fig. 5A](#), the percentage explanation of PC1

(principal component) was 62.4%, indicating the relation between the samples on this axis. However, the percentage of PC2 explanation is just 8.1%, so the results obtained on this axis were less reliable in interpretation. Therefore, we used the Fig. 5B to verify that the L and C samples had higher similarity. Another explanation might be that climatic characteristics such as temperature and precipitation might also affect the bacterial communities. Seasonal effects have been observed by several authors for different experiments (Sáenz de Miera *et al.*, 2014). This indicates that the separation between C1, C2 and C3 from C0 might be caused by climatic characteristics.

The CCA result indicated that the sample value of the CO₂-addition treatments increased along the arrow direction of nitrate content, urease and polyphenol oxidase. However, the other environmental variables such as pH, total phosphorus, and FDA hydrolysis showed the opposite trend. In addition, the pH value with the longest arrow was most closely related to CO₂ concentration, which suggested that this environmental factor was significant in inducing the differentiation of microbial community. All of these OTUs were closer to the L, M, H, and E groups than they were to the C group, which confirmed that these OTUs were significantly influenced by CO₂ concentration. This also indicated the changes of environmental factors such as nitrate nitrogen, urease and polyphenol oxidase, which might bring about a more suitable environment for these OTUs. Moreover, the correlations between pH and OTU7 (*Lysobacter*), OTU10 (*Streptophyta*), OTU13 (*Thermosulfurimonas*), OTU14 (*Gp6*), OTU15 (*Subdivision3_genera_incertae_sedis*), OTU16 (*Kofteria*) were significant and the relative abundance of *Acidobacteria_Gp6* was significantly and positively correlated with soil pH. Synthesizing Figs. 6A and 6B indicates that the arrows were longer for the OTU10 (*Streptophyta*) which belonged to *Cyanobacteria*, and the arrow was also longer for OTU16 (*Kofteria*) belonging to the phylum *Proteobacteria*. This implies that these two OTUs were important variables. Moreover, there was a high correlation between organic matter, dehydrogenase and OTU1 (*Sphingomonas*), OTU4 (*Massilia*), and OTU5 (*Janthinobacterium*). Unfortunately, all the OTUs had less correlation with protease, Olsen phosphorus, total phosphorus and FDA hydrolysis. The CCA results also indicated a gap between the control group and the groups under different CO₂ gas stress conditions. All of these results confirmed that the observed changes of *Proteobacteria* and *Acidobacteria* phyla were significantly impacted by CO₂ injection.

CONCLUSION

Ventilation of CO₂ gas significantly affected soil physiochemical properties and enzyme activity (lower pH and FDA hydrolysis activity, as well as increased nitrate nitrogen content and polyphenol oxidase activity), and resulted in lowered microbial diversity by favoring *Proteobacteria* phylum, such as *Methylophilus*, *Methylobacillus*, *Methylovorus*, and *Methylothera*. The promotion of these organisms might be linked to the increasing some methyl organic compounds in this experiment under CO₂ leakage stress. In contrast, the abundance of *Acidobacteria* decreased and the reason might be correlated with the comprehensive changes of soil parameters, such as soil phosphorus concentration, mineral element contents, and soil enzyme activity. Our data also indicated that the effect of

CO₂ leakage on the soil environment was significant, and that some of the effects were unrecoverable. Moreover, this study indicated that high CO₂ leakage can impact the soil microbial community and pose a risk to the ecological health of soil overlying CO₂ storage reservoirs.

ACKNOWLEDGEMENTS

The authors thank LetPub (<http://www.LetPub.com>) for its linguistic assistance during manuscript preparation.

ADDITIONAL INFORMATION AND DECLARATIONS

Funding

This work was supported by the Fundamental Research Funds for the Central Universities (2017XKQY070). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Grant Disclosures

The following grant information was disclosed by the authors:
Fundamental Research Funds for the Central Universities: 2017XKQY070.

Competing Interests

The authors declare there are no competing interests.

Author Contributions

- Jing Ma conceived and designed the experiments, performed the experiments, analyzed the data, wrote the paper, prepared figures and/or tables, reviewed drafts of the paper.
- Wangyuan Zhang performed the experiments, analyzed the data, prepared figures and/or tables.
- Shaoliang Zhang and Qiyang Feng contributed reagents/materials/analysis tools.
- Qianlin Zhu contributed reagents/materials/analysis tools, prepared figures and/or tables.
- Fu Chen conceived and designed the experiments, reviewed drafts of the paper.

Data Availability

The following information was supplied regarding data availability:

The raw data has been supplied as a [Supplementary File](#).

Supplemental Information

Supplemental information for this article can be found online at <http://dx.doi.org/10.7717/peerj.4024#supplemental-information>.

REFERENCES

- Acosta-Martínez V, Dowd S, Sun Y, Allen V. 2008.** Tag-encoded pyrosequencing analysis of bacterial diversity in a single soil type as affected by management and land use. *Soil Biology and Biochemistry* **40**:2762–2770 DOI [10.1016/j.soilbio.2008.07.022](https://doi.org/10.1016/j.soilbio.2008.07.022).
- Acosta-Martínez V, Dowd S, Sun Y, Wester D, Allen V. 2010.** Pyrosequencing analysis for characterization of soil bacterial populations as affected by an integrated livestock-cotton production system. *Applied Soil Ecology* **45**:13–25 DOI [10.1016/j.apsoil.2010.01.005](https://doi.org/10.1016/j.apsoil.2010.01.005).
- Andrzej R, Katarzyna S. 2013.** Environmental hazards caused by carbon capture and storage (CCS) technologies. *Polish Journal of Environmental Studies* **22**:205–211.
- Awasthi A, Singh M, Soni SK, Singh R, Kalra A. 2014.** Biodiversity acts as insurance of productivity of bacterial communities under abiotic perturbations. *ISME Journal* **8**:2445–2452 DOI [10.1038/ismej.2014.91](https://doi.org/10.1038/ismej.2014.91).
- Bablu P, Sumathi S. 2015.** Biodegradation of dimethyl phthalate ester using free cells, entrapped cells of *Variovorax* sp. BS1 and cell free enzyme extracts: a comparative study. *International Biodeterioration and Biodegradation* **97**:179–187 DOI [10.1016/j.ibiod.2014.11.004](https://doi.org/10.1016/j.ibiod.2014.11.004).
- Beaubien SE, Ciotoli G, Coombs P, Dictor MC, Krüger M, Lombardi S, Pearce JM, West JM. 2008.** The impact of a naturally occurring CO₂ gas vent on the shallow ecosystem and soil chemistry of a Mediterranean pasture (Latera, Italy). *International Journal of Greenhouse Gas Control* **2**:373–387 DOI [10.1016/j.ijggc.2008.03.005](https://doi.org/10.1016/j.ijggc.2008.03.005).
- Beulig F, Urich T, Nowak M, Trumbore SE, Gleixner G, Gilfillan GD, Fjelland KE, Küsel K. 2016.** Altered carbon turnover processes and microbiomes in soils under long-term extremely high CO₂ exposure. *Nature Microbiology* **1**:Article 15025 DOI [10.1038/nmicrobiol.2015.25](https://doi.org/10.1038/nmicrobiol.2015.25).
- Bhattacharyya P, Roy KS, Das M, Ray S, Balachandar D, Karthikeyan S, Nayak AK, Mohapatra T. 2016.** Elucidation of rice rhizosphere metagenome in relation to methane and nitrogen metabolism under elevated carbon dioxide and temperature using whole genome metagenomic approach. *Science of the Total Environment* **542**:886–898 DOI [10.1016/j.scitotenv.2015.10.154](https://doi.org/10.1016/j.scitotenv.2015.10.154).
- Blagodatskaya E, Blagodatsky S, Dorodnikov M, Kuzyakov Y. 2010.** Elevated atmospheric CO₂ increases microbial growth rates in soil: results of three CO₂ enrichment experiments. *Global Change Biology* **16**:836–848 DOI [10.1111/j.1365-2486.2009.02006.x](https://doi.org/10.1111/j.1365-2486.2009.02006.x).
- Chen F, Yang YJ, Ma YJ, Hou HP, Zhang SL, Ma J. 2016.** Effects of CO₂ leakage on soil bacterial community from simulated CO₂-EOR areas. *Environmental Science: Processes & Impacts* **18**:547–554 DOI [10.1039/c5em00571j](https://doi.org/10.1039/c5em00571j).
- Chen F, Zhang WY, Ma J, Yang YJ, Zhang SL, Chen R. 2017.** Experimental study on the effects of underground CO₂ leakage on soil microbial consortia. *International Journal of Greenhouse Gas Control* **63**:241–248 DOI [10.1016/j.ijggc.2017.05.017](https://doi.org/10.1016/j.ijggc.2017.05.017).
- Chen RR, Senbayram M, Blagodatsky S, Myachina O, Dittert K, Lin XG, Blagodatskaya E, Kuzyakov Y. 2014.** Soil C and N availability determine the priming effect:

- microbial N mining and stoichiometric decomposition theories. *Global Change Biology* **20**:2356–2367 DOI [10.1111/gcb.12475](https://doi.org/10.1111/gcb.12475).
- Duan BL, Zhang YB, Xu G, Chen J, Paquette A, Peng SM. 2015.** Long-term responses of plant growth, soil microbial communities and soil enzyme activities to elevated CO₂ and neighbouring plants. *Agricultural and Forest Meteorology* **213**:91–101 DOI [10.1016/j.agrformet.2015.06.017](https://doi.org/10.1016/j.agrformet.2015.06.017).
- Dumbrell AJ, Nelson M, Helgason T, Dytham C, Fitter AH. 2010.** Relative roles of niche and neutral processes in structuring a soil microbial community. *ISME Journal* **4**:337–345 DOI [10.1038/ismej.2009.122](https://doi.org/10.1038/ismej.2009.122).
- Dunbar J, Gallegos-Graves LV, Steven B, Mueller R, Hesse C, Zak DR, Kuske CR. 2014.** Surface soil fungal and bacterial communities in aspen stands are resilient to eleven years of elevated CO₂ and O₃. *Soil Biology and Biochemistry* **76**:227–234 DOI [10.1016/j.soilbio.2014.05.027](https://doi.org/10.1016/j.soilbio.2014.05.027).
- Fernández-Montiel I, Pedescoll A, Bécares E. 2016.** Microbial communities in a range of carbon dioxide fluxes from a natural volcanic vent in Campo de Calatrava, Spain. *International Journal of Greenhouse Gas Control* **50**:70–79 DOI [10.1016/j.ijggc.2016.04.017](https://doi.org/10.1016/j.ijggc.2016.04.017).
- Fernández-Montiel I, Touceda M, Pedescoll A, Gabilondo R, Prieto-Fernández A, Bécares E. 2015.** Short-term effects of simulated below-ground carbon dioxide leakage on a soil microbial community. *International Journal of Greenhouse Gas Control* **36**:51–59 DOI [10.1016/j.ijggc.2015.02.012](https://doi.org/10.1016/j.ijggc.2015.02.012).
- Fracchia L, Dohrmann AB, Martinotti MG, Tebbe CC. 2006.** Bacterial diversity in a finished compost and vermicompost: differences revealed by cultivation-independent analyses of PCR-amplified 16S rRNA genes. *Applied Microbiology and Biotechnology* **71**:942–952 DOI [10.1007/s00253-005-0228-y](https://doi.org/10.1007/s00253-005-0228-y).
- Frerichs J, Oppermann BI, Gwosdz S, Möller I, Herrmann M, Krüger M. 2013.** Microbial community changes at a terrestrial volcanic CO₂ vent induced by soil acidification and anaerobic microhabitats within the soil column. *FEMS Microbiology Ecology* **84**:60–74 DOI [10.1111/1574-6941.12040](https://doi.org/10.1111/1574-6941.12040).
- Gabilondo R, Bécares E. 2014.** The effects of natural carbon dioxide seepage on edaphic protozoan communities in Campo de Calatrava, Ciudad Real, Spain. *Soil Biology and Biochemistry* **68**:133–139 DOI [10.1016/j.soilbio.2013.09.028](https://doi.org/10.1016/j.soilbio.2013.09.028).
- Gogleva AA, Kaparullina EN, Doronina NV, Trotsenko YA. 2011.** *Methylobacillus arboreus* sp. nov. and *Methylobacillus gramineus* sp. nov. novel non-pigmented obligately methylophilic bacteria associated with plants. *Systematic and Applied Microbiology* **34**:477–481 DOI [10.1016/j.syapm.2011.03.005](https://doi.org/10.1016/j.syapm.2011.03.005).
- Guan YS. 1986.** *Soil enzyme and its study method [M]*. Beijing: China Agriculture Press, 274–320 [In Chinese].
- Horner-Devine MC, Carney KM, Bohannan B. 2004.** An ecological perspective on bacterial biodiversity. *Proceedings of the Royal Society of London. Series B: Biological Sciences* **271**:113–122 DOI [10.1098/rspb.2003.2549](https://doi.org/10.1098/rspb.2003.2549).
- Hu Y, Wang L, Tang YS, Li YL, Chen JH, Xi XF, Zhang YN, Fu XH, Wu JH, Sun Y. 2014.** Variability in soil microbial community and activity between coastal and riparian

- wetlands in the Yangtze River estuary—potential impacts on carbon sequestration. *Soil Biology and Biochemistry* **70**:221–228 DOI [10.1016/j.soilbio.2013.12.025](https://doi.org/10.1016/j.soilbio.2013.12.025).
- Janssen PH. 2006.** Identifying the dominant soil bacterial taxa in libraries of 16S rRNA and 16S rRNA genes. *Applied and Environmental Microbiology* **72**:1719–1728 DOI [10.1128/AEM.72.3.1719-1728.2006](https://doi.org/10.1128/AEM.72.3.1719-1728.2006).
- Jones RT, Robeson MS, Lauber CL, Hamady M, Knight R, Fierer N. 2009.** A comprehensive survey of soil acidobacterial diversity using pyrosequencing and clone library analyses. *ISME Journal* **3**:442–453 DOI [10.1038/ismej.2008.127](https://doi.org/10.1038/ismej.2008.127).
- Khan MI, Lee J, Yoo K, Kim S, Park J. 2015.** Improved TNT detoxification by starch addition in a nitrogen-fixing *Methylophilus*-dominant aerobic microbial consortium. *Journal of Hazardous Materials* **300**:873–881 DOI [10.1016/j.jhazmat.2015.08.032](https://doi.org/10.1016/j.jhazmat.2015.08.032).
- Ko D, Yoo G, Yun ST, Chung H. 2016.** Impacts of CO₂ leakage on plants and microorganisms: a review of results from CO₂ release experiments and storage sites. *Greenhouse Gases: Science and Technology* **6**:319–338 DOI [10.1002/ghg.1593](https://doi.org/10.1002/ghg.1593).
- Kolb S. 2009.** Aerobic methanol-oxidizing bacteria in soil. *FEMS Microbiology Letters* **300**:1–10 DOI [10.1111/j.1574-6968.2009.01681.x](https://doi.org/10.1111/j.1574-6968.2009.01681.x).
- Krüger M, Jones D, Frerichs J, Oppermann B, West J, Coombs P, Green K, Barlow K, Lister R, Shaw R, Strutt M, Möller I. 2011.** Effects of elevated CO₂ concentrations on the vegetation and microbial populations at a terrestrial CO₂ vent at Laacher See, Germany. *International Journal of Greenhouse Gas Control* **5**:1093–1098 DOI [10.1016/j.ijggc.2011.05.002](https://doi.org/10.1016/j.ijggc.2011.05.002).
- Lee SH, Cho JC. 2011.** Group-specific PCR primers for the phylum Acidobacteria designed based on the comparative analysis of 16S rRNA gene sequences. *Journal of Microbiological Methods* **86**:195–203 DOI [10.1016/j.mimet.2011.05.003](https://doi.org/10.1016/j.mimet.2011.05.003).
- Lemos LN, Fulthorpe RR, Triplett EW, Roesch LFW. 2011.** Rethinking microbial diversity analysis in the high throughput sequencing era. *Journal of Microbiological Methods* **86**:42–51 DOI [10.1016/j.mimet.2011.03.014](https://doi.org/10.1016/j.mimet.2011.03.014).
- Li Q, Song R, Liu X, Liu G, Sun Y. 2016.** Monitoring of carbon dioxide geological utilization and storage in China: a review. *Acid Gas Extraction for Disposal and Related Topics* **33**:1–358.
- Liu JJ, Sui YY, Yu ZH, Yao Q, Shi Y, Chu HY, Jin J, Liu XB, Wang GH. 2016.** Diversity and distribution patterns of acidobacterial communities in the black soil zone of northeast China. *Soil Biology and Biochemistry* **95**:212–222 DOI [10.1016/j.soilbio.2015.12.021](https://doi.org/10.1016/j.soilbio.2015.12.021).
- Mandić-Mulec I, Gorenc K, Gams Petrisic M, Faganeli J, Ogrinc N. 2012.** Methanogenesis pathways in a stratified eutrophic alpine lake (Lake Bled, Slovenia). *Limnology and Oceanography* **57**:868–880 DOI [10.4319/lo.2012.57.3.0868](https://doi.org/10.4319/lo.2012.57.3.0868).
- Maček I, Dumbrell AJ, Nelson M, Fitter AH, Vodnik D, Helgason T. 2011.** Local adaptation to soil hypoxia determines the structure of an arbuscular mycorrhizal fungal community in roots from natural CO₂ springs. *Applied and Environmental Microbiology* **77**:4770–4777 DOI [10.1128/AEM.00139-11](https://doi.org/10.1128/AEM.00139-11).

- McFarland JW, Waldrop MP, Haw M. 2013.** Extreme CO₂ disturbance and the resilience of soil microbial communities. *Soil Biology and Biochemistry* **65**:274–286 DOI [10.1016/j.soilbio.2013.04.019](https://doi.org/10.1016/j.soilbio.2013.04.019).
- Morales SE, Holben WE. 2014.** Simulated geologic carbon storage leak reduces bacterial richness and alters bacterial community composition in surface soil. *Soil Biology and Biochemistry* **76**:286–296 DOI [10.1016/j.soilbio.2014.05.018](https://doi.org/10.1016/j.soilbio.2014.05.018).
- Naether A, Foesel BU, Naegele V, Wust PK, Weinert J, Bonkowski M, Alt F, Oelmann Y, Polle A, Lohaus G, Gockel S, Hemp A, Kalko EKV, Linsenmair KE, Pfeiffer S, Renner S, Schoning I, Weisser WW, Wells K, Fischer M, Overmann J, Friedrich MW. 2012.** Environmental factors affect acidobacterial communities below the subgroup level in grassland and forest soils. *Applied and Environmental Microbiology* **78**:7398–7406 DOI [10.1128/AEM.01325-12](https://doi.org/10.1128/AEM.01325-12).
- Naidu VSGR, Sarathambal C. 2012.** Effect of elevated and atmosphere CO₂ on soil microorganisms, soil respiration and enzyme activities. *Journal of Soil Biology & Ecology* **32**:82–92.
- Navarrete AA, Kuramae EE, De Hollander M, Pijl AS, Van Veen JA, Tsai SM. 2013.** Acidobacterial community responses to agricultural management of soybean in Amazon forest soils. *FEMS Microbiology Ecology* **83**:607–621 DOI [10.1111/1574-6941.12018](https://doi.org/10.1111/1574-6941.12018).
- Norman RJ, Edberg JC, Stucki JW. 1985.** Determination of nitrate in soil extracts by dual-wavelength ultraviolet spectrophotometry. *Soil Science Society of America Journal* **49**:1182–1185 DOI [10.2136/sssaj1985.03615995004900050022x](https://doi.org/10.2136/sssaj1985.03615995004900050022x).
- Oppermann BI, Michaelis W, Blumenberg M, Frerichs J, Schulz HM, Shippers A, Beaubien SE, Krüger M. 2010.** Soil microbial community changes as a result of long-term exposure to a natural CO₂ vent. *Geochimica et Cosmochimica Acta* **74**:2697–2716 DOI [10.1016/j.gca.2010.02.006](https://doi.org/10.1016/j.gca.2010.02.006).
- Österreicher Cunha P, Molinaro BS, Feijó IVA, Vargas Jr EA, Guimarães JRD. 2015.** Experimental evaluation of CO₂ percolation effects on subsurface soil microbiota. *International Journal of Greenhouse Gas Control* **32**:135–146 DOI [10.1016/j.ijggc.2014.11.009](https://doi.org/10.1016/j.ijggc.2014.11.009).
- Patil RH, Colls JJ, Steven MD. 2010.** Effects of CO₂ gas as leaks from geological storage sites on agro-ecosystems. *Energy* **35**:4587–4591 DOI [10.1016/j.energy.2010.01.023](https://doi.org/10.1016/j.energy.2010.01.023).
- Penn K, Wu DY, Eisen JA, Ward N. 2006.** Characterization of bacterial communities associated with deep-sea corals on Gulf of Alaska seamounts. *Applied and Environmental Microbiology* **72**:1680–1683 DOI [10.1128/AEM.72.2.1680-1683.2006](https://doi.org/10.1128/AEM.72.2.1680-1683.2006).
- Phillips RP, Meier IC, Bernhardt ES, Grandy AS, Wickings K, Finzi AC. 2012.** Roots and fungi accelerate carbon and nitrogen cycling in forests exposed to elevated CO₂. *Ecology Letters* **15**:1042–1049 DOI [10.1111/j.1461-0248.2012.01827.x](https://doi.org/10.1111/j.1461-0248.2012.01827.x).
- Prosser JI. 2007.** Microorganisms cycling soil nutrients and their diversity. In: Van Elsas JD, Jansson JK, Trevors JT, eds. *Modern soil microbiology*. Second edition. Boca Raton: CRC Press, 237–261.
- Ross DJ, Tate KR, Newton PCD, Wilde RH, Clark H. 2000.** Carbon and nitrogen pools and mineralization in a grassland gley soil under elevated carbon dioxide at a natural CO₂ spring. *Global Change Biology* **6**:779–790 DOI [10.1046/j.1365-2486.2000.00357.x](https://doi.org/10.1046/j.1365-2486.2000.00357.x).

- Rousk J, Bååth E, Brookes PC, Lauber CL, Lozupone C, Caporaso G, Knight R, Fierer N. 2010. Soil bacterial and fungal communities across a pH gradient in an arable soil. *ISME Journal* 4:1340–1351 DOI 10.1038/ismej.2010.58.
- Sáenz de Miera LE, Arroyo P, Calabuig EL, Falagán J, Ansola G. 2014. High-throughput sequencing of 16S RNA genes of soil bacterial communities from a naturally occurring CO₂ gas vent. *International Journal of Greenhouse Gas Control* 29:176–184 DOI 10.1016/j.ijggc.2014.08.014.
- Sáenz de Miera LE, Arroyo P, Calabuig EL, Ansola G. 2016. Effects of varying CO₂ flows on bacterial communities in mesocosms created from two soils. *International Journal of Greenhouse Gas Control* 46:205–214 DOI 10.1016/j.ijggc.2016.01.013.
- Sarathambal C, Rathore M, Jaggi D, Kumar B. 2016. Response of soil enzymes to elevated CO₂ and temperature in weeds associated with rice-wheat cropping system. *Indian Journal of Weed Science* 48(1):29–32 DOI 10.5958/0974-8164.2016.00006.X.
- Sardans J, Penuelas J, Estiarte M. 2008. Changes in soil enzymes related to C and N cycle and in soil C and N content under prolonged warming and drought in a Mediterranean shrubland. *Applied Soil Ecology* 39:223–235 DOI 10.1016/j.apsoil.2007.12.011.
- Schloss P, Westcott SL, Ryabin T, Hall JR, Hartmann M, Hollister EB, Lesniewski RA, Oakley BB, Parks DH, Robinson CJ, Sah JW, Stres B, Thallinger GG, Horn DJV, Weber CF. 2009. Introducing mothur: open-source, platform-independent, community-supported software for describing and comparing microbial communities. *Applied and Environmental Microbiology* 75:7537–7541 DOI 10.1128/AEM.01541-09.
- Shi CE, Ai FX, Wang CR, Yan SB, Che YC. 2016. Effects of elevated atmospheric CO₂ and O₃ on soil enzyme activities and microbial biomass. *Journal of Agro-Environment Science* 35(6):1103–1109 [In Chinese].
- Šibanc N, Dumbrell AJ, Mandić-Mulec I, Maček I. 2014. Impacts of naturally elevated soil CO₂ concentrations on communities of soil archaea and bacteria. *Soil Biology and Biochemistry* 68:348–356 DOI 10.1016/j.soilbio.2013.10.018.
- Sinsabaugh RL. 2010. Phenol oxidase, peroxidase and organic matter dynamics of soil. *Soil Biology and Biochemistry* 42:391–404 DOI 10.1016/j.soilbio.2009.10.014.
- Strom EV, Dinarieva TY, Netrusov AI. 2001. *Methylobacillus flagellatus* KT contains a novel cbo-type cytochrome oxidase. *FEBS Letters* 505:109–112 DOI 10.1016/S0014-5793(01)02795-8.
- Tait K, Stahl H, Taylor P, Widdicombe S. 2015. Rapid response of the active microbial community to CO₂ exposure from a controlled sub-seabed CO₂ leak in Ardmucknish Bay (Oban, Scotland). *International Journal of Greenhouse Gas Control* 38:171–181 DOI 10.1016/j.ijggc.2014.11.021.
- Taylor JD, Ellis R, Milazzo M, Hall-Spencer JM, Cunliffe M. 2014. Intertidal epilithic bacteria diversity changes along a naturally occurring carbon dioxide and pH gradient. *FEMS Microbiology Ecology* 89:670–678 DOI 10.1111/1574-6941.12368.
- Tian D, Xin M, Zha LS, Yang W, Zou X, Hou X. 2013. Review of impact of CO₂ leakage from geological storage on near-surface terrestrial ecological system. *Journal of Ecology and Rural Environment* 29:137–145.

- Tringe SG, Von Mering C, Kobayashi A, Salamov AA, Chen K, Chang HW, Podar M, Short JM, Mathur EJ, Detter JC, Bork P, Hugenholtz P, Rubin EM. 2005. Comparative metagenomics of microbial communities. *Science* 308:554–557 DOI 10.1126/science.1107851.
- Van Der Heijden MGA, Bardgett RD, Van Straalen NM. 2008. The unseen majority: soil microbes as drivers of plant diversity and productivity in terrestrial ecosystems. *Ecology Letters* 11:296–310 DOI 10.1111/j.1461-0248.2007.01139.x.
- West JM, Jones DG, Annunziatellis A, Barlow TS, Beaubien SE, Bond A, Breward N, Coombs P, De Angelis D, Gardner A, Gemeni V, Graziani S, Green KA, Gregory S, Gwosdz S, Hannis S, Kirk K, Koukouzas N, Krüger M, Libertini S, Lister TR, Lombardi S, Metcalfe R, Pearce JM, Smith KL, Steven MD, Thatcher K, Ziogou F. 2015. Comparison of the impacts of elevated CO₂ soil gas concentrations on selected European terrestrial environments. *International Journal of Greenhouse Gas Control* 42:357–371 DOI 10.1016/j.ijggc.2015.07.020.
- West JM, Pearce JM, Bentham M, Maul P. 2005. Environmental issues and the geological storage of CO₂. *European Environment* 15:250–259 DOI 10.1002/eet.388.
- West JM, Pearce JM, Coombs P, Ford JR, Scheib C, Colls JJ, Smith KL, Steven MD. 2009. The impact of controlled injection of CO₂ on the soil ecosystem and chemistry of an English lowland pasture. *Energy Procedia* 1:1863–1870 DOI 10.1016/j.egypro.2009.01.243.
- Zhang BH, Zhang Q, Geng CX, Jing JJ. 2016. Influence on physical and chemical properties of soil for the leakage of carbon dioxide during geological storage. *Guangzhou Chemical Industry* 44(7):156–159.
- Zhang Y, Cong J, Lu H, Li G, Qu Y, Su X, Zhou J, Li D. 2014. Community structure and elevational diversity patterns of soil Acidobacteria. *Journal of Environmental Sciences (China)* 26:1717–1724 DOI 10.1016/j.jes.2014.06.012.
- Yue J, Shi Y, Zheng X, Huang G, Zhu J. 2007. The influence of free-air CO₂ enrichment on microorganisms in paddy soil in the rice-growing season. *Applied Soil Ecology* 35:154–162 DOI 10.1016/j.apsoil.2006.05.005.
- Zhou X, Apple ME, Dobeck LM, Cunningham AB, Spangler LH. 2013. Observed response of soil O₂ concentration to leaked CO₂ from an engineered CO₂ leakage experiment. *International Journal of Greenhouse Gas Control* 16:116–128 DOI 10.1016/j.ijggc.2013.03.005.
- Ziogou F, Gemenia V, Koukouzasa N, De Angelis D, Libertini S, Beaubien SE, Lombardi S, West JM, Jones DG, Coombs P, Barlow TS, Gwosdz S, Krüger M. 2013. Potential environmental impacts of CO₂ leakage from the study of natural analogue sites in Europe. *Energy Procedia* 37:3521–3528 DOI 10.1016/j.egypro.2013.06.245.