

# Lawrence Berkeley Laboratory

UNIVERSITY OF CALIFORNIA

Submitted to Physical Review Letters

OBSERVATION OF HIGH MOMENTUM PROTONS FROM LIMITING  
TARGET FRAGMENTATION

J. V. Geaga, S. A. Chessin, J. Y. Grossiord, J. W. Harris,  
D. L. Hendrie, L. S. Schroeder, R. N. Treuhaft, and  
K. Van Bibber

April 1980


**TWO-WEEK LOAN COPY**

*This is a Library Circulating Copy  
which may be borrowed for two weeks.  
For a personal retention copy, call  
Tech. Info. Division, Ext. 6782.*

RECEIVED  
LAWRENCE  
BERKELEY LABORATORY

JUN 20 1980

LIBRARY AND  
DOCUMENTS SECT



LBL-10265 e.2



## DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

OBSERVATION OF HIGH MOMENTUM PROTONS  
FROM LIMITING TARGET FRAGMENTATION\*

J. V. Geaga, S. A. Chessin, J. Y. Grossiord,\*\* J. W. Harris,  
D. L. Hendrie, L. S. Schroeder, R. N. Treuhaft, K. Van Bibber†

Nuclear Science Division  
Lawrence Berkeley Laboratory  
University of California  
Berkeley, CA 94720

ABSTRACT

We report on measurements of the inclusive distributions of protons produced at  $180^\circ$  in the momentum range  $0.3 \leq p \leq 1.0$  GeV/c. Proton, alpha, carbon, and argon beams in the range of kinetic energies  $0.4 \leq T \leq 2.1$  GeV/n (4.89 GeV for protons) were incident on C, Al, Cu, Sn, and Pb targets. The dependences of the cross sections on the projectile and target mass and on the incident energy are presented. Limiting behavior is found at energies above 1. - 2. GeV/n. Features suggestive of nuclear correlations are discussed.

---

\* This work is supported by the Nuclear Physics Division of the U. S. Department of Energy under Contract No. W-7405-ENG-48.

\*\* Present address: Institut de Physique Nucleaire de Lyon, France.

† Present address: Department of Physics, Stanford University.

Nuclear collisions are studied with the hope of learning details of nuclear structure or reaction mechanisms in the collisions. Precise measurement of  $(e,e')$ ,  $(e,e'p)$ ,  $(p,2p)$ , and  $(\gamma,p)$ <sup>(1)</sup> reactions have provided much information on the distributions of nucleon momenta inside nuclei up to 300-400 MeV/c. Predictions using known single particle wavefunctions are in good agreement with the data. Attempts to relate the large momenta observed in  $(\gamma,p)$ <sup>(2)</sup> and backward  $(p,p')$ <sup>(3)</sup> reactions directly to high internal momenta indicate the presence of high momentum components in nuclei far above those expected from single particle wavefunctions. A quasi-elastic description was used<sup>(4)</sup> to relate backward momenta from the  $(p,p')$  reactions to internal momenta. It was stated that production from nuclei could result only from the internal motion of nucleons since backward production is forbidden in free nucleon-nucleon collisions. The difficulties involved in extracting information on high internal momenta, which are associated with deviations from mean field motion in the nucleus, have been pointed out by Gottfried.<sup>(5)</sup> In addition, Amado and Woloshyn<sup>(6)</sup> have shown that final state interactions do not permit a simple interpretation of the backward momenta in the  $(p,p')$  reactions.<sup>(3)</sup> The determination of large internal momenta is still a fundamental experimental and theoretical problem. Nevertheless, the high momenta observed in these types of reactions provide valuable information on the production mechanisms, and possibly on internal momentum distributions.

We have recently concluded a systematic study of the inclusive distributions of high momentum protons produced at  $180^\circ$  using the full range of beams and energies available from the Bevalac facility. Earlier measurements<sup>(7)</sup> at angles  $\theta_{lab} \geq 120^\circ$  using high energy  $\nu$ ,  $\gamma$ ,  $\pi$ , and  $p$  beams revealed that the shapes of the spectra of protons in the momentum interval  $0.3 \leq p \leq 1.0$  GeV/c were insensitive to the type of the projectile and its

energy. We find that at the lower incident energies, the cross sections rise rapidly with increasing bombarding energy. However, we also find that beyond 1.-2. GeV/n, the distributions in the momentum region  $0.4 \leq p \leq 1.0$  GeV/c become relatively insensitive to the further increase of the incident energy. In comparison, projectile fragmentation studies found that fragment distributions for  $p \leq 400$  MeV/c in the projectile frame already became limiting somewhere between beam energies 0.4 and 1.05 GeV/n.<sup>(8)</sup> Projectile fragmentation studies together with target fragmentation measurements provide systematic data spanning the two momentum ranges  $0. \leq p \leq .4$  GeV/c and  $0.4 \leq p \leq 1.0$  GeV/c. We find that the limiting proton distributions encompassing these two momentum ranges are approximated better by two distinct overlapping Gaussians rather than by a single Gaussian. In this parametrization, both Gaussians have a mean at zero momentum. The second Gaussian, which has a larger dispersion, starts to dominate in the region 300-400 MeV/c. Results from theoretical investigations suggest that the second Gaussian is a manifestation of nuclear correlations.

The experiment was performed at the LBL Bevalac. The beams used were (0.8, 1.05, 2.1, 3.5, 4.89) GeV protons, (1.05, 2.1) GeV/n alphas, (0.4, 1.05, 2.1) GeV/n carbon, and (1.05, 1.83) GeV/n argon. The targets were C, Al, Cu, Sn, and Pb. Protons at  $180^\circ$  were measured in the momentum range  $0.3 \leq p \leq 1.0$  GeV/c with a magnetic spectrometer. Details of the apparatus are reported in Ref. 9. We estimate the overall absolute uncertainties in the inclusive cross sections to be  $\pm 15$  percent.

The energy dependence of the cross sections is shown in Fig. 1. Results from other experiments at 90 MeV,<sup>(10)</sup> 600 MeV,<sup>(3)</sup> 7.71 GeV,<sup>(11)</sup> and 400 GeV<sup>(12)</sup> have been included. The data indicates limiting behavior of the distributions in the range of momenta  $0.4 \leq p \leq 1.0$  GeV/c starting between 1.0

and 2.0 GeV/n. A slight rise is found however in the 400 GeV data measured at  $160^\circ$ . The spectrum at 90 MeV, commonly attributed to pre-equilibrium decay<sup>(13)</sup> at low energies, has an elastic kinematic limit at 392 MeV/c. A lack of dependence of the proton cross sections on the incident energy is exhibited in data for all targets with proton beams having kinetic energies  $T \geq 2.0$  GeV, strongly suggestive that limiting target fragmentation<sup>(14)</sup> has set in. The hypothesis of limiting fragmentation states that the cross sections for the production of target fragments in the lab and projectile fragments in the projectile rest frame become independent of incident energy. The energy ranges spanned by the heavier projectiles are not as extensive but the features of the data are similar.

The dependence of the proton production cross sections on projectile mass for a carbon target at an incident energy of 1.05 GeV/n is shown in Fig. 2. The shapes of the proton spectra in the momentum range  $0.4 \leq p \leq 1.0$  GeV/c are seen to be roughly independent of projectile mass. Plotted in the same figure (in the projectile rest frame) are data from Papp<sup>(15)</sup> for  $C+C \rightarrow p(2.5^\circ)$  at 1.05 GeV/n. The break found in the  $C+C \rightarrow p$  data at about 350 MeV/c is similar to that found in extracted momentum distributions from  $(\gamma, p)$ <sup>(2)</sup> data. Recent theoretical investigations<sup>(16-18)</sup> show that the internal momentum distributions in nuclei can be approximated by two overlapping Gaussians when short range correlations are included. The low momentum Gaussian is dominated by the single particle characteristics of the momentum distributions and the second Gaussian emerges when short range correlations are considered. The solid line is a double Gaussian parametrization of the  $C+C \rightarrow p$  data. If the internal momentum distributions are indeed correctly specified by the models, then one is led to conjecture that the shapes of the measured distributions in the momentum region  $0.4 \leq p \leq 1.0$  GeV/c are a consequence of the presence of nuclear correlations.

In order to further study the dependence of these cross sections on projectile and target mass and energy, we have parametrized our data in the momentum range  $0.4 \leq p \leq 1.0$  GeV/c in the form  $d\sigma/dp^3 = \sigma_T / (\sqrt{2\pi})^3 \alpha^3 \times e^{-(p^2/2\alpha^2)}$  (where  $\sigma_T = \int (d\sigma/dp^3) d^3p$ ). The fits are good for incident kinetic energies  $T \geq 1.05$  GeV/n having typical uncertainties of  $\pm 5$  MeV/c for  $\alpha$  and  $\pm 10$  percent for  $\sigma_T$ . The dependence of  $\alpha$  on the target and projectile masses is shown in Fig. 3(a). The mass dependences are relatively weak. The dependence on incident energy is more dramatic and the limiting nature of the cross sections is reflected in this dependence. A limiting value of  $220 \pm 10$  MeV/c is indicated by data at various backward angles and energies above 5 GeV.<sup>(7,11,12)</sup> For comparison, the distributions in the low momentum region  $p \leq 400$  MeV/c, where one would expect the first Gaussian, should be well described by the formulation of Goldhaber<sup>(19)</sup> which has been successful in describing fragment momenta from projectile fragmentation. Employing Fermi momentum values from quasi-elastic electron scattering measurements,<sup>(20)</sup> one gets correspondingly smaller values of  $\alpha$  in this low momentum region ranging from 76 MeV/c for Li to 119 MeV/c for Pb. For the higher momentum region, the dependence of  $\sigma_T$  on the masses at 1.05 GeV/n incident energy is shown in Fig. 3(b). An approximate  $A_p^{2/3} A_t^{4/3}$  ( $p = \text{proj}$ ,  $t = \text{tgt}$ ) dependence is found for  $\sigma_T$  at 1.05 GeV/n and 2.1 GeV/n. The  $A_p$  dependence is expected in a limiting fragmentation model<sup>(21)</sup> where the projectile is seen as a contracted disk by the target nucleus. In such a model, one could interpret the measured momenta as reflecting information of the target. The  $A_t$  dependence argues against incoherent collisions and indicates the involvement of more than one target nucleon in backward nucleon production. Inclusive proton data in the fragmentation region from  $p+d$  and  $p+\alpha$  reactions<sup>(22,23)</sup> show a similar break in the distributions at about 300 MeV/c. This would argue against a multiple scattering cascade as the dominant mechanism, at least for light

nuclei. Definitive theoretical work on the contributions of multiple scattering is, however, still necessary to justify this conclusion.

It has been noted<sup>(8)</sup> that results of measurements of fast backward target fragments and fast forward projectile fragments indicate the presence of short range structure. Of the various approaches<sup>(4,24-33)</sup> to parametrize these data, very attractive are models which include scattering with correlated nucleons. A model by Fujita and Hüfner<sup>(24)</sup> includes both initial correlations between nucleons in the target and final correlations between two nucleons. The final state interaction with the rest of the  $A-2$  nucleons is however neglected. Fujita has extended this model to include multiple correlations.<sup>(34)</sup> The improved model describes well the incident energy dependence of backward proton spectra in the region  $0.4 \leq p \leq 1.0$  GeV/c for the intermediate energy range 0.6 to 1.0 GeV. Theoretical work by Frankfurt and Strikman<sup>(29)</sup> and Yukawa and Furui<sup>(30)</sup> indicate that backward spectra induced by high energy probes are mainly composed of spectator nucleons from the breakup of correlated pairs in the nucleus. As such, these backward nucleons potentially reflect direct information of the nuclear wavefunction.

In conclusion, we find that our data spans a transition region wherein the inclusive distributions of protons in the momentum range  $0.4 \leq p \leq 1.0$  GeV/c are found to exhibit limiting behavior. The data, together with recent theoretical descriptions of internal momentum distributions in nuclei and reaction mechanisms involving correlated nucleons, strongly suggest that the limiting shapes of the measured spectra are related to the presence of correlations in the nucleus.

We wish to thank the Bevatron crew for support during the measurements, and Miklos Gyulassy for helpful theoretical discussions. We also wish to thank Craig Wuest and Gloria Laskowich for their participation during some stages of the experiment.



REFERENCES


- ( 1) D. J. S. Findlay and R. O. Owens, Nucl. Phys. A 292, 53 (1977) and references therein.
- ( 2) D. J. S. Findlay et al., Phys. Lett. 74B, 305 (1978).
- ( 3) S. Frankel et al., Phys. Rev. Lett. 36, 642 (1976).
- ( 4) S. Frankel, Phys. Rev. Lett. 38, 1338 (1977).
- ( 5) K. Gottfried, Ann. Phys. 21, 29 (1963).
- ( 6) R. D. Amado and R. M. Woloshyn, Phys. Lett. 69B, 400 (1977).
- ( 7) G. A. Leksin, Int. Conf. High Energy Physics, ITEP-147 (1976) and references therein.
- ( 8) A. S. Goldhaber and H. H. Heckman, Ann. Rev. Nucl. Part. Sci., 28, 161 (1978) and references therein.
- ( 9) L. S. Schroeder et al., Phys. Rev. Lett. 43, 1787 (1979).
- (10) J. R. Wu et al., Phys. Rev. C 19, 698 (1979).
- (11) A. M. Baldin, Preprint JINR-P1-11302, Dubna (1978).
- (12) Y. D. Bayukov et al., Phys. Rev. C 20, 764 (1979).
- (13) M. Blann, Ann. Rev. Nucl. Sci. 25, 123 (1975).
- (14) J. Benecke et al., Phys. Rev. 188, 2159 (1969).
- (15) J. Papp, LBL-3633, Berkeley (1975), Ph.D. Thesis (unpubl.).
- (16) J. G. Zabolitsky and W. Ey, Phys. Lett. 76B, 527 (1978).
- (17) O. Bohigas and S. Stringari, Preprint ORSAY IPNO/TH 79-59, (1979).
- (18) J. W. Van Orden et al., Preprint U. of MD. TR 80-046, (1979).
- (19) A. S. Goldhaber, Phys. Lett. 53B, 306 (1974).
- (20) E. J. Moniz et al., Phys. Rev. Lett. 26, 445 (1971).
- (21) Perl, M. L. (1974), High Energy Hadron Physics, John Wiley & Sons, New York, p. 201.
- (22) A. M. Baldin et al., Preprint JINR-P1-11168, Dubna (1977).

- (23) L. M. Anderson, LBL-6769, Berkeley (1977), Ph.D. Thesis (unpubl.).
- (24) T. Fujita and J. Hüfner, Nucl. Phys. A 314, 317 (1979).
- (25) G. Berlad et al., Preprint Technion PH-79-78 (1979).
- (26) V. V. Burov et al., Phys. Lett. 67B, 46 (1977).
- (27) T. Fujita, Phys. Rev. Lett. 39, 174 (1977).
- (28) I. A. Schmidt and R. Blankenbecler, Phys. Rev. D 15, 3321 (1977).
- (29) L. L. Frankfurt and M. I. Strikman, Phys. Lett. 83B, 407 (1979).
- (30) T. Yukawa and S. Furui, Phys. Rev. C 20, 2316 (1979).
- (31) R. H. Landau and M. Gyulassy, Phys. Rev. C 19, 149 (1979).
- (32) H. B. Mathis and Meng Ta-chung, Phys. Rev. C 18, 952 (1978).
- (33) V. K. Lukyanov and A. I. Titov, Sov. J. Part. Nucl. 10, 321 (1979).
- (34) T. Fujita, Nucl. Phys. A 324, 409 (1979).

FIGURE CAPTIONS


- Fig. 1. Proton inclusive distributions;  $\Delta, \blacklozenge, \circ$  this expt.;  $\square$  Ref. 12;  $\nabla$  Ref. 11; — Ref. 3; ---- Ref. 10.
- Fig. 2. Proton inclusive distributions;  $\bullet$  Ref. 15, plotted in the rest frame of the projectile;  $\nabla, \diamond, \square$  this expt., plotted in the lab frame; — double Gaussian representation of  $C + C \rightarrow p$  data (solid points) with values of parameters indicated in N(P).
- Fig. 3. (a)  $\blacklozenge$  energy dependence of  $\alpha$ ,  $\circ$  target dependence of  $\alpha$ ,  $\square$  projectile dependence of  $\alpha$  where A denotes both  $A_p$  and  $A_t$  (b) plot of parameter  $\sigma_T$  vs.  $A_p^{2/3} A_t^{4/3}$ , solid line shows trend expected if dependence of  $\sigma_T$  is  $A_p^{2/3} A_t$ .







XBL 7912 - 13375

Fig. 1



XBL 7912 - 13374

Fig. 2



XBL 7912-13376

Fig. 3