


Contents lists available at [ScienceDirect](http://ScienceDirect.com)

Neurobiology of Aging

journal homepage: www.elsevier.com/locate/neuagingExome sequencing identifies 2 novel presenilin 1 mutations (p.L166V and p.S230R) in British early-onset Alzheimer's disease [☆]Celeste Sassi ^{a,b,*}, Rita Guerreiro ^{a,b}, Raphael Gibbs ^{a,b}, Jinhui Ding ^b, Michelle K. Lupton ^c, Claire Troakes ^c, Katie Lunnon ^c, Safa Al-Sarraj ^c, Kristelle S. Brown ^d, Christopher Medway ^d, Jenny Lord ^d, James Turton ^d, David Mann ^e, Julie Snowden ^f, David Neary ^f, Jeniffer Harris ^f, Jose Bras ^a, ARUK Consortium¹, Kevin Morgan ^d, John F. Powell ^c, Andrew Singleton ^b, John Hardy ^a^a University College London (UCL) Institute of Neurology, London, UK^b Laboratory of Neurogenetics, National Institute on Aging, National Institutes of Health, Bethesda, MD, USA^c Institute of Psychiatry, King's College London, London, UK^d School of Molecular Medical Sciences, Institute of Genetics, Queen's Medical Centre, University of Nottingham, Nottingham, UK^e Institute of Brain, Behaviour, and Mental Health, The University of Manchester, Manchester, UK^f Cerebral Function Unit Greater Manchester Neuroscience Centre, Manchester, UK

ARTICLE INFO

Article history:

Received 16 March 2014

Accepted 22 April 2014

Available online 2 May 2014

Keywords:

Early-onset Alzheimer's disease

APP

PSEN1

PSEN2

British cohort

ABSTRACT

Early-onset Alzheimer's disease (EOAD) represents 1%–2% of the Alzheimer's disease (AD) cases, and it is generally characterized by a positive family history and a rapidly progressive symptomatology. Rare coding and fully penetrant variants in amyloid precursor protein (*APP*), presenilin 1 (*PSEN1*), and presenilin 2 (*PSEN2*) are the only causative mutations reported for autosomal dominant AD. Thus, in this study we used exome sequencing data to rapidly screen rare coding variability in *APP*, *PSEN1*, and *PSEN2*, in a British cohort composed of 47 unrelated EOAD cases and 179 elderly controls, neuropathologically proven. We report 2 novel and likely pathogenic variants in *PSEN1* (p.L166V and p.S230R). A comprehensive catalog of rare pathogenic variants in the AD Mendelian genes is pivotal for a premortem diagnosis of autosomal dominant EOAD and for the differential diagnosis with other early onset dementias such as frontotemporal dementia (FTD) and Creutzfeldt-Jakob disease (CJD).

© 2014 The Authors. Published by Elsevier Inc. All rights reserved.

[☆] This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/3.0/>).

* Corresponding author at: Neurogenetics Laboratory, National Institutes of Health, 35 Convent Drive, Bethesda, MD 20892, USA and UCL Institute of Neurology Queen Square London WC1N 3BG, UK. Tel.: +44 (0)20 3456 7890; fax: +44 (0)20 7278 5069.

E-mail address: celeste.sassi.10@ucl.ac.uk (C. Sassi).

¹ The Alzheimer's Research UK (ARUK) Consortium: Peter Passmore, David Craig, Janet Johnston, Bernadette McGuinness, Stephen Todd, Queen's University Belfast, UK; Reinhard Heun, Royal Derby Hospital, UK; Heike Kölsch, University of Bonn, Germany; Patrick G. Kehoe, University of Bristol, UK; Nigel M. Hooper, University of Leeds, UK; Emma R.L.C. Vardy, University of Newcastle, UK; David M. Mann, Stuart Pickering-Brown, University of Manchester, UK; Kristelle Brown, James Lowe, Kevin Morgan, University of Nottingham, UK; A. David Smith, Gordon Wilcock, Donald Warden, University of Oxford (OPTIMA), UK; Clive Holmes, University of Southampton, UK.

1. Introduction

Early-onset Alzheimer's disease (EOAD), with onset of symptoms before 65 years of age, accounts for 1%–2% of the Alzheimer's disease (AD) cases. The disease commonly clusters within families and presents a rapid and severe progression. Generally, EOAD is a polygenic and complex disease. On the contrary, 10% of EOAD cases present a Mendelian autosomal dominant pattern of inheritance and are caused by rare and fully penetrant mutations in amyloid precursor protein (*APP*), presenilin 1 (*PSEN1*), and presenilin 2 (*PSEN2*), leading to A β ₄₂ overproduction (Goate et al., 1991; Pastor and Goate, 2004; Raux et al., 2005; Sherrington et al., 1995). *PSEN1* is the most common cause of EOAD (18%–50%), with 185 mutations reported (www.molgen.database/). By contrast, *APP* and *PSEN2* have been described in 5% and <1% of the EOAD cases, respectively (www.molgen.database/). Given the pivotal role of these genes in AD pathogenesis, a detailed catalog of pathogenic variants and an algorithm for their classification are fundamental premises for an effective genetic screening of the patients.

2. Methods

2.1. Cases and controls

Our cohort was composed of 47 independent EOAD (age at onset ≤ 65 years) autopsy proven cases and 179 elderly (>60 years), unrelated controls. Three patients reported a positive family history of AD. The other patients were referred as apparently sporadic EOAD cases.

All the patients and controls were Caucasian, mostly from the UK (London, Manchester, Nottingham, and Edinburgh) and to a lesser extent from North America. The average age at diagnosis was 57.7 years (range 41–64 years) for the EOAD patients and the mean age of ascertainment was 78 years (range 60–102 years) for the controls (Table 1).

Written informed consent was obtained for each individual and the study was approved by the appropriate institutional review boards.

2.2. Exome sequencing

Library preparation for next-generation sequencing was performed according to the NimbleGen (Roche NimbleGen v2) and TruSeq (Illumina) sample-preparation protocols. DNA libraries were then hybridized to exome-capture probes with NimbleGen SeqCap EZ Human Exome Library, version 2.0 (Roche NimbleGen) or TruSeq (Illumina). Each capture method covers the *APP*, *PSEN1*, and *PSEN2* loci. Exome-enriched libraries were sequenced on the Illumina HiSeq 2000 using 2×100 bp paired end read cycles.

2.3. Bioinformatics

Sequence alignment and variant calling was performed against the reference human genome (UCSC hg19). Paired end sequence reads (2×100 bp paired end read cycles) were aligned using the Burrows-Wheeler aligner (Li and Durbin, 2009). Format conversion and indexing were performed with Picard (www.picard.sourceforge.net/index.shtml). The Genome Analysis Toolkit was used to recalibrate base quality scores, perform local re-alignments around indels, and to call and filter the variants (McKenna et al., 2010). VCFtools was used to annotate gene information for the remaining novel variants. In total, 450,987 single-nucleotide variants and 41,678 indels were called. We used snpEff v3.2 software to annotate the variants. Variants were checked against established databases (1000 Genomes Project and dbSNP v.134). The protein coding effects of variants was predicted using SIFT, Polyphen2, and SeattleSeq Annotation (gvs.gs.washington.edu/SeattleSeqAnnotation). All variants within the coding regions of *APP*, *PSEN1*, and *PSEN2* were annotated for both cases and controls.

2.4. Sanger sequencing

All rare variants identified by whole-exome sequencing in the candidate genes were validated by Sanger sequencing.

Primers for exons harboring rare variants were designed in Primer3 (<http://bioinfo.ut.ee/primer3-0.4.0/>) using UCSC (<http://genome.ucsc.edu/>)

reference sequences NM_000484 (*APP*), NM_000021.3 (*PSEN1*), and NM_000447.2 (*PSEN2*).

Purified sequences were analyzed on an ABI 3730 DNA Analyzer (Applied Biosystems, CA, USA) and electropherograms were visualized in Sequencher software (version 4.2 Gene Codes Corporation, MI, USA).

2.5. APOE genotyping

APOE genotypes comprising the *APOE* $\epsilon 2$, $\epsilon 3$, and $\epsilon 4$ alleles were assayed using the TaqMan method (Applied Biosystems Inc [ABI], Foster City, CA, USA). Single nucleotide polymorphism-specific primers and probes were designed by ABI (TaqMan genotyping assays).

3. Results

We report the experience of a mutation screening in *APP*, *PSEN1*, and *PSEN2* in 47 EOAD unrelated cases and 179 elderly controls neuropathologically proven, from the UK. 9 EOAD cases (19%) and 5 controls (2.8%) carried at least 1 rare coding variant in the genes studied.

In our EOAD cohort, we identified a total of 8 rare coding variants in *APP*, *PSEN1*, and *PSEN2*, absent in controls. Of these, 3 have been previously reported in familial AD (p.C410Y and p.113+1delG in *PSEN1* and p.V717L in *APP*) (Campion et al., 1995; De Jonghe et al., 2001; Murrell et al., 2000; Tysoe et al., 1998) and 4 are novel variants: 2 detected in *PSEN1* (p.L166V and p.S230R) and 2 in *APP* (p.K496Q and p.620L). In addition, we report a rare and likely tolerated polymorphism in *PSEN2* (p.S130L, rs63750197), present in 1 case and 1 control (Table 2).

PSEN1 p.L166V maps to the third transmembrane domain (TM3) of PS1, on the α -helix surface and corresponds to a conserved domain in *PSEN2* (p.L172), where no variants or mutations have been reported. Moreover, it has been described as possibly damaging by in silico predictions (SIFT and Polyphen2). This variant was detected in a very early onset case, presenting typical and rapidly progressing AD features with extrapyramidal signs (parkinsonism and myoclonus). At the age of 42 years, this patient developed difficulties in memory, necessitating his retirement from work. Over the next 2 years, he presented a decline in cognitive function, culminating in his medical referral. His memory loss had become more severe, he had difficulties in verbal expression, reading, writing, and calculation, and he was spatially disoriented. He had become generally apathetic. Neurologic examination revealed extrapyramidal signs and myoclonus. Neuropsychological examination revealed profound impairments in memory, language, perceptuospatial skills, and praxis. Social skills were well preserved, and he was frustrated by his difficulties.

He was reviewed at regular intervals over the following 4 years. During that time there was dramatic decline in his physical and cognitive skills. He showed marked parkinsonism and myoclonus. Speech was unintelligible, and he showed profound perceptual and spatial impairments. A single photon emission computerized tomography (SPECT) scan, 4 years after symptom onset, showed temporoparietal hypoperfusion. The profile remained typical of Alzheimer's disease. He died aged 50 years, 8 years after onset of

Table 1
Cohort

Cohort	n	Diagnosis	Sequencing strategy	Age (y) mean \pm SD (range)	Male (%)	<i>APOE</i> $\epsilon 4+$ (%)
EOAD	47	Clinical and neuropathological	Exome sequencing	57.7 (41–64)	55	57
Controls	179	Clinical and neuropathological	Exome sequencing	78 (60–102)	55	40

Key: EOAD, early-onset Alzheimer's disease; SD, standard deviation.

Table 2
Rare coding variants found in *APP*, *PSEN1*, and *PSEN2* in 47 EOAD cases and 179 elderly controls. *APP* (NM_000484); presenilins 1 and 2, *PSEN1* (NM_000021.3), and *PSEN2* (NM_000447.2)

Variant interpretation	Gene	Variant	Minor allele	Status	SIFT and/or polyphen	EOAD (n = 47)		Controls (n = 179)		Comment	
						Count (%)	(AAO-AAAD) (y)	Genotype	APOE		Count (%)
Known causative mutations											
<i>APP</i> p.V717L	<i>APP</i>	p.V717L	A	rs63750264	Possibly damaging	1 (2)	59	C/A	ε3ε3	Causal mutation in FAD	
<i>PSEN1</i> p.C410Y	<i>PSEN1</i>	p.C410Y	A	rs661	Possibly damaging	1 (2)	49–57	G/A	ε3ε3	Causal mutation in FAD	
<i>PSEN1</i> p.113 + 1delG, splice5	<i>PSEN1</i>	p.113 + 1delG, splice5	delG	rs63751475	Possibly damaging	2 (4.2)	41–51	G/-	ε4ε4, ε3ε3	Causal mutation in FAD	
Probable pathogenic variants ^a											
<i>PSEN1</i> p.L166V	<i>PSEN1</i>	p.L166V	G	Novel	Possibly damaging	1 (2)	42–50	C/G	ε3ε3	L166del, L166P, L166R causal mutations in FAD; L166H causal mutation in a pseudo-sporadic AD case	
Likely rare benign polymorphisms											
<i>PSEN1</i> p.S230R	<i>PSEN1</i>	p.S230R	G	Novel	Possibly damaging	1 (2)	56–66	T/G	ε3ε3	S230I reported in a French EOAD family	
<i>APP</i> p.K496Q	<i>APP</i>	p.K496Q	G	Novel	Possibly damaging	1 (2)	NA	T/G	ε3ε4	Exon 11	
<i>APP</i> p.620L	<i>APP</i>	p.620L	A	Novel	Tolerated	1 (2)	65	G/A	ε3ε4	Exon 14	
<i>PSEN2</i> p.S130L	<i>PSEN2</i>	p.S130L	T	rs63750197	Possibly damaging	1 (2)	61–65	C/T	ε3ε3		
									1 (0.5)	C/T	ε2ε2

Key: AAD, age at death; AAO, age at onset; AD, Alzheimer's disease; APP, amyloid precursor protein; FAD, familial Alzheimer's disease; PSEN1, presenilin 1; PSEN2, presenilin 2.
^a Classification based on the algorithm proposed by [Guerreiro et al. \(2010\)](#).

symptoms. The neuropathological examination revealed advanced AD lesions (Braak VI, CERAD C). There was no relevant previous medical history and no clear history of dementia in any first-degree relative, still alive, and in good health conditions.

Finally, the *PSEN1* p.L166V was not found in 179 healthy elderly neuropathologically confirmed controls, or in any control present in genomic variant databases (dbSNP v.137, ESP).

PSEN1 p.S230R is a nonconservative amino acid change in the fifth transmembrane domain (TM5), on the α -helix surface of PS1. It corresponds to a conserved residue in *PSEN2* (p.S236), where no variants or mutations have been described and was predicted as possibly damaging by SIFT and Polyphen2. The carrier presented features consistent with the reported clinical phenotype in most *PSEN1* mutations. The patient was diagnosed at 58 years, with a 2-year history of problems in memory. Neurologic examination was entirely normal. His behavior was socially appropriate and he was insightful into his difficulties. Neuropsychological examination confirmed the problems in memory and language. Performance in other cognitive domains was well preserved. Functional imaging using SPECT showed left posterior hypoperfusion. The patient's father had died in his early 60s of a similar dementing illness. The patient was clinically diagnosed with Alzheimer's disease, which was suspected to be familial.

He was reviewed at regular intervals. His cognitive skills deteriorated gradually. His memory became worse; he had increased difficulties in verbal expression and visuospatial tasks. He presented parkinsonian signs and myoclonus. A repeat SPECT scan revealed bilateral parietal hypoperfusion, more marked on the left side. Latterly, he became severely parkinsonian and mute. He died aged 66 years, 10 years after onset of symptoms. The neuropathological features revealed a fully developed Alzheimer's disease (Braak VI, CERAD C).

Finally, although we cannot rule out a pathogenic role for *APP* p.K496Q and p.620L, we suggest they are likely to be rare benign polymorphisms, as they cluster outside exon 16 and 17, where all pathogenic mutations have been reported up to date (www.molgen.database/).

4. Discussion

Mutations in *APP*, *PSEN1*, and *PSEN2* are the only fully penetrant variants described for EOAD. Thus, in this study we used exome-sequencing data to investigate rare coding variability in *APP*, *PSEN1*, and *PSEN2*, in a British cohort composed of 47 unrelated EOAD cases and 179 elderly controls. In this cohort, 6 patients of 47 (12.7%) and none of the controls carried a rare coding variant in *PSEN1*. This confirms that *PSEN1* is the major gene involved in EOAD and suggests that rare coding variability in this gene is commonly detrimental. Among the variants detected in *PSEN1*, 2 are novel (p.L166V and p.S230R), likely harmful and classified as probable pathogenic, following the algorithm proposed by [Guerreiro et al. \(2010\)](#). These variants have been detected in 2 patients who presented the typical hallmarks of *PSEN1* mutations: early onset of symptoms, aggressive symptomatology, and extrapyramidal signs (parkinsonism and myoclonus).

PSEN1 p.L166 codon seems to be a very vulnerable site for mutations: 4 different causal mutations (p.L166del, p.L166H, p.L166P, and p.L166R) have been previously described in this same residue in 3 familial AD cases (p.L166R, p.L166P, and p.L166del) ([Ezquerro et al., 2000](#); [Knight et al., 2007](#); [Moehlmann et al., 2002](#)) and in 1 pseudo-sporadic case (p.L166H) ([Pantieri et al., 2005](#)). From these, p.L166P is known to be a very aggressive mutation: it causes onset of symptoms before the third decade of age, significantly increases A β ₄₂ production, and impairs NOTCH1 signaling in a cell culture system ([Moehlmann et al., 2002](#)). Furthermore, the amino acid

position 166, located in the TM3 of PS1 shows a significant phylogenetic conservation across different species and in homologous proteins such as PS2, suggesting that the position is of functional significance.

In our cohort, the *PSEN1* p.L166V variant has been detected in a patient with very young onset of age (42 years at diagnosis), thus supporting the severe effects of an amino acid substitution in this conserved residue. Surprisingly, the patient's parents were not affected by Alzheimer's disease. This suggests that the p.L166V may be a de novo mutation or that other genetic factors may influence the phenotype of *PSEN1* mutations.

PSEN1 p.S230R clusters in the TM5 domain of PS1 and is the homologous of a conserved residue in *PSEN2* (p.S236). The patient carrying this variant presented a family history of AD. Recently, a different missense mutation in this codon (p.S230I) has been reported as possible pathogenic in a French family with EOAD (Wallon et al., 2012). Thus, the p.S230R is likely to be a novel mutation for early onset, autosomal dominant Alzheimer's disease.

In summary, we confirm that *PSEN1* is commonly associated with EOAD and we report 2 novel and likely pathogenic mutations detected in this gene (p.L166V and p.S230R). A comprehensive catalog of rare pathogenic variants in Alzheimer's disease Mendelian genes is pivotal: (1) for a premortem diagnosis of EOAD patients; (2) for the differential diagnosis of other types of early onset dementias (FTD and CJD); and (3) for the identification of at risk relatives who may be potential candidates for clinical trials.

Disclosure statement

The authors have nothing to disclose.

Acknowledgements

This study was supported by the Alzheimer's Research UK, the Medical Research Council, the Wellcome Trust, the Wellcome Trust/MRC Joint Call in Neurodegeneration Award (WT089698) to the UK Parkinson's Disease Consortium (whose members are from the University College London Institute of Neurology, the University of Sheffield, and the MRC Protein Phosphorylation Unit at the University of Dundee), grants (P50 AG016574, U01 AG006786, and R01 AG18023), the National Institute for Health Research Biomedical Research Unit in Dementia at University College London Hospitals, University College London; the National Institutes of Health Research, the Wellcome Trust Medical Research Council, an anonymous charitable foundation, the Fondation pour la Recherche sur Alzheimer, the Big Lottery (to Dr Morgan); a fellowship from Alzheimer's Research UK (to Dr Guerreiro); and the Intramural Research Programs of the National Institute on Aging and the National Institute of Neurological Disease and Stroke, National Institutes of Health (Department of Health and Human Services Project number, ZO1 AG000950-10). The MRC London Neurodegenerative Diseases Brain Bank and the Manchester Brain Bank from Brains for Dementia Research are jointly funded from ARUK and AS via ABBUK Ltd.

References

- Campion, D., Flaman, J.M., Brice, A., Hannequin, D., Dubois, B., Martin, C., Moreau, V., Charbonnier, F., Didierjean, O., Tardieu, S., 1995. Mutations of the presenilin 1 gene in families with early-onset Alzheimer's disease. *Hum. Mol. Genet.* 4, 2373–2377.
- De Jonghe, C., Esselens, C., Kumar-Singh, S., Craessaerts, K., Serneels, S., Checler, F., Annaert, W., Van Broeckhoven, C., De Strooper, B., 2001. Pathogenic APP

- mutations near the gamma-secretase cleavage site differentially affect Abeta secretion and APP C-terminal fragment stability. *Hum. Mol. Genet.* 10, 1665–1671.
- Ezquerro, M., Carnero, C., Blesa, R., Oliva, R., 2000. A novel presenilin 1 mutation (Leu166Arg) associated with early-onset Alzheimer disease. *Arch. Neurol.* 57, 485–488.
- Goate, A., Chartier-Harlin, M.C., Mullan, M., Brown, J., Crawford, F., Fidani, L., Giuffra, L., Haynes, A., Irving, N., James, L., 1991. Segregation of a missense mutation in the amyloid precursor protein gene with familial Alzheimer's disease. *Nature* 349, 704–706. <http://dx.doi.org/10.1038/349704a0>.
- Guerreiro, R.J., Baquero, M., Blesa, R., Boada, M., Brás, J.M., Bullido, M.J., Calado, A., Crook, R., Ferreira, C., Frank, A., Gómez-Isla, T., Hernández, I., Lleó, A., Machado, A., Martínez-Lage, P., Masdeu, J., Molina-Porcel, L., Molinuevo, J.L., Pastor, P., Pérez-Tur, J., Relvas, R., Oliveira, C.R., Ribeiro, M.H., Rogaeva, E., Sa, A., Samaranch, L., Sánchez-Valle, R., Santana, I., Tarraga, L., Valdivieso, F., Singleton, A., Hardy, J., Clarimón, J., 2010. Genetic screening of Alzheimer's disease genes in Iberian and African samples yields novel mutations in presenilins and APP. *Neurobiol. Aging* 31, 725–731. <http://dx.doi.org/10.1016/j.neurobiolaging.2008.06.012>.
- Knight, W.D., Kennedy, J., Mead, S., Rossor, M.N., Beck, J., Collinge, J., Mummery, C., 2007. A novel presenilin 1 deletion (p.L166del) associated with early onset familial Alzheimer's disease. *Eur. J. Neurol.* 14, 829–831. <http://dx.doi.org/10.1111/j.1468-1331.2007.01857.x>.
- Li, H., Durbin, R., 2009. Fast and accurate short read alignment with Burrows-Wheeler transform. *Bioinformatics* 25, 1754–1760. <http://dx.doi.org/10.1093/bioinformatics/btp324>.
- McKenna, A., Hanna, M., Banks, E., Sivachenko, A., Cibulskis, K., Kernytzky, A., Garimella, K., Altshuler, D., Gabriel, S., Daly, M., DePristo, M.A., 2010. The Genome Analysis Toolkit: a MapReduce framework for analyzing next-generation DNA sequencing data. *Genome Res.* 20, 1297–1303. <http://dx.doi.org/10.1101/gr.107524.110>.
- Moehlmann, T., Winkler, E., Xia, X., Edbauer, D., Murrell, J., Capell, A., Kaether, C., Zheng, H., Ghetti, B., Haass, C., Steiner, H., 2002. Presenilin-1 mutations of leucine 166 equally affect the generation of the Notch and APP intracellular domains independent of their effect on Abeta 42 production. *Proc. Natl. Acad. Sci. U.S.A.* 99, 8025–8030. <http://dx.doi.org/10.1073/pnas.112686799>.
- Murrell, J.R., Hake, A.M., Quaid, K.A., Farlow, M.R., Ghetti, B., 2000. Early-onset Alzheimer disease caused by a new mutation (V717L) in the amyloid precursor protein gene. *Arch. Neurol.* 57, 885–887.
- Pantieri, R., Pardini, M., Cecconi, M., Dagna-Bricarelli, F., Vitali, A., Piccini, A., Russo, R., Borghi, R., Tabaton, M., 2005. A novel presenilin 1 L166H mutation in a pseudo-sporadic case of early-onset Alzheimer's disease. *Neurol. Sci.* 26, 349–350. <http://dx.doi.org/10.1007/s10072-005-0499-1>.
- Pastor, P., Goate, A.M., 2004. Molecular genetics of Alzheimer's disease. *Curr. Psychiatry Rep.* 6, 125–133.
- Raux, G., Guyant-Maréchal, L., Martin, C., Bou, J., Penet, C., Brice, A., Hannequin, D., Frebourg, T., Campion, D., 2005. Molecular diagnosis of autosomal dominant early onset Alzheimer's disease: an update. *J. Med. Genet.* 42, 793–795. <http://dx.doi.org/10.1136/jmg.2005.033456>.
- Sherrington, R., Rogaev, E.I., Liang, Y., Rogaeva, E.A., Levesque, G., Ikeda, M., Chi, H., Lin, C., Li, G., Holman, K., Tsuda, T., Mar, L., Foncin, J.F., Bruni, A.C., Montesi, M.P., Sorbi, S., Rainero, I., Pinessi, L., Nee, L., Chumakov, I., Pollen, D., Brookes, A., Sanseau, P., Polinsky, R.J., Wasco, W., Da Silva, H.A., Haines, J.L., Pericak-Vance, M.A., Tanzi, R.E., Roses, A.D., Fraser, P.E., Rommens, J.M., St George-Hyslop, P.H., 1995. Cloning of a gene bearing missense mutations in early-onset familial Alzheimer's disease. *Nature* 375, 754–760. <http://dx.doi.org/10.1038/375754a0>.
- Tysoe, C., Whittaker, J., Xuereb, J., Cairns, N.J., Cruts, M., Van Broeckhoven, C., Wilcock, G., Rubinsztein, D.C., 1998. A presenilin-1 truncating mutation is present in two cases with autopsy-confirmed early-onset Alzheimer disease. *Am. J. Hum. Genet.* 62, 70–76. <http://dx.doi.org/10.1086/301672>.
- Wallon, D., Rousseau, S., Rovelet-Lecrux, A., Quillard-Muraine, M., Guyant-Maréchal, L., Martinaud, O., Pariente, J., Puel, M., Rollin-Sillaire, A., Pasquier, F., Le Ber, I., Sarazin, M., Croisile, B., Boutoleau-Brettonnière, C., Thomas-Antérion, C., Paquet, C., Moreaud, O., Gabelle, A., Sellal, F., Sauvée, M., Laquerrière, A., Duyckaerts, C., Delisle, M.-B., Streichenberger, N., Lannes, B., Frebourg, T., Hannequin, D., Campion, D., collaborators of GMAJ project, 2012. The French series of autosomal dominant early onset Alzheimer's disease cases: mutation spectrum and cerebrospinal fluid biomarkers. *J. Alzheimers Dis.* 30, 847–856. <http://dx.doi.org/10.3233/JAD-2012-120172>.

Web references

- 1000 Genomes. <http://www.1000genomes.org/>.
- dbSNP. <http://www.ncbi.nlm.nih.gov/projects/SNP/>.
- ESP. <http://evs.gs.washington.edu/EVS/>.
- Molgen. <http://www.molgen.ua.ac.be/ADMutations/>.
- Picard. <http://picard.sourceforge.net/index.shtml>.
- Polyphen 2. <http://genetics.bwh.harvard.edu/pph2/>.
- SeattleSeq. <http://snp.gs.washington.edu/SeattleSeqAnnotation138/>.
- SIFT. <http://sift.jcvi.org/>.