

Hindawi Publishing Corporation  
Discrete Dynamics in Nature and Society  
Volume 2012, Article ID 137471, 14 pages  
doi:10.1155/2012/137471

## Research Article

# Oscillation and Nonoscillation Criteria for Nonlinear Dynamic Systems on Time Scales

**Shanliang Zhu and Chunyun Sheng**

*College of Mathematics and Physics, Qingdao University of Science and Technology,  
Qingdao 266061, China*

Correspondence should be addressed to Shanliang Zhu, zhushanliang77@163.com

Received 10 April 2012; Accepted 23 June 2012

Academic Editor: Jinde Cao

Copyright © 2012 S. Zhu and C. Sheng. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

We consider the nonlinear dynamic system  $x^\Delta(t) = a(t)g(y(t))$ ,  $y^\Delta(t) = -f(t, x^\sigma(t))$ . We establish some necessary and sufficient conditions for the existence of oscillatory and nonoscillatory solutions with special asymptotic properties for the system. We generalize the known results in the literature. Some examples are included to illustrate the results.

## 1. Introduction

In this paper we investigate the nonlinear two-dimensional dynamic system:

$$x^\Delta(t) = a(t)g(y(t)), \quad y^\Delta(t) = -f(t, x^\sigma(t)), \quad t \in [t_0, \infty)_{\mathbb{T}}, \quad (1.1)$$

where  $a(t)$  is a nonnegative, rd-continuous function which is defined for  $t \in [t_0, \infty)_{\mathbb{T}} = [t_0, \infty) \cap \mathbb{T}$ . Here,  $\mathbb{T}$  is a time scale unbounded from above. We assume throughout that  $g: \mathbb{R} \rightarrow \mathbb{R}$  is a continuous function with  $ug(u) > 0$  for  $u \neq 0$ , and  $f(t, u): [t_0, \infty)_{\mathbb{T}} \times \mathbb{R} \rightarrow \mathbb{R}$  is continuous as a function of  $u \in \mathbb{R}$  with sign property  $uf(t, u) > 0$  for  $u \neq 0$  and  $t \in [t_0, \infty)_{\mathbb{T}}$ .

By the solution of system (1.1), we mean a pair of nontrivial real-valued functions  $(x(t), y(t))$  which has property  $x, y \in C_{\text{rd}}^1([t_0, \infty)_{\mathbb{T}}, \mathbb{R})$  and satisfies system (1.1) for  $t \in [t_0, \infty)_{\mathbb{T}}$ . Our attention is restricted to those solutions  $(x(t), y(t))$  of system (1.1) which exist on some half-line  $[t_x, \infty)_{\mathbb{T}}$  and satisfy  $\sup\{|x(t)| + |y(t)| : t \geq t_x\} > 0$  for any  $t_x \geq t_0$ . As usual, a continuous real-valued function defined on  $[T_0, \infty)$  is said to be oscillatory if it has arbitrarily large zeros, otherwise it is said to be nonoscillatory. A solution  $(x(t), y(t))$  of system (1.1) is called oscillatory if both  $x(t)$  and  $y(t)$  are oscillatory functions, and otherwise it will be called nonoscillatory. System (1.1) is called oscillatory if its solutions are oscillatory.

The theory of time scales, which has recently received a lot of attention, was introduced by Hilger in his Ph.D. thesis in 1990 in order to unify continuous and discrete analysis (see [1]). Not only can this theory of the so-called “dynamic equations” unify the theories of differential equations and difference equations, but also extend these classical cases to cases “in between”, for example, to the so-called  $q$ -difference equations and can be applied on other different types of time scales. Since Hilger formed the definition of derivatives and integral on time scales, several authors have expounded on various aspects of this new theory; see the survey paper by Agarwal et al. [2] and references cited therein. A book on the subject of time scales (see [3]) summarizes and organizes much of time scale calculus. The reader is referred to Chapter 1 in [3] for the necessary time scale definitions and notations used throughout this paper.

The system (1.1) includes two-dimensional linear/nonlinear differential and difference systems, which were investigated in the literature, see for example [4–9] and the references therein.

On the other hand, the system (1.1) reduces to some important second-order dynamic equations in the particular case, for example

$$\left[ \frac{x^\Delta(t)}{a(t)} \right]^\Delta + b(t)f(x^\sigma(t)) = 0, \quad (1.2)$$

$$x^{\Delta\Delta}(t) + b(t)|x^\sigma(t)|^{\lambda-1}x^\sigma(t) = 0, \quad \lambda > 0,$$

where  $b(t)$  is rd-continuous on  $[t_0, \infty)_{\mathbb{T}}$ . In recent years there has been much research activity concerning the oscillation and nonoscillation of solutions of dynamic equation (1.2) on time scales. We refer the reader to the recent papers [10–13] and the references therein. However, most of previous studies for the system (1.1) have been restricted to the case where  $f(t, u) = b(t)f(u)$ , for example [4–8, 14–18] and the references therein. Erbe and Mert [14, 17] obtained some oscillation results for the system (1.1). Fu and Lin [15] obtained some oscillation and nonoscillation criteria for the linear dynamic system (1.1).

Since there are few works about oscillation and nonoscillation of dynamic systems on time scales (see [15]), motivated by [9, 14, 15], in this paper we investigate oscillatory and nonoscillatory properties for the system (1.1) in the case of general  $f(t, u)$  in which  $t$  and  $u$  are not necessarily separable. In the next section, by means of appropriate hypotheses on  $f(t, u)$  and fixed point theorem, we establish some new sufficient and necessary conditions for the existence of nonoscillatory solutions with special asymptotic properties for the system (1.1). In Section 3, we obtain sufficient and necessary conditions for all solutions of the system (1.1) to be oscillatory via the results in Section 2 and some inequality techniques without using Riccati transformation. Our results not only unify the known results of differential and difference systems but also extend and improve the existing results of dynamic systems on time scales in the literature.

## 2. Nonoscillation Results

In this section, we generalize and improve some results of [7–9, 15, 18]. Some necessary and sufficient conditions are given for the system (1.1) to admit the existence of nonoscillatory solutions with special asymptotic properties. These results will be used for the next section. Additional hypotheses on  $g(u)$  and  $f(t, u)$  are needed for this purpose.

(H<sub>1</sub>) For any positive constant  $l$  and  $L$  with  $l < L$ , there exist positive constants  $h$  and  $H$ , depending possibly on  $l$  and  $L$ , such that  $l \leq |u| \leq L$  implies

$$hf(t, l) \leq |f(t, u)| \leq Hf(t, L), \quad t \in [t_0, \infty)_{\mathbb{T}}. \quad (2.1)$$

(H<sub>2</sub>) There exists a positive constant  $k$  such that  $g(uv) \geq kg(u)g(v)$  for  $uv > 0$ .

(H<sub>3</sub>) For any positive constant  $l$  and  $L$  with  $l < L$ , there exist positive constants  $h$  and  $H$ , depending possibly on  $l$  and  $L$ , such that  $l \leq |u| \leq L$  implies

$$hf(t, l\theta(t)) \leq |f(t, u\theta(t))| \leq Hf(t, L\theta(t)), \quad t \in [t_0, \infty)_{\mathbb{T}}, \quad (2.2)$$

where  $\theta(t)$  is a positive nondecreasing function.

For convenience, we will employ the following notation:

$$A(s, t) = \int_s^t a(\tau) \Delta\tau, \quad s, t \in [t_0, \infty)_{\mathbb{T}}. \quad (2.3)$$

**Theorem 2.1.** *Assume that  $g$  is nondecreasing and that (H<sub>1</sub>) holds. Then system (1.1) has a nonoscillatory solution  $(x(t), y(t))$  such that  $\lim_{t \rightarrow \infty} x(t) = \alpha \neq 0$  and  $\lim_{t \rightarrow \infty} y(t) = 0$  if and only if for any positive constant  $d$*

$$\int_{t_0}^{\infty} a(t) g\left(d \int_t^{\infty} |f(s, c)| \Delta s\right) \Delta t < \infty \quad \text{for some } c \neq 0. \quad (2.4)$$

*Proof.* Suppose that  $(x(t), y(t))$  is a nonoscillatory solution of (1.1) such that  $\lim_{t \rightarrow \infty} x(t) = \alpha \neq 0$  and  $\lim_{t \rightarrow \infty} y(t) = 0$ . Without loss of generality, we assume that  $\alpha > 0$ . Then there exist  $t_1 \in [t_0, \infty)_{\mathbb{T}}$  and positive constants  $l$  and  $L$  such that  $l \leq x(t) \leq L$  for  $t \in [t_1, \infty)_{\mathbb{T}}$ . Condition (H<sub>1</sub>) implies that

$$f(t, x^\sigma(t)) \geq hf(t, l) \quad (2.5)$$

for  $t \in [t_1, \infty)_{\mathbb{T}}$  and some constant  $h > 0$ . It follows from the second equation in (1.1) that

$$y(s) - y(t) = - \int_t^s f(\tau, x^\sigma(\tau)) \Delta\tau. \quad (2.6)$$

Let  $s \rightarrow \infty$  and noting that  $\lim_{s \rightarrow \infty} y(s) = 0$ , we have

$$y(t) = \int_t^{\infty} f(\tau, x^\sigma(\tau)) \Delta\tau, \quad t \in [t_1, \infty)_{\mathbb{T}}. \quad (2.7)$$

Thus, from (2.5), (2.7) and the first equation in (1.1), we obtain that

$$\begin{aligned}
\infty &> \lim_{t \rightarrow \infty} x(t) - x(t_1) = \int_{t_1}^{\infty} a(s)g(y(s))\Delta s \\
&= \int_{t_1}^{\infty} a(s)g\left(\int_s^{\infty} f(\tau, x^\sigma(\tau))\Delta\tau\right)\Delta s \\
&\geq \int_{t_1}^{\infty} a(s)g\left(h \int_s^{\infty} f(\tau, l)\Delta\tau\right)\Delta s,
\end{aligned} \tag{2.8}$$

which implies that (2.4) holds.

Conversely, suppose that (2.4) holds, we may assume that  $c > 0$ . In view of  $(H_1)$ , there is a constant  $H > 0$  such that  $c/2 \leq x(t) \leq c$  implies

$$f(t, x^\sigma(t)) \leq Hf(t, c), \quad t \in [t_0, \infty)_{\mathbb{T}}. \tag{2.9}$$

Since (2.4) holds, we can choose  $t_1 \in [t_0, \infty)_{\mathbb{T}}$  large enough such that

$$\int_{t_1}^{\infty} a(s)g\left(H \int_s^{\infty} f(\tau, c)\Delta\tau\right)\Delta s \leq \frac{c}{2}. \tag{2.10}$$

Let  $BC[t_0, \infty)_{\mathbb{T}}$  be the Banach space of all real-valued rd-continuous functions on  $[t_0, \infty)_{\mathbb{T}}$  endowed with the norm  $\|x\| = \sup_{t \in [t_0, \infty)_{\mathbb{T}}} |x(t)| < \infty$ . We defined a bounded, convex, and closed subset  $\Omega$  of  $BC[t_0, \infty)_{\mathbb{T}}$  as

$$\Omega = \left\{ x \in BC[t_0, \infty)_{\mathbb{T}} : \frac{c}{2} \leq x(t) \leq c \right\}. \tag{2.11}$$

Define an operator  $\Gamma : \Omega \rightarrow BC[t_0, \infty)_{\mathbb{T}}$  as follows:

$$(\Gamma x)(t) = \begin{cases} c - \int_t^{\infty} a(s)g\left(\int_s^{\infty} f(\tau, x^\sigma(\tau))\Delta\tau\right)\Delta s, & t \in [t_1, \infty)_{\mathbb{T}}, \\ c - \int_{t_1}^{\infty} a(s)g\left(\int_s^{\infty} f(\tau, x^\sigma(\tau))\Delta\tau\right)\Delta s, & t \in [t_0, t_1]_{\mathbb{T}}. \end{cases} \tag{2.12}$$

Now we show that  $\Gamma$  satisfies the assumptions of Schauder's fixed-point theorem (see [19, Corollary 6]).

(i) We will show that  $\Gamma x \in \Omega$  for any  $x \in \Omega$ . In fact, for any  $x \in \Omega$  and  $t \in [t_1, \infty)_{\mathbb{T}}$ , in view of (2.10), we get

$$\begin{aligned} c \geq (\Gamma x)(t) &= c - \int_t^\infty a(s)g\left(\int_s^\infty f(\tau, x^\sigma(\tau))\Delta\tau\right)\Delta s \\ &\geq c - \int_{t_1}^\infty a(s)g\left(H \int_s^\infty f(\tau, c)\Delta\tau\right)\Delta s \\ &\geq c - \frac{c}{2} = \frac{c}{2}. \end{aligned} \tag{2.13}$$

Similarly, we can prove that  $c/2 \leq (\Gamma x)(t) \leq c$  for any  $x \in \Omega$  and  $t \in [t_0, t_1)_{\mathbb{T}}$ . Hence,  $\Gamma x \in \Omega$  for any  $x \in \Omega$ .

(ii) We prove that  $\Gamma$  is a completely continuous mapping. First, we consider the continuity of  $\Gamma$ . Let  $x_n \in \Omega$  and  $\|x_n - x\| \rightarrow 0$  as  $n \rightarrow \infty$ , then  $x \in \Omega$  and  $|x_n(t) - x(t)| \rightarrow 0$  as  $n \rightarrow \infty$  for any  $t \in [t_0, \infty)_{\mathbb{T}}$ . Consequently, by the continuity of  $g$  and  $f$ , for any  $t \in [t_1, \infty)_{\mathbb{T}}$ , we have

$$\lim_{n \rightarrow \infty} \left| a(t) \left[ g\left(\int_t^\infty f(\tau, x_n^\sigma(\tau))\Delta\tau\right) - g\left(\int_t^\infty f(\tau, x^\sigma(\tau))\Delta\tau\right) \right] \right| = 0. \tag{2.14}$$

From (2.9), we obtain that

$$a(t) \left| g\left(\int_t^\infty f(\tau, x_n^\sigma(\tau))\Delta\tau\right) - g\left(\int_t^\infty f(\tau, x^\sigma(\tau))\Delta\tau\right) \right| \leq 2a(t)g\left(H \int_t^\infty f(\tau, c)\Delta\tau\right). \tag{2.15}$$

On the other hand, from (2.12) we have

$$|(\Gamma x_n)(t) - (\Gamma x)(t)| \leq \int_{t_1}^\infty a(s) \left| g\left(\int_s^\infty f(\tau, x_n^\sigma(\tau))\Delta\tau\right) - g\left(\int_s^\infty f(\tau, x^\sigma(\tau))\Delta\tau\right) \right| \Delta s \tag{2.16}$$

for  $t \in [t_0, t_1)_{\mathbb{T}}$  and

$$|(\Gamma x_n)(t) - (\Gamma x)(t)| \leq \int_t^\infty a(s) \left| g\left(\int_s^\infty f(\tau, x_n^\sigma(\tau))\Delta\tau\right) - g\left(\int_s^\infty f(\tau, x^\sigma(\tau))\Delta\tau\right) \right| \Delta s \tag{2.17}$$

for  $t \in [t_1, \infty)_{\mathbb{T}}$ . Therefore, from (2.16) and (2.17), we have

$$\|\Gamma x_n - \Gamma x\| \leq \int_{t_1}^\infty a(s) \left| g\left(\int_s^\infty f(\tau, x_n^\sigma(\tau))\Delta\tau\right) - g\left(\int_s^\infty f(\tau, x^\sigma(\tau))\Delta\tau\right) \right| \Delta s. \tag{2.18}$$

Referring to Chapter 5 in [20], we see that the Lebesgue dominated convergence theorem holds for the integral on time scales. Then, from (2.14) and (2.15), (2.18) yields  $\lim_{n \rightarrow \infty} \|\Gamma x_n - \Gamma x\| = 0$ , which implies that  $\Gamma$  is continuous on  $\Omega$ .

Next, we show that  $\Gamma\Omega$  is uniformly cauchy. In fact, for any  $\varepsilon > 0$ , take  $t_2 \in [t_1, \infty)_{\mathbb{T}}$  and  $t_2 > t_1$  such that

$$\int_{t_2}^{\infty} a(s)g\left(H \int_s^{\infty} f(\tau, c)\Delta\tau\right)\Delta s \leq \varepsilon. \quad (2.19)$$

Then for any  $x \in \Omega$  and  $t, r \in [t_2, \infty)_{\mathbb{T}}$ , we have

$$\begin{aligned} |(\Gamma x)(t) - (\Gamma x)(r)| &\leq \left| \int_t^{\infty} a(s)g\left(H \int_s^{\infty} f(\tau, c)\Delta\tau\right)\Delta s \right| \\ &\quad + \left| \int_r^{\infty} a(s)g\left(H \int_s^{\infty} f(\tau, c)\Delta\tau\right)\Delta s \right| \\ &\leq 2 \int_{t_2}^{\infty} a(s)g\left(H \int_s^{\infty} f(\tau, c)\Delta\tau\right)\Delta s \leq 2\varepsilon. \end{aligned} \quad (2.20)$$

This means that  $\Gamma\Omega$  is uniformly cauchy.

Finally, we prove that  $\Gamma\Omega$  is equicontinuous on  $[t_0, t_2]_{\mathbb{T}}$  for any  $t_2 \in [t_0, \infty)_{\mathbb{T}}$ . Without loss of generality, we set  $t_2 > t_1$ . For any  $x \in \Omega$ , we have  $|(\Gamma x)(t) - (\Gamma x)(r)| \equiv 0$  for  $t, r \in [t_0, t_1]_{\mathbb{T}}$  and

$$\begin{aligned} |(\Gamma x)(t) - (\Gamma x)(r)| &= \left| \int_t^{\infty} a(s)g\left(\int_s^{\infty} f(\tau, x^\sigma(\tau))\Delta\tau\right)\Delta s - \int_r^{\infty} a(s)g\left(\int_s^{\infty} f(\tau, x^\sigma(\tau))\Delta\tau\right)\Delta s \right| \\ &\leq \left| \int_t^r a(s)g\left(H \int_s^{\infty} f(\tau, c)\Delta\tau\right)\Delta s \right| \end{aligned} \quad (2.21)$$

for  $t, r \in [t_1, t_2]_{\mathbb{T}}$ .

Now, we see that for any  $\varepsilon > 0$ , there exists  $\delta > 0$  such that when  $t, r \in [t_0, t_2]_{\mathbb{T}}$  with  $|t - r| < \delta$ ,  $|(\Gamma x)(t) - (\Gamma x)(r)| < \varepsilon$  for any  $x \in \Omega$ . This means that  $\Gamma\Omega$  is equicontinuous on  $[t_0, t_2]_{\mathbb{T}}$  for any  $t_2 \in [t_0, \infty)_{\mathbb{T}}$ . By Arzela-Ascoli theorem (see [19, Lemma 4]),  $\Gamma\Omega$  is relatively compact. From the above, we have proved that  $\Gamma$  is a completely continuous mapping.

By Schauder's fixed point theorem, there exists  $x \in \Omega$  such that  $\Gamma x = x$ . Therefore, we have

$$x(t) = c - \int_t^{\infty} a(s)g\left(\int_s^{\infty} f(\tau, x^\sigma(\tau))\Delta\tau\right)\Delta s, \quad t \in [t_1, \infty)_{\mathbb{T}}. \quad (2.22)$$

Set

$$y(t) = \int_t^{\infty} f(\tau, x^\sigma(\tau))\Delta\tau, \quad t \in [t_1, \infty)_{\mathbb{T}}. \quad (2.23)$$

Then  $\lim_{t \rightarrow \infty} y(t) = 0$  and  $y^\Delta(t) = -f(t, x^\sigma(t))$ . On the other hand,

$$x(t) = c - \int_t^\infty a(s)g(y(s))\Delta s, \quad (2.24)$$

which implies that  $\lim_{t \rightarrow \infty} x(t) = c$  and  $x^\Delta(t) = a(t)g(y(t))$ . The proof is complete.  $\square$

**Corollary 2.2.** *Suppose that  $g$  is nondecreasing and that  $(H_1)$  and  $(H_2)$  hold. Then system (1.1) has a nonoscillatory solution  $(x(t), y(t))$  such that  $\lim_{t \rightarrow \infty} x(t) = \alpha \neq 0$  and  $\lim_{t \rightarrow \infty} y(t) = 0$  if and only if for some  $c \neq 0$*

$$\int_{t_0}^\infty a(t)g\left(\int_t^\infty |f(s, c)|\Delta s\right)\Delta t < \infty. \quad (2.25)$$

**Theorem 2.3.** *Suppose that  $\lim_{t \rightarrow \infty} A(t_0, t) = \infty$  and  $g$  is nondecreasing. Suppose further that  $(H_3)$  holds. Then system (1.1) has a nonoscillatory solution  $(x(t), y(t))$  such that  $\lim_{t \rightarrow \infty} (x(t)/A(t_0, t)) = \alpha \neq 0$  and  $\lim_{t \rightarrow \infty} y(t) = \beta \neq 0$  if and only if for some  $c \neq 0$*

$$\int_{t_0}^\infty |f(t, cA(t_0, \sigma(t)))|\Delta t < \infty. \quad (2.26)$$

*Proof.* Suppose that  $(x(t), y(t))$  is a nonoscillatory solution of (1.1) such that  $\lim_{t \rightarrow \infty} (x(t)/A(t_0, t)) = \alpha \neq 0$  and  $\lim_{t \rightarrow \infty} y(t) = \beta \neq 0$ . We may assume that  $\alpha > 0$ . Hence, there exist  $t_1 \in [t_0, \infty)_{\mathbb{T}}$  and positive constant  $l, L$  such that  $lA(t_0, t) \leq x(t) \leq LA(t_0, t)$  and  $y^\Delta(t) < 0$ ,  $y(t) > \beta$  for  $t \in [t_1, \infty)_{\mathbb{T}}$ . By condition  $(H_3)$ , there exists a constant  $h > 0$  such that  $f(t, x^\sigma(t)) \geq hf(t, lA(t_0, \sigma(t)))$  for  $t \in [t_1, \infty)_{\mathbb{T}}$ . According to the second equation in (1.1), we have

$$\infty > y(t_1) - \beta = \int_{t_1}^\infty f(t, x^\sigma(t))\Delta t \geq h \int_{t_1}^\infty f(t, lA(t_0, \sigma(t)))\Delta t, \quad (2.27)$$

which implies that (2.26) holds with  $c = l$ .

Conversely, Let (2.26) holds for some  $c = 2p$ , where  $p > 0$ . By  $(H_3)$ , there exists a constant  $H > 0$  such that  $p \leq u \leq 2p$  implies  $f(t, uA(t_0, \sigma(t))) \leq Hf(t, 2pA(t_0, \sigma(t)))$  for  $t \in [t_0, \infty)_{\mathbb{T}}$ . Take  $t_1 \in [t_0, \infty)_{\mathbb{T}}$  so large that

$$H \int_{t_1}^\infty f(t, cA(t_0, \sigma(t)))\Delta t \leq d, \quad (2.28)$$

where  $d = g^{-1}(c)/2$ . We introduce  $BC[t_1, \infty)_{\mathbb{T}}$  be the partially ordered Banach space of all real-valued and rd-continuous functions  $x(t)$  with the norm  $\|x\| = \sup_{t \in [t_1, \infty)_{\mathbb{T}}} (|x(t)|/A(t_1, t))$ , and the usual pointwise ordering  $\leq$ .

Define

$$\Omega = \{x \in BC[t_1, \infty)_{\mathbb{T}} : g(d)A(t_1, t) \leq x(t) \leq g(2d)A(t_1, t)\}. \quad (2.29)$$

It is easy to see that  $\Omega$  is a bounded, convex, and closed subset of  $BC[t_1, \infty)_{\mathbb{T}}$ . Let us further define an operator  $\Gamma : \Omega \rightarrow BC[t_1, \infty)_{\mathbb{T}}$  as follows:

$$(\Gamma x)(t) = \int_{t_1}^t a(s)g\left(d + \int_s^\infty f(\tau, x^\sigma(\tau))\Delta\tau\right)\Delta s, \quad t \in [t_1, \infty)_{\mathbb{T}}. \quad (2.30)$$

Since it can be shown that  $\Gamma$  is continuous and sends  $\Omega$  into a relatively compact subset of  $\Omega$ , the Schauder's fixed point theorem ensures that the existence of an  $x \in \Omega$  such that  $x = \Gamma x$ , this is

$$x(t) = \int_{t_1}^t a(s)g\left(d + \int_s^\infty f(\tau, x^\sigma(\tau))\Delta\tau\right)\Delta s, \quad t \in [t_1, \infty)_{\mathbb{T}}. \quad (2.31)$$

Set

$$y(t) = d + \int_t^\infty f(\tau, x^\sigma(\tau))\Delta\tau, \quad t \in [t_1, \infty)_{\mathbb{T}}. \quad (2.32)$$

Then  $\lim_{t \rightarrow \infty} y(t) = d$  and  $y^\Delta(t) = -f(t, x^\sigma(t))$ . On the other hand, by L'Hôpital's Rule (see [15, Lemma 2.11]), we have

$$\begin{aligned} \lim_{t \rightarrow \infty} \frac{x(t)}{A(t_0, t)} &= \lim_{t \rightarrow \infty} \frac{a(t)g\left(d + \int_t^\infty f(\tau, x^\sigma(\tau))\Delta\tau\right)}{a(t)} \\ &= \lim_{t \rightarrow \infty} g\left(d + \int_t^\infty f(\tau, x^\sigma(\tau))\Delta\tau\right) = g(d) \neq 0, \end{aligned} \quad (2.33)$$

and  $x^\Delta(t) = a(t)g(y(t))$ . The proof is complete.  $\square$

*Remark 2.4.* Theorems 2.1 and 2.3 extend and improve essentially the known results of [7–9, 15, 18].

### 3. Oscillation Results

In this section, we need some additional conditions to guarantee that the system (1.1) has oscillatory solutions.

(H<sub>4</sub>) There exists a continuous nondecreasing function  $\varphi : \mathbb{R} \rightarrow \mathbb{R}$  such that

$$\operatorname{sgn} \varphi(u) = \operatorname{sgn} u, \quad \int^{\pm\infty} \frac{du}{g(\varphi(u))} < \infty, \quad (3.1)$$

and  $|f(t, u)| \geq |f(t, l)| |\varphi(u)|$ ,  $t \in [t_0, \infty)_{\mathbb{T}}$ ,  $|u| \geq u_0$  for some constants  $u_0 > 0$  and  $l \neq 0$  with  $\operatorname{sgn} l = \operatorname{sgn} u$ .



(H<sub>5</sub>) There exists a continuous nondecreasing function  $\varphi : [-M, M] \rightarrow \mathbb{R}$ ,  $M > 0$  being a constant, such that

$$\operatorname{sgn} \varphi(v) = \operatorname{sgn} v, \quad \int_0^{\pm M} \frac{dv}{\varphi(g(v))} < \infty, \quad (3.2)$$

and  $|f(t, uv)| \geq k|f(t, u)|\varphi(v)$ ,  $t \in [t_0, \infty)_{\mathbb{T}}$ ,  $u \neq 0$ ,  $0 < |v| < v_0$  for some positive constant  $k > 0$  and  $v_0 > 0$ .

**Theorem 3.1.** *Suppose that  $\lim_{t \rightarrow \infty} A(t_0, t) = \infty$  and  $g$  is nondecreasing. Suppose further that (H<sub>1</sub>), (H<sub>2</sub>) and (H<sub>4</sub>) hold. Then system (1.1) is oscillatory if and only if for all  $c \neq 0$*

$$\int_{t_0}^{\infty} a(t)g\left(\int_t^{\infty} |f(s, c)|\Delta s\right)\Delta t = \infty. \quad (3.3)$$

*Proof.* If (3.3) does not hold, by Theorem 2.1, system (1.1) has a nonoscillatory solution  $(x(t), y(t))$  such that  $\lim_{t \rightarrow \infty} x(t) = \alpha \neq 0$  and  $\lim_{t \rightarrow \infty} y(t) = 0$ .

Conversely, suppose that (3.3) holds and that (1.1) has a nonoscillatory solution  $(x(t), y(t))$  for  $t \in [t_0, \infty)_{\mathbb{T}}$ . We may assume that  $x(t) > 0$  for  $t \in [t_1, \infty)_{\mathbb{T}}$ , where  $t_1 \in [t_0, \infty)_{\mathbb{T}}$ . Since  $\lim_{t \rightarrow \infty} A(t_0, t) = \infty$ , it is easy to show that  $y(t) > 0$ ,  $t \in [t_1, \infty)_{\mathbb{T}}$ . From the second equation in (1.1), we have  $y^\Delta(t) < 0$ ,  $t \in [t_1, \infty)_{\mathbb{T}}$ . Hence,  $\lim_{t \rightarrow \infty} y(t) \geq 0$ . It follows from the first equation in (1.1) that  $x^\Delta(t) > 0$ ,  $t \in [t_1, \infty)_{\mathbb{T}}$ , and  $\lim_{t \rightarrow \infty} x(t) = \infty$  by Theorem 2.1. Integrating the second equation in (1.1) from  $t$  to  $\infty$  yields that

$$y(t) \geq \int_t^{\infty} f(s, x^\sigma(s))\Delta s, \quad t \in [t_1, \infty)_{\mathbb{T}}. \quad (3.4)$$

By (3.4), (H<sub>2</sub>) and in view of nondecreasing  $\varphi$ , it follows that

$$\frac{x^\Delta(t)}{g(\varphi(x^\sigma(t)))} = \frac{a(t)g(y(t))}{g(\varphi(x^\sigma(t)))} \geq \frac{a(t)g\left(\int_t^{\infty} f(s, x^\sigma(s))\Delta s\right)}{g(\varphi(x^\sigma(t)))} \geq ka(t)g\left(\int_t^{\infty} \frac{f(s, x^\sigma(s))}{\varphi(x^\sigma(s))}\Delta s\right) \quad (3.5)$$

for  $t \in [t_1, \infty)_{\mathbb{T}}$ .

Since (H<sub>4</sub>) holds and  $\lim_{t \rightarrow \infty} x(t) = \infty$ , there is  $t_2 \in [t_1, \infty)_{\mathbb{T}}$  and  $l > 0$  such that

$$\frac{f(t, x^\sigma(t))}{\varphi(x^\sigma(t))} \geq f(t, l) \quad (3.6)$$

for  $t \in [t_2, \infty)_{\mathbb{T}}$ . From (3.5) and (3.6), we get

$$\frac{x^\Delta(t)}{g(\varphi(x^\sigma(t)))} \geq ka(t)g\left(\int_t^{\infty} f(s, l)\Delta s\right). \quad (3.7)$$

Integrating (3.7) from  $t_2$  to  $t$ , we have

$$\int_{t_2}^t \frac{x^\Delta(s)}{g(\varphi(x^\sigma(s)))} \Delta s \geq k \int_{t_2}^t a(s) g\left(\int_s^\infty f(\tau, l) \Delta \tau\right) \Delta s. \quad (3.8)$$

Since  $g$ ,  $\varphi$ , and  $x$  are nondecreasing, we obtain

$$\int_{t_2}^t \frac{x^\Delta(s)}{g(\varphi(x^\sigma(s)))} \Delta s \leq \int_{t_2}^t \frac{x^\Delta(s)}{g(\varphi(x(s)))} \Delta s. \quad (3.9)$$

By (H<sub>4</sub>), (3.8) and (3.9), we get

$$k \int_{t_2}^t a(s) g\left(\int_s^\infty f(\tau, l) \Delta \tau\right) \Delta s \leq \int_{x(t_2)}^{x(t)} \frac{du}{g(\varphi(u))} < \infty, \quad (3.10)$$

which contradicts (3.3) when  $t \rightarrow \infty$ . The proof is complete.  $\square$

**Theorem 3.2.** *Suppose that  $\lim_{t \rightarrow \infty} A(t_0, t) = \infty$  and (H<sub>5</sub>) holds. Suppose further that  $f(t, u)$  is nondecreasing in  $u$  for each fixed  $t \in [t_0, \infty)_{\mathbb{T}}$  and  $g$  is nondecreasing. Then system (1.1) is oscillatory if and only if for all  $c \neq 0$*

$$\int_{t_0}^\infty |f(t, cA(t_0, \sigma(t)))| \Delta t = \infty. \quad (3.11)$$

*Proof.* If (3.11) does not hold, by Theorem 2.3, system (1.1) has a nonoscillatory solution  $(x(t), y(t))$  such that  $\lim_{t \rightarrow \infty} (x(t)/A(t_0, t)) = \alpha \neq 0$  and  $\lim_{t \rightarrow \infty} y(t) = \beta \neq 0$ .

Conversely, suppose that (3.11) holds and that system (1.1) has a nonoscillatory solution  $(x(t), y(t))$  for  $t \in [t_0, \infty)_{\mathbb{T}}$ . We assume that  $x(t) > 0$  for  $t \in [t_1, \infty)_{\mathbb{T}}$ , where  $t_1 \in [t_0, \infty)_{\mathbb{T}}$ . Then by same argument in the proof of Theorem 3.1, we have  $x^\Delta(t) > 0$ ,  $y^\Delta(t) < 0$ ,  $y(t) > 0$  eventually. We claim that (3.11) implies  $\lim_{t \rightarrow \infty} y(t) = 0$ . In fact, if  $\lim_{t \rightarrow \infty} y(t) = \beta > 0$ , then  $y(t) \geq \beta$  for  $t \in [t_1, \infty)_{\mathbb{T}}$ . According to the first equation in (1.1), we get

$$\begin{aligned} x^\sigma(t) &= x(t_1) + \int_{t_1}^{\sigma(t)} a(s) g(y(s)) \Delta s \\ &\geq \int_{t_1}^t a(s) g(y(s)) \Delta s + \int_t^{\sigma(t)} a(s) g(y(s)) \Delta s \\ &\geq g(y(t)) [A(t_1, t) + \mu(t) a(t)] = g(y(t)) A(t_1, \sigma(t)) \geq g(\beta) A(t_1, \sigma(t)). \end{aligned} \quad (3.12)$$

Integrating the second equation in (1.1) from  $t_1$  to  $\infty$ , we have

$$\beta - y(t_1) = - \int_{t_1}^\infty f(s, x^\sigma(s)) \Delta s \leq - \int_{t_1}^\infty f(s, g(\beta) A(t_1, \sigma(s))) \Delta s = -\infty, \quad (3.13)$$

which is a contradiction. Hence,  $\lim_{t \rightarrow \infty} y(t) = 0$ .

By (H<sub>5</sub>), we have

$$kf(t, A(t_1, \sigma(t))) \leq kf\left(t, \frac{x^\sigma(t)}{g(y(t))}\right) \leq \frac{f(t, x^\sigma(t))}{\varphi(g(y(t)))} = \frac{-y^\Delta(t)}{\varphi(g(y(t)))}. \quad (3.14)$$

From (3.14), it follows

$$\int_{t_1}^t kf(s, A(t_1, \sigma(s)))\Delta s \leq - \int_{t_1}^t \frac{y^\Delta(s)}{\varphi(g(y(s)))} \Delta s. \quad (3.15)$$

Hence,

$$k \int_{t_1}^t f(s, A(t_1, \sigma(s)))\Delta s \leq - \int_{y(t_1)}^{y(t)} \frac{du}{\varphi(g(u))}. \quad (3.16)$$

In view of (H<sub>5</sub>) and (3.11), this is a contradiction. The proof is complete. □

*Remark 3.3.* Theorems 3.1 and 3.2 improve the existing results of [15, 18].

*Example 3.4.* Consider the system:

$$x^\Delta(t) = |y(t)|^{1/\alpha-1}y(t), \quad y^\Delta(t) = -\frac{t^v|x^\sigma(t)|^{\gamma-1}x^\sigma(t)}{1+t^u|x^\sigma(t)|^m}, \quad (3.17)$$

where  $\mathbb{T} = a\mathbb{N} = \{an \mid n \in \mathbb{N}\}$ ,  $a, m, \gamma, u, \alpha > 0$  and  $v$  are constants as well as  $\gamma > m$ .

Let

$$a(t) = 1, \quad g(y) = |y|^{(1/\alpha)-1}y, \quad f(t, x) = \frac{t^v|x|^{\gamma-1}x}{1+t^u|x|^m}. \quad (3.18)$$

It is easy to see that  $g(y)$  is increasing and for  $0 < l \leq x \leq L$ ,  $\gamma \geq m$ ,

$$\begin{aligned} f(t, l) &\leq f(t, x) \leq f(t, L), \\ f(t, lt) &\leq f(t, xt) \leq f(t, Lt). \end{aligned} \quad (3.19)$$

For  $u > v + 2\alpha$ , we have

$$\begin{aligned}
\int_{t_0}^{\infty} a(t)g\left(\int_t^{\infty} |f(s,c)|\Delta s\right)\Delta t &= \int_{t_0}^{\infty} \left(\int_t^{\infty} \frac{s^v|c|^\gamma}{1+s^u|c|^m}\Delta s\right)^{1/\alpha} \Delta t \\
&\leq |c|^{(\gamma-m)/\alpha} \int_{t_0}^{\infty} \left(\int_t^{\infty} s^{v-u}\Delta s\right)^{1/\alpha} \Delta t \\
&\leq |c|^{(\gamma-m)/\alpha} a^{(v-u+\alpha+1)/\alpha} \sum_{n=n_0}^{\infty} \sum_{r=n}^{\infty} r^{(v-u)/\alpha} \\
&= |c|^{(\gamma-m)/\alpha} a^{(v-u+\alpha+1)/\alpha} \sum_{n=n_0}^{\infty} n^{(v-u+\alpha)/\alpha} < \infty,
\end{aligned} \tag{3.20}$$

that is, (2.25) holds. By Corollary 2.2, system (3.17) has a nonoscillatory solution  $(x(t), y(t))$  such that  $\lim_{t \rightarrow \infty} x(t) \neq 0$  and  $\lim_{t \rightarrow \infty} y(t) = 0$ .

On the other hand, For  $u + m > v + \gamma + 1$ , we obtain

$$\begin{aligned}
\int_a^{\infty} |f(t, cA(a, \sigma(t)))| \Delta t &= \int_a^{\infty} \frac{t^{v+\gamma}|c|^\gamma}{1+t^{u+m}|c|^m} \Delta t \\
&\leq |c|^{(\gamma-m)} \int_a^{\infty} t^{v+\gamma-u-m} \Delta t \\
&= |c|^{(\gamma-m)} a^{v+\gamma-u-m+1} \sum_{n=1}^{\infty} n^{v+\gamma-u-m} < \infty.
\end{aligned} \tag{3.21}$$

Hence, (2.26) holds. By Theorem 2.3, system (3.17) has a nonoscillatory solution  $(x(t), y(t))$  such that  $\lim_{t \rightarrow \infty} (x(t)/t) \neq 0$  and  $\lim_{t \rightarrow \infty} y(t) \neq 0$ .

*Example 3.5.* Consider the system:

$$x^\Delta(t) = \frac{1}{t}y^5(t), \quad y^\Delta(t) = -\frac{t^3|x^\sigma(t)|^{4/3}x^\sigma(t)}{1+t^3(x^\sigma(t))^2}, \tag{3.22}$$

where  $\mathbb{T} = a^{\mathbb{N}_0}$ ,  $\mathbb{N}_0 = \{0\} \cup \mathbb{N}$  and  $a > 1$ .

Let

$$a(t) = \frac{1}{t}, \quad g(y) = y^5, \quad f(t, x) = \frac{t^3|x|^{4/3}x}{1+t^3x^2}. \tag{3.23}$$

Obviously,  $f(t, x)$  is increasing in  $x$  for fixed  $t$ , and taking  $\varphi(u) = |u|^{-2/3}u$ , we have

$$\begin{aligned}
|f(t, x)| &\geq |f(t, \operatorname{sgn} x)| |x|^{1/3}, \quad |x| \geq 1, \\
\int^{\pm\infty} \frac{du}{g(\varphi(u))} &= \int^{\pm\infty} \frac{du}{u^{5/3}} < \infty.
\end{aligned} \tag{3.24}$$

On the other hand, we obtain

$$\begin{aligned} \int_1^\infty a(t)g\left(\int_t^\infty |f(s,c)|\Delta s\right)\Delta t &= \int_1^\infty \frac{1}{t}\left(\int_t^\infty \frac{s^3|c|^{7/3}}{1+s^3|c|^2}\Delta s\right)^5\Delta t \\ &\leq (a-1)^6|c|^{35/3}\sum_{n=0}^\infty\left(\sum_{r=n}^\infty \frac{a^{4r}}{1+a^{3r}|c|^2}\right)^5 = \infty, \end{aligned} \quad (3.25)$$

that is, (3.3) holds. Hence, system (3.22) is oscillatory by Theorem 3.1.

*Example 3.6.* Consider the system:

$$x^\Delta(t) = y(t), \quad y^\Delta(t) = -b(t)|x^\sigma(t)|^\lambda \operatorname{sgn} x^\sigma(t) \quad (3.26)$$

on a time scale  $\mathbb{T}$  which contains only isolated points and is unbounded above. Here,  $a(t) = 1$ ,  $g(y) = y$ ,  $0 < \lambda < 1$ ,  $f(t, x) = b(t)|x|^\lambda \operatorname{sgn} x$ ,  $b(t)$  is a nonnegative rd-continuous function on  $[t_0, \infty)_{\mathbb{T}}$ .

We take  $\varphi(x) = |x|^{\lambda-1}x$ ,  $0 < |x| \leq 1$ , then all conditions of Theorem 3.2 are satisfied. Hence, system (3.26) is oscillatory if and only if

$$\int_{t_0}^\infty \sigma^\lambda(t)b(t)\Delta t = \infty. \quad (3.27)$$

On the other hand, system (3.26) can be written in the Emden-Fowler equation:

$$x^{\Delta\Delta}(t) + b(t)|x^\sigma(t)|^\lambda \operatorname{sgn} x^\sigma(t) = 0. \quad (3.28)$$

Since we do not assume that  $\lambda$  is a quotient of odd positive integers, (3.28) includes the equation studied in [21]. Theorem 3.2 generalizes and improves Theorem 7 of [21].

## Acknowledgment

The authors sincerely thank the reviewers for their valuable suggestions and useful comments that have lead to the present improved version of the original paper.

## References

- [1] S. Hilger, "Analysis on measure chains—a unified approach to continuous and discrete calculus," *Results in Mathematics*, vol. 18, no. 1-2, pp. 18–56, 1990.
- [2] R. Agarwal, M. Bohner, D. O'Regan, and A. Peterson, "Dynamic equations on time scales: a survey," *Journal of Computational and Applied Mathematics*, vol. 141, no. 1-2, pp. 1–26, 2002.
- [3] M. Bohner and A. Peterson, *Dynamic Equations on Time Scales: An Introduction with Applications*, Birkhäuser, Boston, Mass, USA, 2001.
- [4] R. P. Agarwal, W.-T. Li, and P. Y. H. Pang, "Asymptotic behavior of nonlinear difference systems," *Applied Mathematics and Computation*, vol. 140, no. 2-3, pp. 307–316, 2003.

- [5] A. Lomtadze and N. Partsvania, "Oscillation and nonoscillation criteria for two-dimensional systems of first order linear ordinary differential equations," *Georgian Mathematical Journal*, vol. 6, no. 3, pp. 285–298, 1999.
- [6] J. R. Graef and E. Thandapani, "Oscillation of two-dimensional difference systems," *Computers & Mathematics with Applications*, vol. 38, no. 7-8, pp. 157–165, 1999.
- [7] W.-T. Li, "Classification schemes for nonoscillatory solutions of two-dimensional nonlinear difference systems," *Computers & Mathematics with Applications*, vol. 42, no. 3–5, pp. 341–355, 2001.
- [8] J. D. Mirzov, *Asymptotic Behaviour of Solutions of Systems of Non-Linear Non-Autonomous Ordinary Differential Equations*, Maikop, Russia, 1993.
- [9] J. Jiang and X. Li, "Oscillation and nonoscillation of two-dimensional difference systems," *Journal of Computational and Applied Mathematics*, vol. 188, no. 1, pp. 77–88, 2006.
- [10] E. Akin-Bohner, M. Bohner, and S. H. Saker, "Oscillation criteria for a certain class of second order Emden-Fowler dynamic equations," *Electronic Transactions on Numerical Analysis*, vol. 27, pp. 1–12, 2007.
- [11] Y. Şahiner, "Oscillation of second-order delay differential equations on time scales," *Nonlinear Analysis: Theory, Methods and Applications*, vol. 63, no. 5–7, pp. e1073–e1080, 2005.
- [12] D.-X. Chen, "Oscillation and asymptotic behavior for  $n$ th-order nonlinear neutral delay dynamic equations on time scales," *Acta Applicandae Mathematicae*, vol. 109, no. 3, pp. 703–719, 2010.
- [13] L. Erbe and A. Peterson, "Recent results concerning dynamic equations on time scales," *Electronic Transactions on Numerical Analysis*, vol. 27, pp. 51–70, 2007.
- [14] L. Erbe and R. Mert, "Some new oscillation results for a nonlinear dynamic system on time scales," *Applied Mathematics and Computation*, vol. 215, no. 7, pp. 2405–2412, 2009.
- [15] S.-C. Fu and M.-L. Lin, "Oscillation and nonoscillation criteria for linear dynamic systems on time scales," *Computers & Mathematics with Applications*, vol. 59, no. 8, pp. 2552–2565, 2010.
- [16] Y. Xu and Z. Xu, "Oscillation criteria for two-dimensional dynamic systems on time scales," *Journal of Computational and Applied Mathematics*, vol. 225, no. 1, pp. 9–19, 2009.
- [17] L. Erbe and R. Mert, "Oscillation for a nonlinear dynamic system on time scales," *Journal of Difference Equations and Applications*, vol. 17, no. 9, pp. 1333–1350, 2011.
- [18] D. R. Anderson, "Oscillation and nonoscillation criteria for two-dimensional time-scale systems of first-order nonlinear dynamic equations," *Electronic Journal of Differential Equations*, vol. 24, pp. 1–13, 2009.
- [19] Z.-Q. Zhu and Q.-R. Wang, "Existence of nonoscillatory solutions to neutral dynamic equations on time scales," *Journal of Mathematical Analysis and Applications*, vol. 335, no. 2, pp. 751–762, 2007.
- [20] M. Bohner and A. Peterson, *Advances in Dynamic Equations on Time Scales*, Birkhäuser, Boston, Mass, USA, 2003.
- [21] E. Akin-Bohner and J. Hoffacker, "Oscillation properties of an Emden-Fowler type equation on discrete time scales," *Journal of Difference Equations and Applications*, vol. 9, no. 6, pp. 603–612, 2003.



# Hindawi

Submit your manuscripts at  
<http://www.hindawi.com>

