

RESEARCH ARTICLE

Open Access

Effect of apoptosis in neural stem cells treated with sevoflurane

Jianlei Qiu^{1,2}, Pengcai Shi³, Wude Mao⁴, Yuyi Zhao¹, Wenshuai Liu⁵ and Yuelan Wang^{2,3*}

Abstract

Background: At present, sevoflurane inhalation anesthesia used on infants is well-known. But long-time exposure to inhalation anesthetic could cause neurologic disorder, especially nerve degeneration in infant and developing brain. The central nervous system degeneration of infants could affect the memory and cognitive function. γ -Aminobutyric acid (GABA) is a known inhibitory neurotransmitter in central nervous system. Inhalation anesthetic sevoflurane may activate GABA_A receptor to inhibit central nervous system, leading to apoptosis of neural degeneration, cognitive dysfunction in the critical period of brain development.

Methods: Neural stem cells were derived from Wistar embryos, cultured *in vitro*. Third generation of neural stem cells were randomly divided into four groups according to cultured suspension: Sevoflurane group (Group S), GABA_A receptor antagonists, Bicuculline group (Group B), Sevoflurane + GABA_A receptor antagonists, Bicuculline group (Group S + B), dimethyl sulphoxide (DMSO) group (Group D). Group B and Group D did not receive sevoflurane preconditioning. Group S and Group S + B were pretreated with 1 minimum alveolar concentration (MAC) sevoflurane for 0 h, 3 h, 6 h, and 12 h. Group S + B and Group B were pretreated with bicuculline (10 μ M). Group D was treated with DMSO (10 μ L/mL). After treatments above, all groups were cultured for 48 h. Then we measured the cells viability by Cell Counting Kit (CCK-8) assay, cytotoxicity by Lactate Dehydrogenase (LDH) assay, apoptosis ratio with Annexin V/propidium iodide (PI) staining by flow cytometry, and the expression of GABA_AR, anti-apoptotic protein Bcl-2, pro-apoptotic protein Bax and Caspase-3 by western blotting.

Results: After exposing to sevoflurane for 0 h, 3 h, 6 h, and 12 h with 1MAC, we found that cell viability obviously decreased and cytotoxicity increased in time-dependent way. And Annexin V/PI staining indicated increased apoptosis ratio by flow cytometry. The protein level of GABA_A receptor, pro-apoptotic protein Bax and apoptosis protein Caspase-3 increased; while anti-apoptotic protein Bcl-2 decreased. And bicuculline could reverse all detrimental results caused by sevoflurane.

Conclusion: Sevoflurane can inhibit the central nervous system by activating GABA_A, resulting in apoptosis of neural stem cells, thus leading to the NSCs degeneration.

Keywords: Sevoflurane, γ -Aminobutyric acid, Apoptosis, Neural Stem Cells

Background

Sevoflurane anesthesia in infant rats can result in long-term cognitive impairment, possibly by inhibiting neurogenesis [1]. The hippocampus is critical for memory consolidation and is one of only two mammalian brain regions where neural stem cells (NSCs) are renewed continuously throughout life. Thereby, further studies

with sevoflurane exposure of NSCs are necessary to confirm whether sevoflurane can influence apoptosis of neural cells.

GABA is an important endogenous amino acid of the central nervous system [2,3], which could activate GABA_A receptor to produce inhibitory effects in the release of neurotransmitter [4]. Sevoflurane repress spontaneous pre-motor neuronal activity by enhancing GABA_A receptor function [5] thus enhancing the inhibitory effects of central nervous system and further inducing apoptosis of neural stem cells.

* Correspondence: wylgdf@163.com

²School of Medicine, Shandong University, Ji'nan, Shandong, China

³Department of Anesthesiology, Shandong Provincial Qianfoshan Hospital, Ji'nan, Shandong, China

Full list of author information is available at the end of the article

Apoptosis is directly regulated by Bcl-2 family proteins, including anti-apoptosis protein Bcl-2 and Bcl-xL and pro-apoptotic protein Bax and Bak [6], which finally lead to the activation of Caspase-3 [7]. However, whether neural stem cells exposed to sevoflurane cause significance of apoptosis is still unknown. In present study, we use NSCs exposed to sevoflurane to simulate infant sevoflurane anesthesia. Exploration of the mechanisms underneath can provide a novel way to protect our central nervous system against sevoflurane.

Methods

Materials

Cells Counting Kit-8 (CCK-8) and Annexin V-FITC Apoptosis Detection Kit were from Dojindo (Japan). Cells culture medium (DMEM/F12), B-27[®] Supplement without Vitamin A, and fetal bovine serum, were from Gibco (USA). Lactate Dehydrogenase (LDH) was from Beyotime (China). Epidermal Growth Factor (EGF), Fibroblast Growth Factor-basic (bFGF) were from PeproTech (USA).

The following antibodies were purchased as indicated: Rabbit anti-bcl-2 antibody, Mouse anti-bax antibody, Rabbit anti-caspase-3 and Rabbit anti-GAPDH polyclonal antibody were all from Abcam (USA). Rabbit anti-GABA_A R α was from Santa Cruz (USA). Bicuculline was from Sigma (USA).

NSCs culture

All experimental procedures were approved by the Ethics Committee of Dezhou People's Hospital. Every effort was made to minimize the number of animals used and the suffering of the animals. All animals were used only once.

NSCs were derived from Wistar rat embryonic brains, cultured, and expanded. The rat embryos were decapitated on embryonic day 17 (E17), and the hippocampus was quickly removed and placed on a dish with ice cold Hank's balanced salt solution containing penicillin (100 U/ml)/streptomycin (100 μ g/ml) [8,9]. Next, the hippocampus was minced, incubated in 0.25% trypsin-ethylenediaminetetraacetic acid (trypsin/EDTA) for 5 min at 37°C and then incubated with 0.05% DNase for 5 min at 37°C. Minced cells were then centrifuged, and the cell pellet was washed twice with Dulbecco's phosphate buffered saline (DPBS). The isolated cells were seeded at 2×10^4 cells/ml on 100-mm dishes with medium mixture F-12 (DMEM/F12, 1:1), 2% B-27, 1% penicillin (100 U/ml)/streptomycin (100 μ g/ml), 20 ng/ml bFGF, 20 ng/ml EGF, maintained at 37°C in a 5% CO₂ atmosphere. Medium, growth factors, and supplements were changed every 3 days during culture process.

Sevoflurane and bicuculline treatment

NSCs culture plates were put into an airtight glass chamber with inlet and outlet connectors. The inlet port of the chamber was connected to an anesthesia machine (Cicero-EM 8060, Drager, Germany). Sevoflurane was delivered into chamber by a sevoflurane vaporizer attached to the anesthesia machine. The concentrations of sevoflurane in the chamber were detected at the chamber outlet port by using a gas monitor (PM 8060, Drager, Germany). NSCs were divided into four groups: sevoflurane (1MAC) group (Group S), bicuculline group (Group B), and a combination of sevoflurane and bicuculline group (Group S + B), dimethyl sulphoxide (Group D). The Group S was treated with sevoflurane (1MAC) mixed with 95% air/5% CO₂ at 6 L/min for 0 h, 3 h, 6 h, and 12 h. The Group B was treated with bicuculline (10 μ M) and 95% air/5% CO₂ at 6 L/min for 0 h, 3 h, 6 h, and 12 h. The Group S + B was treated with bicuculline (10 μ M) and sevoflurane mixed with 95% air/5% CO₂ at 6 L/min for 0 h, 3 h, 6 h, and 12 h. The Group D was treated with dimethyl sulphoxide (10 μ L/mL) for 0 h, 3 h, 6 h, and 12 h.

CCK-8 assay and lactate dehydrogenase (LDH) release assay

A CCK-8 10 μ l of the kit reagent was added to the live cells into 96-well plates, and the suspension NSCs (100 μ l/well) were incubated for 3 h at 37°C 5% CO₂. Cell viability was measured by the absorbance at 450 nm [1].

A colorimetric assay kit was used to quantify the LDH released from the cultured NSCs. After treated with sevoflurane, bicuculline, and DMSO, NSCs were collected, then centrifuged at 400 g for 5 min. Suck the supernatant, and add 150 μ L LDH release reagent to mix completely, then incubate at 37°C 5% CO₂ for 1 h. After centrifuged at 400 g for 5 min again, suck the supernatant. Add 120 μ L supernatant to 96-well plates. Cell cytotoxicity was measured by the absorbance at 490 nm. All results were normalized to the optical densities (OD) [1].

Measurement of apoptosis by Annexin V/PI staining

Annexin V/PI staining was measured with an FITC Annexin V Apoptosis Detection Kit [10]. After treated with sevoflurane and Bicuculline, the NSCs were washed twice with PBS, and the concentration was adjusted to 1×10^6 cells/ml with Annexin V binding solution. Then remove 195 μ l cell suspensions to a new tube, add 5 μ l Annexin V-FITC combination, add 10 μ l PI solution. Cells were incubated in dark room for 15 min at room temperature before adding 300 μ l Annexin V binding solution. Apoptosis ratio was measured by flow cytometry.

Western blotting analysis

For the preparation of total cell extracts, samples were washed twice with cold PBS and lysed in appropriate amount of radio immunoprecipitation assay (RIPA) buffer supplemented with phenylmethanesulfonyl fluoride (PMSF) [11-13]. The lysate was collected and protein concentration was determined using a bicinchoninic acid protein assay kit. Equal amounts of protein were denatured and separated on 10% SDS-PAGE gels and then transferred to polyvinylidene difluoride membranes (Bio-Rad, Hercules, CA) at 100 V for 1 h. After blocking with skim milk (5%), proteins were incubated with respective primary antibodies in blocking solution, according to the manufacturer’s recommendations. The appropriate horseradish peroxidase-conjugated secondary antibody was added to the filters followed by incubation for 2 h at room temperature with an appropriate dilution. After sequential washing of the membranes in T-PBS to remove excess secondary antibody, the signals were detected by chemiluminescence using the electrochemiluminescence (ECL) system. Relative band densities of the

various proteins were measured from scanned films using Image J Software.

Statistical analysis

Representative experiments from at least three independent experiments are shown. Statistical analysis was performed using the SPSS 19.0 statistics package. All data are expressed as mean ± SD. Statistical differences were assessed using Student’s t-tests or one-way analysis of variance (ANOVA), where appropriate among groups. A *P*-value <0.05 was considered statistically significant.

Results

Viability of NSCs with sevoflurane by CCK-8 assays

To evaluate the viability of NSCs after sevoflurane treatment, the NSCs were incubated in 1MAC of sevoflurane for 0 h, 3 h, 6 h, and 12 h, and cell viability was measured using the CCK-8 assays. Compared with 0 h, the viability of NSCs exposed to sevoflurane at 1 MAC decreased in time-dependent way in Group S. Cell viability decreased 8% (*P* <0.05) (Figure 1A) at 3 h post exposure, 15% at




Figure 1 Effects of sevoflurane on cell viability and cytotoxicity. NSCs were exposed to sevoflurane with 1 MAC at 0, 3, 6, and 12 h. **A:** Cell viability was measured by CCK-8 assays. Compared with 0 h in Group S, **P* <0.05; Compared with 0 h in Group B, &*P* >0.05; Compared with 0 h in Group D, #*P* >0.05; compared with the corresponding time point in Group S, ###*P* <0.05, &&*P* <0.05 and ***P* <0.05 in Group S+B. **B:** Cytotoxicity was measured by LDH assays. Compared with 0 h in Group S, **P* <0.05; Compared with 0 h in Group B, &*P* >0.05; Compared with 0 h in Group D, #*P* >0.05; compared with the corresponding time point in Group S, ###*P* <0.05, &&*P* <0.05, ***P* <0.05 in Group S+B. Data are representative of 6 independent experiments.

6 h ($P < 0.05$) (Figure 1A), and 28% at 12 h ($P < 0.05$) (Figure 1A). Compared with corresponding time point of Group S, the viability of NSCs increased in Group S + B ($P < 0.05$) (Figure 1A). Compared with 0 h, the viability of NSCs exposed to DMSO for 3 h, 6 h, and 12 h did not change obviously in Group D ($P > 0.05$) (Figure 1A). Compared with 0 h, the viability of NSCs exposed to bicuculline for 3 h, 6 h, and 12 h did not change obviously in Group B ($P > 0.05$) (Figure 1A).

Cytotoxicity of NSCs with Sevoflurane by LDH assays

Cytotoxicity of Sevoflurane on NSCs was measured by LDH assays. LDH release in the culture media increased in Group S along with the time from 0 hour to 12 h. Cytotoxicity increased 1.2-fold at 3 h, 1.4-fold at 6 h ($P < 0.05$) (Figure 1B), and 1.8-fold at 12 h ($P < 0.05$) (Figure 1B). LDH release in the culture media was decreased in Group S + B compared with corresponding time of Group S, ($P < 0.05$) (Figure 1B). Compared with 0 h, the cytotoxicity of NSCs exposed to DMSO for 3 h, 6 h, and 12 h did not change obviously in Group D ($P > 0.05$) (Figure 1B). Compared with 0 h, the viability of NSCs exposed to bicuculline for 3 h, 6 h, and 12 h did not change obviously in Group B ($P > 0.05$) (Figure 1A).

Apoptosis of NSCs with sevoflurane by annexin V/PI staining

Annexin V/PI staining was measured by flow cytometry (Figure 2). The results were similar to the CCK-8 assay. Compared with 0 h, the NSCs exposed to sevoflurane for 3 to 12 hours displayed a decreased proportion of early and late apoptotic cells in Group S ($P < 0.05$) (Figure 2), and the ratio of apoptosis could be reversed by bicuculline at all times points in Group S + B ($P < 0.05$) (Figure 2).

Expression of GABA_A receptor in group S and group S + B

Activation of GABA_A receptor in response to sevoflurane exposure was reported to decrease the viability of NSCs. The protein level of GABA_A receptor in NSCs was measured by western blotting. The expression of GABA_A receptor exposed to sevoflurane was higher in Group S and Group S + B, compared with 0 h ($P < 0.05$) (Figure 3). The expression of GABA_A receptor was higher in Group S compared with Group S + B pre-treated with bicuculline for 2 h at 3 h, 6 h, and 12 h ($P < 0.05$) (Figure 3).

Expression of Bcl-2, Bax and caspase-3 with the treatments of sevoflurane and bicuculline

Western blotting showed that exposure to 1MAC sevoflurane in Group S induced the down regulation of Bcl-2 (Figure 4), and up regulation of Bax (Figure 4) and Caspase-3 (Figure 5), compared with 0 h ($P < 0.05$). In Group S + B, the changes of Bcl-2, Bax and Caspase-3 expression was a relatively mild compared with Group S ($P < 0.05$).

Discussion

Sevoflurane is a new type of fluorinated inhalation anesthetic allowing for more rapid emergence than previous drugs, due to its lower blood/gas solubility [14,15]. Nowadays, more and more infants adopt sevoflurane inhalation anesthesia, while a series of problems arises. Recent studies showed that young children exposed to general anesthesia are at greater risk of learning deficits in adolescence [16]. It showed that postnatal exposure of common sevoflurane anesthetic could cause widely neuronal apoptosis subsequent learning and memory disorders [17].





Figure 2 Bicuculline increases neuronal viability and reduces apoptosis in sevoflurane neurotoxicity. NSCs were incubated with bicuculline before sevoflurane treatment. The cells were then labeled by Annexin V-FITC and propidium iodide (PI) in Group S and Group S + B. The apoptosis was determined by flow cytometry. **A:** Q1 and Q3 quadrants were represented for normal cells and death cells with red color; Q2 and Q4 quadrants were represented for viable apoptotic cells and non-viable apoptotic cells with blue color. **B:** The histogram represents the percentage of apoptotic cells. Compared with 0 h in Group S, * $P < 0.05$. In Group S + B, compared with the corresponding time point in Group S, # $P < 0.05$, ## $P < 0.05$ and & $P < 0.05$. Data are representative of 3 independent experiments.



The study of Statmann et al. demonstrated that early inhalational anesthesia can induce long-term, mainly hippocampus-dependent cognitive dysfunction in rodents [18]. Postnatal hippocampal neurogenesis sustains hippocampal function throughout life [19], thus protection of the hippocampal neurogenesis is very significant. Cognitive impairment following anesthetic exposure may be associated with suppression of survival, proliferation of hippocampal neural stem cells. In this study, we confirmed in rat hippocampus that the viability of NSCs exposed to sevoflurane decreased, and cytotoxicity of sevoflurane increased.

Previous studies have shown that inhalational anesthetics, such as sevoflurane and isoflurane, are very helpful for reduction of perioperative mortality [20,21]. But sevoflurane anesthetics may also cause memory impairment by

neurons lost in hippocampus through cells apoptosis [17]. And, recent reports also showed that propofol and sevoflurane would trigger widespread apoptotic neurodegeneration throughout the developing brain [22-24]. However, the mechanism remained unknown. Our studies approved that exposure to sevoflurane could cause apoptosis of NSCs, and apoptosis rate increased in a time-dependent way.

Many researches had given us possible mechanisms of anesthetic-induced apoptosis of NSCs. Some anesthetic agents induce apoptosis by blocking N-methyl-D-aspartate (NMDA) receptor and activating GABA_A receptor [25]. As a combination of NMDA receptor antagonist and GABA_A receptor agonists, inhalation anesthetic kills immature brain cells [26,27]. But the mechanisms of sevoflurane-induced apoptosis of NSCs are currently unknown. In our study, we found that the expression of GABA_A reporters increased after exposure to sevoflurane in time-dependent way by western blotting. That is to say, sevoflurane could activate the GABA_A receptors. Therefore, we assume that stimulation of GABA by consistently duration of sevoflurane may induce apoptosis.

Bcl-2 protein family is an important factor to regulate cell intrinsic pathway of apoptosis. The family consists of two members: one is anti-apoptotic protein Bcl-2, the other is pro-apoptotic molecular Bcl-2-associated X protein (Bax). The protein Bax received the apoptosis signal could transfer to mitochondrial outer membrane, open the channel of permeability, release cytochrome c, and induce apoptosis finally. The protein bcl-2 could combine with Bax to change its spatial conformation to improve the apoptosis [28,29]. Expression of Bcl-2 or related anti-apoptotic proteins block cell death in response to many varieties [30]. In sevoflurane treated rat pup, the expression of Bcl-2 was suppressed, suggesting that anesthetic exposure may accelerate the physiological apoptosis [31]. In our study, we found that exposure under sevoflurane could increase the expression of pro-apoptotic molecular Bax, while decrease Bcl-2 expression by western blotting. That is to say, sevoflurane could cause apoptosis of NSCs by increasing Bax and decreasing Bcl-2.

Caspase-3 is a member of the Caspase family of aspartate-specific cysteine proteases that plays a central role in the execution of the apoptotic program [31]. The GABA agonist ethanol activated Caspase-3 in progenitor cells [32]. In sevoflurane treated rat pup, the expression of Caspase-3 increased, suggesting that anesthetic exposure may accelerate the physiological apoptosis [33]. The cellular expression of Bcl-2 and activated Caspases likely mediate the increased vulnerability to neuronal apoptosis during synaptogenesis [34]. Recent report indicated that it could active caspase-3 and decrease neuronal in the neonatal hippocampus, but did not affect subsequent behavioural performances in juvenile rats which exposed to 2.3% sevoflurane for 6 h




Figure 4 Effect of Bcl-2 and Bax expression in Group S and Group S + B. The expressions of Bcl-2 and Bax were detected with exposure to sevoflurane. **A** and **C**: Representative Western blotting of Bcl-2 and Bax expressions in Group S and Group S + B. **B** and **D**: Densitometric analysis of Bcl-2 and Bax expressions relative to that of GAPDH. Compared with 0 h in Group S, *P <0.05. In Group S + B, compared with the corresponding time point in Group S, #P <0.05, ##P <0.05 and &P <0.05. Data are representative of 3 independent experiments.

[35]. Consistently, we found that caspase-3 activated under exposure of sevoflurane by western blotting.

Bicuculline, which acts on the GABA_A receptor recognition site, is specific GABA_A receptor blockers [36]. Thereby it could selectively block GABA_A receptor, inhibit activation of GABA_A. Our study showed that GABA_A receptors could be prevented by bicuculline. Furthermore, bicuculline could block apoptosis of NSCs by increasing expression of Bcl-2 and decreasing expression of Bax and Caspase-3. For the first time, we provide a novel therapeutic way to protect NSCs against sevoflurane anesthesia.

Conclusions

This study indicates that exposure to sevoflurane causes the apoptosis of NSCs, and long-time exposure could aggravate the damage. Activation of GABA_A receptor can induce the apoptosis by increasing Bax and caspase-3 and decreasing Bcl-2. The specific blocker of GABA_A receptor bicuculline could ameliorate the apoptosis. Our results suggest that GABA_A receptor may be an important receptor in NSCs exposing to sevoflurane, inhibition of GABA_A receptor may be a novel and effective target.




Figure 5 Effect of Caspase-3 expression in Group S and Group S + B. The expressions of Bcl-2 and Bax were detected with exposure to sevoflurane. **A**: Representative Western blotting of Caspase-3 expression in Group S and Group S + B. **B**: Densitometric analysis of Caspase-3 expression relative to that of GAPDH. Compared with 0 h in Group S, *P <0.05. In Group S + B, compared with the corresponding time point in Group S, #P <0.05, ##P <0.05 and &P <0.05. Data are representative of 3 independent experiments.

Abbreviations

GABA: γ -Aminobutyric acid; CCK-8: Cell counting kit-8; LDH: Lactate dehydrogenase; PI: Propidium iodide; MAC: Minimum alveolar concentration; NSCs: Neural stem cells; EGF: Epidermal growth Factor; bFGF: Fibroblast growth factor-basic; OD: Optical densities; NMDA: N-methyl-D-aspartate; RIPA: Radio immunoprecipitation assay; PMSF: Phenylmethanesulfonyl fluoride; SDS: Sodium dodecyl sulfate; PAGE: Polyacrylamide gel electrophoresis; ECL: Electrochemiluminescence; ANOVA: Analysis of variance.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

JQ, PS, WM, YZ and WL carried out all the experiments, participated in the molecular biology studies, JQ carried out the immunoassays, participated in performed the statistical analysis and drafted the manuscript. YW participated in the design and conceived of the study, and participated in its design and coordination and helped to draft the manuscript. All authors read and approved the final manuscript.

Acknowledgements

This work was supported by the National Natural Science Foundation of China (81270127). The authors thank Medical Research Center of Qianfoshan Hospital of Shandong Province for equipment support and technical assistance.

Author details

¹Department of Anesthesiology, Dezhou People's Hospital, Dezhou, Shandong, China. ²School of Medicine, Shandong University, Ji'nan, Shandong, China. ³Department of Anesthesiology, Shandong Provincial Qianfoshan Hospital, Ji'nan, Shandong, China. ⁴Department of Anesthesiology, Jiaozhou Central Hospital of Qingdao, Qingdao, Shandong, China. ⁵Department of Emergency, Dezhou People's Hospital, Dezhou, Shandong, China.

Received: 22 September 2014 Accepted: 24 February 2015

Published online: 07 March 2015

References

- Nie H, Peng Z, Lao N, Dong H, Xiong L. Effects of sevoflurane on self-renewal capacity and differentiation of cultured neural stem cells. *Neurochem Res.* 2013;38:1758–67.
- Satin LS, Kinard TA. Neurotransmitters and their receptors in the islets of Langerhans of the pancreas: what messages do acetylcholine, glutamate, and GABA transmit? *Endocrine.* 1998;8:213–23.
- Gerber 3rd JC, Hare TA. Gamma-aminobutyric acid in peripheral tissue, with emphasis on the endocrine pancreas: presence in two species and reduction by streptozotocin. *Diabetes.* 1979;28:1073–6.
- DeFeudis FV. GABA and cardiovascular regulation. *Rev Clin Basic Pharm.* 1985;5:141–58.
- Stucke AG, Zuperku EJ, Krolo M, Brandes IF, Hopp FA, Kampine JP, et al. Sevoflurane enhances gamma-aminobutyric acid type A receptor function and overall inhibition of inspiratory premotor neurons in a decerebrate dog model. *Anesthesiology.* 2005;103:57–64.
- Gui D, Guo Y, Wang F, Liu W, Chen J, Chen Y, et al. Astragaloside IV, a novel antioxidant, prevents glucose-induced podocyte apoptosis *in vitro* and *in vivo*. *PLoS One.* 2012;7:e39824.
- Zhang M, Shan H, Chang P, Wang T, Dong W, Chen X, et al. Hydrogen sulfide offers neuroprotection on traumatic brain injury in parallel with reduced apoptosis and autophagy in mice. *PLoS One.* 2014;9:e87241.
- Choi MR, Jung KH, Park JH, Das ND, Chung MK, Choi IG, et al. Ethanol-induced small heat shock protein genes in the differentiation of mouse embryonic neural stem cells. *Arch Toxicol.* 2011;85:293–304.
- Han D, Choi MR, Jung KH, Kim N, Kim SK, Chai JC, et al. Global transcriptome profiling of genes that are differentially regulated during differentiation of mouse embryonic neural stem cells into astrocytes. *J Mol Neurosci.* 2015;55:109–25.
- Rieger AM, Nelson KL, Konowalchuk JD, Barreda DR. Modified annexin V/propidium iodide apoptosis assay for accurate assessment of cell death. *J Vis Exp.* 2011;24:50.
- Liu M, Gu C, Wang Y. Upregulation of the tight junction protein occludin: effects on ventilation-induced lung injury and mechanisms of action. *BMC Pulm Med.* 2014;14:94.
- Wang Y, Minshall RD, Schwartz DE, Hu G. Cyclic stretch induces alveolar epithelial barrier dysfunction via calpain-mediated degradation of p120-catenin. *Am J Physiol Lung Cell Mol Physiol.* 2011;301:L197–206.
- Zhao T, Liu M, Gu C, Wang X, Wang Y. Activation of c-Src tyrosine kinase mediated the degradation of occludin in ventilator-induced lung injury. *Respir Res.* 2014;15:158.
- Ebert TJ, Robinson BJ, Uhrich TD, Mackenthun A, Pichotta PJ. Recovery from sevoflurane anesthesia: a comparison to isoflurane and propofol anesthesia. *Anesthesiology.* 1998;89:1524–31.
- Robinson BJ, Uhrich TD, Ebert TJ. A review of recovery from sevoflurane anaesthesia: comparisons with isoflurane and propofol including meta-analysis. *Acta Anaesthesiol Scand.* 1999;43:185–90.
- Wildier RT, Flick RP, Sprung J, Katusic SK, Barbaresi WJ, Mickelson C, et al. Early exposure to anesthesia and learning disabilities in a population-based birth cohort. *Anesthesiology.* 2009;110:796–804.
- Satomoto M, Satoh Y, Terui K, Miyao H, Takishima K, Ito M, et al. Neonatal exposure to sevoflurane induces abnormal social behaviors and deficits in fear conditioning in mice. *Anesthesiology.* 2009;110:628–37.
- Stratmann G, Sall JW, May LD, Bell JS, Magnusson KR, Rau V, et al. Isoflurane differentially affects neurogenesis and long-term neurocognitive function in 60-day-old and 7-day-old rats. *Anesthesiology.* 2009;110:834–48.
- Trouche SBB, Roulet P, Rampon C. Recruitment of adult-generated neurons into functional hippocampal networks contributes to updating and strengthening of spatial memory. *Proc Natl Acad Sci U S A.* 2009;106:5919–24.
- Landoni G, Rodseth RN, Santini F, Ponschab M, Ruggeri L, Szekely A, et al. Randomized evidence for reduction of perioperative mortality. *J Cardiothorac Vasc Anesth.* 2012;26:764–72.
- Bignami E, Biondi-Zoccai G, Landoni G, Fochi O, Testa V, Sheiban I, et al. Volatile anesthetics reduce mortality in cardiac surgery. *J Cardiothorac Vasc Anesth.* 2009;23:594–9.
- Tu S, Wang X, Yang F, Chen B, Wu S, He W, et al. Propofol induces neuronal apoptosis in infant rat brain under hypoxic conditions. *Brain Res Bull.* 2011;86:29–35.
- Zhou X, Song FH, He W, Yang XY, Zhou ZB, Feng X, et al. Neonatal exposure to sevoflurane causes apoptosis and reduces nNOS protein expression in rat hippocampus. *Molecular medicine reports.* 2012;6:543–6.
- Pellegrini L, Bennis Y, Velly L, Grandvuillemin I, Pisano P, Bruder N, et al. Erythropoietin protects newborn rat against sevoflurane-induced neurotoxicity. *Paediatr Anaesth.* 2014;24:749–59.
- Fredriksson A, Ponten E, Gordh T, Eriksson P. Neonatal exposure to a combination of N-methyl-D-aspartate and gamma-aminobutyric acid type A receptor anesthetic agents potentiates apoptotic neurodegeneration and persistent behavioral deficits. *Anesthesiology.* 2007;107:427–36.
- Brambrink AM, Evers AS, Avidan MS, Farber NB, Smith DJ, Zhang X, et al. Isoflurane-induced neuroapoptosis in the neonatal rhesus macaque brain. *Anesthesiology.* 2010;112:834–41.
- Stratmann G, May LD, Sall JW, Alvi RS, Bell JS, Ormerod BK, et al. Effect of hypercarbia and isoflurane on brain cell death and neurocognitive dysfunction in 7-day-old rats. *Anesthesiology.* 2009;110:849–61.
- Llambi F, Green DR. Apoptosis and oncogenesis: give and take in the BCL-2 family. *Curr Opin Genet Dev.* 2011;21:12–20.
- Leber B, Geng F, Kale J, Andrews DW. Drugs targeting Bcl-2 family members as an emerging strategy in cancer. *Expert Rev Mol Med.* 2010;12:e28.
- Brunelle JK, Letai A. Control of mitochondrial apoptosis by the Bcl-2 family. *J Cell Sci.* 2009;122:437–41.
- Wang WY, Yang R, Hu SF, Wang H, Ma ZW, Lu Y. N-stearoyl-L-tyrosine ameliorates sevoflurane induced neuroapoptosis via MEK/ERK1/2 MAPK signaling pathway in the developing brain. *Neurosci Lett.* 2013;541:167–72.
- Ieraci A, Herrera DG. Single alcohol exposure in early life damages hippocampal stem/progenitor cells and reduces adult neurogenesis. *Neurobiol Dis.* 2007;26:597–605.
- Gown AM, Willingham MC. Improved detection of apoptotic cells in archival paraffin sections: immunohistochemistry using antibodies to cleaved caspase 3. *J Histochem Cytochem.* 2002;50:449–54.
- Yuan J, Yankner BA. Apoptosis in the nervous system. *Nature.* 2000;407:802–9.

35. Feng X, Liu JJ, Zhou X, Song FH, Yang XY, Chen XS, et al. Single sevoflurane exposure decreases neuronal nitric oxide synthase levels in the hippocampus of developing rats. *Br J Anaesth.* 2012;109:225–33.
36. Palombi PS, Caspary DM. GABAA receptor antagonist bicuculline alters response properties of posteroventral cochlear nucleus neurons. *J Neurophysiol.* 1992;67:738–46.

**Submit your next manuscript to BioMed Central
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

