



# Diuron tolerance and potential degradation by pelagic microbiomes in the Great Barrier Reef lagoon

Florent E. Angly<sup>1</sup>, Olga Pantos<sup>1,2</sup>, Thomas C. Morgan<sup>1</sup>, Virginia Rich<sup>3,4</sup>, Hemerson Tonin<sup>5</sup>, David G. Bourne<sup>5</sup>, Philip Mercurio<sup>5,6</sup>, Andrew P. Negri<sup>5</sup> and Gene W. Tyson<sup>1</sup>

<sup>1</sup> Australian Centre for Ecogenomics, The University of Queensland, St Lucia, Queensland, Australia

<sup>2</sup> Global Change Institute, The University of Queensland, St Lucia, Queensland, Australia

<sup>3</sup> Department of Soil, Water and Environmental Science, The University of Arizona, Tucson, AZ, United States of America

<sup>4</sup> Microbiology Department, The Ohio State University, Columbus, OH, United States of America

<sup>5</sup> Australian Institute of Marine Science, Townsville, Queensland, Australia

<sup>6</sup> National Research Centre for Environmental Toxicology, The University of Queensland, Coopers Plains, Queensland, Australia

## ABSTRACT

Diuron is a herbicide commonly used in agricultural areas where excess application causes it to leach into rivers, reach sensitive marine environments like the Great Barrier Reef (GBR) lagoon and pose risks to marine life. To investigate the impact of diuron on whole prokaryotic communities that underpin the marine food web and are integral to coral reef health, GBR lagoon water was incubated with diuron at environmentally-relevant concentration (8 µg/L), and sequenced at specific time points over the following year. 16S rRNA gene amplicon profiling revealed no significant short- or long-term effect of diuron on microbiome structure. The relative abundance of prokaryotic phototrophs was not significantly altered by diuron, which suggests that they were largely tolerant at this concentration. Assembly of a metagenome derived from waters sampled at a similar location in the GBR lagoon did not reveal the presence of mutations in the cyanobacterial photosystem that could explain diuron tolerance. However, resident phages displayed several variants of this gene and could potentially play a role in tolerance acquisition. Slow biodegradation of diuron was reported in the incubation flasks, but no correlation with the relative abundance of heterotrophs was evident. Analysis of metagenomic reads supports the hypothesis that previously uncharacterized hydrolases carried by low-abundance species may mediate herbicide degradation in the GBR lagoon. Overall, this study offers evidence that pelagic phototrophs of the GBR lagoon may be more tolerant of diuron than other tropical organisms, and that heterotrophs in the microbial seed bank may have the potential to degrade diuron and alleviate local anthropogenic stresses to inshore GBR ecosystems.

Submitted 19 May 2015

Accepted 13 February 2016

Published 8 March 2016

Corresponding author

Florent E. Angly,  
florent.angly@gmail.com

Academic editor

Hauke Smidt

Additional Information and  
Declarations can be found on  
page 13

DOI 10.7717/peerj.1758

© Copyright  
2016 Angly et al.

Distributed under  
Creative Commons CC-BY 4.0

**OPEN ACCESS**

**Subjects** Bioinformatics, Ecology, Genomics, Marine Biology, Microbiology

**Keywords** Diuron, Great barrier reef, Amplicon profiling, Metagenomics, Herbicide, Incubation

## INTRODUCTION

Coral reefs are very rich and diverse ecosystems, though due to both local and global anthropogenic disturbances, they are in a state of gradual decline (*Pandolfi et al., 2003*).

Local impacts such as coastal pollution (Brodie *et al.*, 2012) and overfishing (De'ath *et al.*, 2012) affect both the reef macrobiota (Fabricius, 2005; Sandin *et al.*, 2008) and the microorganisms in their associated microbiomes (Dinsdale *et al.*, 2008; Thurber *et al.*, 2009; Webster *et al.*, 2011). Microorganisms not only underpin the marine food web and system function (Azam *et al.*, 1983; Fenchel, 2008), but also form intimate relationships with corals that are essential for their health (Rohwer *et al.*, 2002; Lesser *et al.*, 2004; Lema, Willis & Bourne, 2012). Hence, changes to microbiome structure and function can compromise the health of coral reefs and their ability to recover from stresses (Ainsworth, Thurber & Gates, 2010; Hughes *et al.*, 2010).

Fishing impact is limited across the Great Barrier Reef (GBR), but its otherwise oligotrophic inshore coastal habitats (Schaffelke *et al.*, 2012) are subject to pollution from land runoff of agricultural, industrial and domestic origin (Packett *et al.*, 2009; Brodie *et al.*, 2012). For example, herbicides that are used to control weeds in the sugarcane plantations of Queensland have been detected in the waterways of the catchments (McMahon *et al.*, 2005; Mitchell, Brodie & White, 2005; Shaw & Müller, 2005), intertidal sediments, and in the waters surrounding inshore coral reefs (Shaw & Müller, 2005; Lewis *et al.*, 2009). The herbicide diuron, or 3-(3,4-dichlorophenyl)-1,1-dimethylurea (DCMU), is consistently detected in the GBR lagoon (Haynes, Müller & Carter, 2000; Shaw *et al.*, 2010; Kennedy *et al.*, 2012b), exceeding the Australian and New Zealand guideline trigger value of 0.2 µg/L at some sites (Smith *et al.*, 2012). Diuron is an effective broad-spectrum herbicide due to its ability to inhibit the photosystem II (PSII) complex of photosynthetic organisms (Metz *et al.*, 1986). Consequently, it poses risks to a wide range of marine eukaryotes including invertebrates (Bellas *et al.*, 2005; Mai *et al.*, 2013), seagrasses (Haynes *et al.*, 2000; Flores *et al.*, 2013), fishes (Mhadhbi & Beiras, 2012), diatoms (Legrand *et al.*, 2006; Magnusson, Heimann & Negri, 2008) and microalgae, both benthic (Magnusson *et al.*, 2012) and in endosymbiotic relationship with corals (Jones *et al.*, 2003; Jones, 2004; Shaw, Brodie & Müller, 2012). Furthermore, diuron's low rate of abiotic degradation by hydrolysis and photo-degradation (Okamura, 2002; Moncada, 2004; Mercurio *et al.*, 2015) results in its accumulation in the marine environment, particularly in sediments (Haynes, Müller & Carter, 2000; Balakrishnan, Takeda & Sakugawa, 2012; Xu *et al.*, 2013).

In addition to its effects on eukaryotes, diuron is also toxic to bacteria. Low concentrations of diuron (1.6–23 µg/L) impairs the photosynthesis of phototrophs such as cyanobacteria (Allen *et al.*, 1983; Deng, Gao & Sun, 2012), while very high concentrations ( $1 \times 10^6$  µg/L) inhibit the growth of bacterial heterotrophs (Fajà *et al.*, 2010). Consequently, exposure to a pulse of diuron through a flooding event can significantly alter microbiome structure (Tlili *et al.*, 2008) and decrease microbial abundance (Ricart *et al.*, 2009). Conversely, diuron exposure can also increase bacterial abundance in wastewater treatment ponds (Sumpono *et al.*, 2003), likely due to heterotrophic bacteria taking advantage of the release of organic compounds by organisms susceptible to diuron, such as diatoms (Proia *et al.*, 2011). An alternative explanation is that some bacteria are able to metabolize diuron and use it as an energy source (Dellamatrice & Monteiro, 2004), as seen in soil, sediments and sludge (Cullington & Walker, 1999; Dellamatrice & Monteiro, 2004; Sørensen, Albers & Aamand, 2008; Stasinakis *et al.*, 2009; Pesce *et al.*, 2012). This biodegradation is catalyzed by

phenylurea hydrolases (PuhAB) (*Turnbull et al., 2001; Khurana et al., 2009*) and proceeds faster than degradation by abiotic means (*Cullington & Walker, 1999; Dellamatrice & Monteiro, 2004; Sørensen, Albers & Aamand, 2008; Stasinakis et al., 2009; Pesce et al., 2010*). Microorganisms are therefore regularly employed in herbicide pollution remediation technologies (*Villaverde et al., 2012; Safi, Awad & El-Nahhal, 2014*).

A recent study found that microbial community variations in the GBR lagoon are primarily driven by riverine effluents (*Angly et al., 2016*) and a year-long seawater simulation experiment presented evidence that microorganisms play a role in the biodegradation of a wide range of PSII herbicides in this region (*Mercurio et al., 2015*). Although the community-wide effects of diuron on marine microorganisms are not characterized, this research suggests that pelagic microbiomes may protect coral reef and seagrass ecosystems by degrading this herbicide. In the present study, we hypothesized that diuron significantly affects the composition of these microbiomes by specifically: (i) inhibiting phototrophs in the short-term; and (ii) sustaining the long-term growth of selected heterotrophs that have the potential to metabolize it, leading to the herbicide's disappearance. To investigate these hypotheses, we collected samples from the simulation study of *Mercurio et al. (2015)* and an inshore GBR location, and characterized their microbiomes (Archaea and Bacteria) using high-throughput 16S rRNA amplicon profiling and metagenomics.

## MATERIALS & METHODS

### Cape Ferguson diuron incubations

*Mercurio et al. (2015)* collected surface seawater (24 L) in sterile containers in the tropical dry season (15 May 2012) from Cape Ferguson, QLD, Australia (latitude  $-19.2673297$ , longitude  $147.0591537$ ) (*Fig. S1*), a site that is  $\sim 17$  km downstream from the Haughton River mouth and where diuron is consistently reported in the wet season (*Lewis et al., 2009; Kennedy et al., 2012a; Kennedy et al., 2012b*). The investigators passed seawater through  $20 \mu\text{m}$  impact filters, dispensed it in 500 mL glass flasks and incubated it during 365 days on a shaking platform at  $25^\circ\text{C}$  either in the dark or in the light (12:12 light day cycle with  $40 \mu\text{mol photons m}^{-2} \text{s}^{-1}$ ), and with or without amendment of diuron (at the ecologically relevant concentration of  $8 \mu\text{g/L}$ ) (*Lewis et al., 2009; Kennedy et al., 2012a; Kennedy et al., 2012b*) (*Figs. S2A and S2B*). The investigators performed each experimental treatment in triplicate (12 flasks in total) and monitored diuron concentration for each flask over the life of the incubation. For more details of the experimental set up and diuron degradation results, see *Mercurio et al. (2015)*.

In the present study, we collected subsamples ( $705 \mu\text{L}$ ) from each flask of the Cape Ferguson incubation experiment at day 0, 2, 7, 28, 120 and 365 for 16S rRNA gene analysis. Each subsample was collected with a pipette after thoroughly shaking the flask,  $5 \mu\text{L}$  were stained with  $5 \mu\text{L}$  of DAPI ( $1 \mu\text{g/mL}$ ) in the dark for 15 min, and observed with a Nikon Ci-L epifluorescence microscope (*Fig. S2C*). This confirmed the presence of DNA-containing cells, a prerequisite for sequencing.

Total DNA was extracted by first centrifuging each sample at  $13,000 \times g$  for 30 min. Each resulting pellet was then resuspended in  $20 \mu\text{L}$  microLYSIS-Plus DNA release buffer

(Microzone, West Sussex, UK) for 30 min at room temperature and incubated using a thermal cycler following the manufacturer's tough cell lysis protocol (65 °C for 15 min; 96 °C for 2 min; 65 °C for 4 min; 96 °C for 1 min; 65 °C for 1 min; 96 °C for 30 s). Control of the absence of microLYSIS-Plus reagent contamination by foreign DNA was carried out by adding a blank sample, containing only the microLYSIS-Plus buffer (without template DNA).

Microbial amplicons were generated by PCR-amplifying the V6–V8 variable regions of the 16S rRNA gene in the total DNA using a universal primer set targeting Archaea and Bacteria (iTAG926F and iTAG1392wR primers) according to the protocol in [Dove et al. \(2013\)](#). These amplicons were paired-end sequenced on an Illumina MiSeq instrument at the Australian Centre for Ecogenomics (ACE).

### Microbiome profiling

Amplicon reads were processed using Hitman (<https://github.com/fangly/hitman>, described in [Angly et al., 2016](#)), a bioinformatic workflow based around the UPARSE methodology ([Edgar, 2013](#)). The following parameters were used: trimming length of 250 bp, minimum quality value of 7 (16 for HiFi sequences), maximum number of expected errors of 3.0 (0.5 for HiFi sequences), OTU clustering at 97% identity (species-level), GOLD ([Bernal, Ear & Kyrpides, 2001](#)) as the reference database for chimera detection, rarefaction depth of 20,000 counts, minimum global alignment of 95% identity (genus-level) for taxonomic annotation using the merged Silva ([Quast et al., 2012](#)) and Greengenes ([McDonald et al., 2012](#)) databases ([https://github.com/fangly/merge\\_gg\\_silva](https://github.com/fangly/merge_gg_silva)), gene-copy number correction with CopyRighter ([Angly et al., 2014](#)), and exclusion of taxa matching “Eukaryota\*” or “\*Chloroplast\*.”

Rarefaction curves were produced using Bio-Community's bc\_accumulate ([Angly, Fields & Tyson, 2014](#)) with 100 random subsets. After taxonomic assignments and gene-copy number correction within Hitman, calculation of  $\alpha$ -diversity was performed using Bio-Community bc\_measure\_alpha at the selected rarefaction depth. A few OTUs of interest, that could not be taxonomically assigned with Hitman, were classified by the RDP Classifier at 80% confidence ([Cole et al., 2009](#)), and Silva's SINA with 95% identity ([Quast et al., 2012](#)).

### Statistical analysis

The significance of changes in taxon relative abundance between sampling points were evaluated using LEfSe ([Segata et al., 2011](#)). The effects of incubation regimen on microbial community structure was assessed by Hellinger-transforming the microbial profiles and analyzing them using the R language (*R Foundation for Statistical Computing, Vienna, Austria, 0000*), specifically using the capscale(), adonis() and rda() functions of the vegan packages ([Dixon, 2003](#)), for PCoA, PERMANOVA and RDA analysis respectively.

### Dunk Island metagenome preparation

An additional seawater sample was collected in the tropical dry season (13 October 2009), north of Dunk Island, QLD, Australia (latitude  $-17.9242918$ , longitude  $146.1429637$ ) ([Fig. S1](#)). This site is  $\sim 15$  km downstream from the Tully River mouth, and exposed to

diuron and other PSII herbicides in comparable concentrations as the Cape Ferguson site (Lewis *et al.*, 2009; Kennedy *et al.*, 2012a; Kennedy *et al.*, 2012b). A 20 L volume was taken from a depth of 5 m and pre-filtered through a 2.7  $\mu\text{m}$  Whatman GF/D filter and a 1.6  $\mu\text{m}$  Whatman GF/A filter to remove particles and most eukaryotic microorganisms. The filtrate was then passed through a 0.22  $\mu\text{m}$  Millipore Express Plus filter to capture the bacterial and archaeal fraction. The filters were folded in half, cells inward, added to a tube containing 20 mL of lysis buffer (40 mM  $\text{Na}_2\text{EDTA}$ , 50 mM Tris pH 8.3 and 0.73 M sucrose, sterilized), stored shipboard at  $-20^\circ\text{C}$  and transferred to  $-80^\circ\text{C}$  on land.

DNA was extracted from the filter using a modified method from Suzuki *et al.* (2004). In brief, the filter was thawed on ice, added 6 mL of lysis buffer with 5 mg/mL lysozyme and the tube was incubated for 30 min at  $37^\circ\text{C}$ , while rotating at 10 rpm. Proteinase K (1.1 mg/mL final concentration) and 10% sodium dodecyl sulfate (1.1% final concentration) were added and the sample was incubated at  $55^\circ\text{C}$  for 2 h, with rotation. The lysate was split in half and DNA was extracted from each using two rounds of phenol:chloroform:isoamyl alcohol (25:24:1, pH 8.0), then one round of chloroform:isoamyl alcohol (24:1). Aqueous phases were pooled and frozen overnight at  $-20^\circ\text{C}$ . The aqueous phase was then cleaned by passage of 15 mL at a time through Amicon Ultra-15 100 kDa spin unit (EMD Millipore, Billerica, MA, USA). The filter was washed once with 8 mL of Tris EDTA buffer (TE, 10 mM, pH 8.0) and recovered with 50  $\mu\text{L}$  of TE (1 mM, pH 8.0). DNA was then further cleaned by precipitation with 70% ethanol, the pellet was washed once with 70% ethanol, air dried, and resuspended in 100  $\mu\text{L}$  TE, for a total yield of 37  $\mu\text{g}$  DNA. The resulting DNA was sequenced on an Illumina (Solexa) Genome Analyzer II instrument at the University of Arizona, producing 25.4 million pairs of 101 bp long reads.

### Read-centric metagenomic screening for phenylurea hydrolases

The Dunk Island metagenomic read pairs were cleaned by removing Illumina adapters with TRIMMOMATIC, merged using PEAR (but keeping unmerged read pairs), 5' end quality-trimmed at the first nucleotide below Q13 and filtered to remove sequences smaller than 60 bp using TRIMMOMATIC. The resulting quality-controlled reads were compared to all known PuhAB phenylurea hydrolase proteins (GI 218764925, 598062302 and 218764905), belonging to the metal-dependent amidohydrolase superfamily (Turnbull *et al.*, 2001; Khurana *et al.*, 2009), using BLASTX (Camacho *et al.*, 2009). The BLAST database also included 55 other closely-related proteins, including other herbicide hydrolases, to ensure the specificity of the results: the MolA molinate hydrolase (Sugrue *et al.*, 2015) (GenBank FN985594), four LibA linuron hydrolases (Bers *et al.*, 2011; Bers *et al.*, 2013) (GenBank JN104629, JN104630, JN104631 and JN104633) and 50 proteins from the metal dependent amidohydrolase superfamily (GI 18655481, 7245484, 23200144, 23200220, 3892028, 22218649, 14719683, 13786715, 28948588, 30749918, 999767, 24987382, 27574194, 30750126, 24371617, 40787177, 15966345, 16124371, 5817646, 22972062, 21222419, 23058081, 24216335, 3912984, 1709955, 33595951, 27375360, 27378941, 22987263, 23105179, 3914514, 16763233, 27377792, 2829648, 18311855, 15612748, 17540282, 17548772, 38108196, 15791459, 15528804, 40063581 and 24371695). Significant similarities ( $E$ -value  $< 1e-6$ ) were extracted and their alignment to the most similar proteins was visually

inspected using Jalview ([Waterhouse et al., 2009](#)). The putative taxonomic affiliation of the Puh-like proteins was established by comparing the metagenomic reads to the NCBI nt database using TBLASTX.

Under the assumption that each distinct identified PuhAB protein is encoded by a different species (the three known Puh proteins are encoded on three distinct genomes), we approximated this species' relative abundance as:  $A \approx R \times G \times 10^{-4} / (P \times M \times L)\%$ , where  $R$  is the number of reads matching *puhAB*,  $M$  is the number of screened metagenomic reads (22,927,633),  $L$  is the average read length  $L$  (93.7 bp),  $P$  is the average length of the *puhAB* genes (1,376 bp),  $G$  is the average genome length in marine microbiomes (2.58 Mbp) ([Angly et al., 2009](#)), and  $S$  is the number of species in inshore GBR water column (643 OTUs in the diuron incubation experiment).

### Contig-centric metagenomic screening for photosystem genes


For this analysis, the Dunk Island metagenomic raw read pairs were cleaned with TRIMMOMATIC by removing Illumina adapters, deleting reads with uncalled bases, truncating their 5' end to a final length of 80 bp, and removing smaller reads. The data were assembled using IDBA-UD ([Peng et al., 2012](#)), and the resulting scaffolds translated into their six possible reading frames. The hmmsearch tool of HMMER3 ([Eddy, 2011](#)) was employed to look for photosystem B proteins in these translated scaffolds using the TIGR001151 PsbA hidden Markov Model profile of TIGRFAMs ([Haft, Selengut & White, 2003](#)). A maximum  $E$ -value of  $1e-50$  was used to retrieve significant matches and their alignment was visualized in Jalview. The taxonomic affiliation of the scaffolds matching PsbA was determined by best BLASTN similarity against the NCBI nr database (minimum identity of 70% over a minimum alignment length of 1,200 bp, i.e., the length of PsbA + 40 amino acids). Nesoni (<https://github.com/Victorian-Bioinformatics-Consortium/nesoni>) and SHRiMP ([Rumble et al., 2009](#)) were used to map the Illumina reads against the metagenomic scaffolds and call single nucleotide polymorphisms (SNPs).

## RESULTS & DISCUSSION

### Microbial dynamics in Cape Ferguson diuron incubation

[Mercurio et al. \(2015\)](#) collected seawater during the tropical dry season at Cape Ferguson, an inshore region of the GBR ([Figs. S1 and S2](#)), to conduct a year-long diuron incubation experiment. In the present study, the 16S rRNA amplicon sequencing of 72 samples taken at set time points from the incubation flasks generated a total of 4.96 million read pairs (NCBI accession [PRJNA276057](#)). Processing through the Hitman bioinformatic pipeline resulted in 3.83 million high-quality sequences (77.3% of the initial amount). Rarefaction at a depth of 20,000 counts per sample provided a sequencing depth-independent view of the diversity of the samples ([Fig. S3, Table S1](#)), collectively containing 4,743 distinct OTUs (97% identity level).

The taxonomic affiliation conducted by best global alignment against the Greengenes database and subsequent gene-copy number correction ([Angly et al., 2014](#)) permitted estimation of changes in the relative abundance of prokaryotic taxa over time ([Fig. S4](#)). When averaging the replicates ([Fig. 1, Fig. S5](#)), the most abundant taxa at the start of



**Figure 1** Heatmap showing the relative abundance of microbial genera over the one-year Cape Ferguson diuron incubations. The four incubation conditions are control + dark (C\_D), control + light (C\_L), diuron + dark (D\_D) and diuron + light (D\_L). The three replicates of each incubation condition were averaged and only microbial genera reaching 1% are indicated.

the incubation (day 0) included the orders Rickettsiales (19%) and Synechococcales (14% average relative abundance), from the Cyanobacteria and Proteobacteria phyla, respectively. The microbiomes were marked by a succession of various taxa over time, as seen in previous work (Fierer *et al.*, 2010). For example, Sphingomonadales increased significantly from an initial average of 0.38% relative abundance (day 0) to dominate the communities with 30% at day 2 (LEfSe;  $\alpha < 0.05$ ). Rhodobacterales-affiliated sequences increased significantly, reaching a maximum relative abundance of 36% on average a week after the start of the incubation (day 7) (LEfSe;  $\alpha < 0.05$ ), and subsequently significantly decreased until day 120 (LEfSe;  $\alpha < 0.05$ ). At the end of the incubation experiment (day 365), Oceanospirillales were very abundant in the control flasks exposed to light, while Thiotrichales dominated the samples incubated in the dark (both control and diuron-treated).

Three predominant OTUs (OTU 12, 13 and 20) characteristic of the flasks kept in the dark could not be assigned to a taxonomic group. Further identification efforts using the RDP Classifier and Silva's SINA suggest that they all belong to the Proteobacteria phylum, more precisely to the *Salinisphaera*, *Coxiella* and GR-WP33-30 taxa (Table S2). The genus *Salinisphaera* includes a recently sequenced species that is adapted to environments with fluctuating conditions (Antunes *et al.*, 2011), while the genus *Coxiella* contains a single species that is highly resistant to environmental stresses such as temperature, osmotic pressure and ultraviolet radiation (Voth & Heinzen, 2007), and representatives of the order GR-WP33-30 were detected in uranium mines (Selenska-Pobell & Radeva, 2004). The robustness of these taxa may be responsible for their success in the dark and likely oligotrophic conditions of the incubation flasks.


### Effect of diuron on microbial profiles

The diuron measurements made by Mercurio *et al.* (2015) in the incubation flasks ranged from an initial 8.77  $\mu\text{g/L}$  (dark conditions, replicate R4) down to 3.78  $\mu\text{g/L}$  (light conditions, replicate R3) after one year of incubation. Here, we included the diuron concentration of each individual flask as an input for a constrained ordination (Fig. 2), which demonstrated a significant influence of incubation time and light exposure, but not of diuron concentrations on the microbial profiles (PERMANOVA,  $p < 0.05$ ). Dissection of the differences between diuron-treated and control flasks for each individual sampling day, PCoA (Fig. S6) confirmed that diuron did not affect microbiome composition significantly (PERMANOVA,  $p < 0.05$ ).

### Resistance of photosynthetic bacteria to diuron


Some phototrophic bacteria are inhibited by diuron, while others are insensitive. For example, photosystems I and II exist in Cyanobacteria and vascular plants, and physiological experiments have demonstrated binding of diuron on the cyanobacterial photosystem II, leading to photosynthesis inhibition (Allen *et al.*, 1983; Gadkari, 1988; Brusslan & Haselkorn, 1989; Deng, Gao & Sun, 2012). Conversely, diuron does not bind to the photosynthetic reaction center of purple bacteria and they may remain unaffected (Sinning, 1992). We thus hypothesized that the majority of phototrophic prokaryotes in the incubation experiment would be affected by diuron toxicity, resulting in their rapid decline.





**Figure 2** OTU-level RDA of the microbiomes (Hellinger-based) in the Cape Ferguson diuron incubations. OTUs are indicated by a red cross and the Greengenes taxonomic affiliation of the most discriminating is shown. Circle size is proportional to incubation time (sampling day). Environmental factors are green arrows depicting light amount in the light and dark treatments, incubation time and measured diuron concentration. Asterisks denote environmental factors that are statistically significant (PERMANOVA;  $p < 0.05$ ). Samples from day 2 and 7 were omitted from this analysis because diuron concentration was not measured on these days.

Cyanobacteria and purple bacteria such as Rhodobacteraceae were prevalent in the incubation flasks, but their relative abundance did not decline between day 0 and 28 (Fig. 2, Figs. S6A–S6D), despite the presence of more than 8.45  $\mu\text{g/L}$  diuron on average during this period, a concentration that markedly inhibits the photosynthesis of diatoms and green algae (Magnusson, Heimann & Negri, 2008; Magnusson et al., 2012). This supports previous reports that Cyanobacteria are less sensitive to PSII herbicides than eukaryotic phototrophs (Lüring & Roessink, 2006). The relative insensitivity of Cyanobacteria in our dataset could be explained by pollution-induced community tolerance (PICT) following chronic exposure to herbicides, which was previously reported for biofilms in a French river (Tlili et al., 2008; Tlili et al., 2011) and for periphyton (a mixture of detritus, algae and



**Figure 3** Sequence alignment of the PsbA proteins predicted from Dunk Island metagenomic scaffolds. Residues are colored based on the Clustal X scheme. Locations previously correlated with herbicide resistance are indicated by a box. The bold sequence corresponds to PEST type 11, implicated in resistance against the irgarol PSII herbicide. The bottom panel represents the number of conserved amino acids at each position and their consensus and the column on the right the BLASTN taxonomic classification of the scaffolds.

microorganisms growing on submerged surfaces) in the GBR lagoon (*Magnusson et al., 2012*) and in a Swedish fjord (*Molander & Blanck, 1992*). The mechanism underpinning this tolerance for diuron is not yet elucidated, but may be related to the evolution and enrichment of high-turnover variants of the PsbA protein upon which diuron and other PSII herbicides such as irgarol 1051 bind (*Eriksson et al., 2009; Deng, Gao & Sun, 2012*).

Metagenomic analysis was undertaken to explore the presence of PsbA variants and the potential for diuron resistance in the GBR lagoon. Since the samples collected during the Cape Ferguson incubation experiment contained too little biomass for comprehensive metagenomic sequencing, we prepared a metagenome from a sample collected during the tropical dry season at Dunk Island, another inshore GBR location (NCBI accession [SRR1819825](#)). Weather, river effluent and diuron exposure data indicate that the Dunk Island and Cape Ferguson samples were both representative of the GBR lagoon during the dry season, when the effects of riverine floodwaters are minimal ([Text S1, Tables S3–S6](#)), and therefore comparable (*Angly et al., 2016*). The Dunk Island metagenome was assembled into ~74,000 scaffolds (771 bp average length, 879 bp N50) from which all putative PsbA protein sequences were identified ([Fig. 3](#)). The introduction of Val219 and Ser264 mutations in PsbA confers PSII herbicide resistance (*Bettini et al., 1987; Mengistu et al., 2000*) and mutations in the PsbA PEST domain (rich in amino-acids P, E, S and T) were previously correlated with resistance in the environment (*Eriksson et al., 2009*). But none of these mutations were detected in the Dunk Island scaffolds of Cyanobacteria (*Synechococcus* and *Prochlorococcus*). Further, a total of ~24,900 reads mapped onto the 12.9 kb long *Prochlorococcus* scaffold, but no SNPs could be identified within the *psbA* gene.

PEST sequence type 11 (RETTENESANAGYK), representing a PEST type hypothesized to confer irgarol tolerance to the Swedish fjord microbiomes (Eriksson *et al.*, 2009), was detected in the metagenomic scaffolds (Fig. 3). However, BLASTN analyses suggest that this sequence was part of a eukaryotic genome (100% query coverage and 100% identity to *Micromonas*, a member of the Prasinophyceae class) (Worden *et al.*, 2009). The Prasinophyceae are composed of unicellular photosynthetic green algae and *Micromonas* (<2  $\mu\text{m}$ ) could have passed through the 1.6  $\mu\text{m}$  wide pores of the filters used during metagenome preparation. Similarly, the Swedish fjord sequencing read containing PEST type 11 (accession AM933747) best matched a eukaryotic genome from another subdivision of the Prasinophyceae (100% query coverage and 99% identity to the Pycnococcaceae family). This suggests that PEST type 11 is a general feature of the Prasinophyceae genome and, at least in the present study, not an adaptive mutation of Bacteria and Archaea to protect against PSII herbicides.

BLASTN investigation of the metagenomic scaffolds from Dunk Island encoding PsbA revealed that they were not only prokaryotic and eukaryotic. Most of them (14 out of 24) were of viral origin (Fig. 3), and 9 out of the 11 PEST types identified in the present study matched some carried by cyanophages. This large PEST type diversity and the propensity of phages to transfer genes to and from their hosts (lateral gene transfer) raises the possibility that cyanophages steer the stability of PsbA in their hosts (Lindell *et al.*, 2004; Zeidner *et al.*, 2005; Sharon *et al.*, 2007). Future research should consider how this may influence the tolerance of phototrophs to PSII herbicides.

### Potential for degradation by heterotrophic bacteria

Another hypothesis formulated in this study was that specific heterotrophic populations would carry genes for the degradation of diuron and take advantage of this resource, leading to their increase in relative abundance over time. During the one-year Cape Ferguson incubation experiment, Mercurio *et al.* (2015) reported 15–31% diuron degradation. They attributed this slow degradation in part to prokaryotic breakdown but, in the present work, we found no significant association between diuron-treated incubation flasks and heterotrophic abundance between day 28 and 365 (Fig. 2, Figs. S6D–S6F). While this evidence goes against our hypothesis of rapid heterotrophic degradation, a similar marine incubation study also detected a lack of diuron degradation over a shorter 42 d timeframe (Thomas, McHugh & Waldock, 2002).

Experiments using microbiomes from soil (Attaway, Paynter & Camper, 1982; Cullington & Walker, 1999; Widehem *et al.*, 2002; Dellamatrice & Monteiro, 2004; Ngigi *et al.*, 2011), activated sludge (Stasinakis *et al.*, 2009) and freshwater sediments (Ellis & Camper, 1982; Pesce *et al.*, 2010) have demonstrated that diuron can be degraded by bacteria belonging to the genera *Pseudomonas* (El-Deeb *et al.*, 2000), *Arthrobacter* (Turnbull *et al.*, 2001; Villaverde *et al.*, 2012), *Mycobacterium* (Khurana *et al.*, 2009), *Variovorax* (Sørensen, Albers & Aamand, 2008), *Bacillus*, *Vagococcus* and *Burkholderia* (Ngigi *et al.*, 2011). Sequences affiliated with some of these taxa, specifically *Burkholderia* and *Pseudomonas*, were detected at ~4% in our marine incubations, but their relative abundance did not change significantly

in response to long-term exposure to this herbicide (Fig. 2, Fig. S1), suggesting that exposure to diuron and any potential degradation did not alter their evolutionary fitness.

To explore the reasons for the lack of rapid heterotrophic degradation of diuron in the incubation experiment and the potential for diuron degradation in the GBR lagoon at large, we looked for phenylurea hydrolase genes, known to degrade diuron (Turnbull *et al.*, 2001; Khurana *et al.*, 2009), in the reads of the Dunk Island metagenome. Two reads had BLASTX similarities to a protein database covering the metal-dependent amidohydrolase superfamily. These reads were more similar to PuhB than to other PSII hydrolases and proteins from the same superfamily, with a high 51% amino acid identity over 54 amino acids (Fig. S7) indicative of the presence of potential phenylurea hydrolase homologs (Rost, 1999). Despite the short length of these metagenomic reads and the potential sequencing errors they contain, the proteins identified here may represent novel phenylurea or other PSII hydrolases, whose existence has previously been suggested (Pesce *et al.*, 2012). Further research will be needed to characterize the sequence, structure and function of this protein and thus confirm this hypothesis. The five top scoring similarities of one of these metagenomic reads (TBLASTX,  $\geq 98\%$  query cover,  $E$  value  $\leq 2e-22$ ) suggest that a bacterium from the Bacteroidetes phylum (Flavobacteriia or Cytophagia order) encodes this PuhB-like protein, while the five top scoring similarities for the other read (TBLASTX,  $\geq 97\%$  identity,  $E$  value  $\leq 5e-12$ ) did not agree on a precise taxonomic origin. Further, calculations (see 'Materials & Methods' section) indicate that this putative hydrolase could be present in low-abundance species, in the tail of the microbial rank-abundance curve ( $\sim 0.01\%$  relative abundance). Overall, these findings suggest that the marine microbial seed bank, "a reservoir of dormant individuals that can potentially be resuscitated" (Lennon & Jones, 2011), may have a potential for herbicide degradation.

Microbial enrichment studies that reported rapid biodegradation of diuron were conducted with a rich substrate or supplemented with alternative sources of carbon and nitrogen, sometimes under the form of soil or sediments (Widehem *et al.*, 2002; Sørensen, Albers & Aamand, 2008). From this evidence, we conclude that resources may be a limiting factor for marine heterotrophs to express their diuron-degrading potential in often oligotrophic marine waters (Schaffelke *et al.*, 2012), as is the case in marine incubations performed without supplementation (Thomas, McHugh & Waldock, 2002; Mercurio *et al.*, 2015). GBR microorganisms are thought to metabolize nutrients from land runoff at inshore sites (Alongi & McKinnon, 2005) and, given that these sites receive high diuron and nutrient input during the wet season (Packett *et al.*, 2009) perhaps along with diuron-degrading species, we predict that heterotrophic diuron degradation may be enhanced episodically in the GBR lagoon.

## CONCLUSIONS

This study used amplicon and metagenomic sequencing to evaluate the effects of a PSII herbicide on the composition of entire prokaryotic communities, rather than selected species. It provides a baseline for future research on the impacts of herbicides on the marine ecosystem by suggesting that the effects of the PSII herbicide diuron on communities of

GBR near-shore pelagic prokaryotes are limited. Metagenomic evidence suggests that prokaryotic heterotrophs in the marine water column may encode potential new herbicide hydrolase genes, though their expression may be limited by scarce environmental resources in the dry season. The apparent tolerance of marine pelagic phototrophs to diuron may have been due to the acquisition of a resistance mechanism following regular exposure to this herbicide. While no PEST sequence mutations in Cyanobacteria could explain this resistance in the present study, resident phages carried various PEST sequence types and could act as a reservoir. In summary, many components of coral reef ecosystems are stressed by herbicides from land runoff, but in contrast, the pelagic microbiome that underpins the marine food web and is integral to reef functioning, may represent an important buffer that mitigates the impacts of local anthropogenic and natural stresses on coral reefs.

## ACKNOWLEDGEMENTS

The authors would like to thank Elke Allers (the University of Arizona) for sample extraction assistance, Nicola Angel (Australian Centre for Ecogenomics) for her assistance with Illumina amplicon sequencing, and Jochen Müller (National Research Centre for Environmental Toxicology), Michael Nefedov and Nancy Lachner (Australian Centre for Ecogenomics) for assistance and advice in the lab.

## ADDITIONAL INFORMATION AND DECLARATIONS

### Funding

This study was supported by the Australian Research Council's Discovery Early Career Research Award to Florent Angly (DE120101213) and Queen Elizabeth II fellowship to Gene Tyson (DP1093175) and the Australian Government's National Environmental Research Program Tropical Ecosystems Hub Project 4.2. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

### Grant Disclosures

The following grant information was disclosed by the authors:

Australian Research Council's Discovery Early Career Research Award: DE120101213.

Queen Elizabeth II fellowship: DP1093175.

Australian Government's National Environmental Research Program Tropical Ecosystems Hub Project 4.2.

### Competing Interests

The authors declare there are no competing interests.

### Author Contributions

- Florent E. Angly performed the experiments, analyzed the data, wrote the paper, prepared figures and/or tables, reviewed drafts of the paper.
- Olga Pantos performed the experiments, wrote the paper, reviewed drafts of the paper.
- Thomas C. Morgan performed the experiments, analyzed the data, reviewed drafts of the paper.

- Virginia Rich and Philip Mercurio conceived and designed the experiments, performed the experiments, reviewed drafts of the paper.
- Hemerson Tonin performed the experiments, reviewed drafts of the paper.
- David G. Bourne, Andrew P. Negri and Gene W. Tyson conceived and designed the experiments, reviewed drafts of the paper.

### DNA Deposition

The following information was supplied regarding the deposition of DNA sequences:

[PRJNA276057](#)

[SRR1819825](#)

### Data Availability

The following information was supplied regarding data availability:

All raw data (DNA sequences and associated metadata) were submitted to GenBank.

### Supplemental Information

Supplemental information for this article can be found online at <http://dx.doi.org/10.7717/peerj.1758#supplemental-information>.

## REFERENCES

- Ainsworth TD, Thurber RV, Gates RD. 2010.** The future of coral reefs: a microbial perspective. *Trends in Ecology & Evolution* 25:233–240 DOI [10.1016/j.tree.2009.11.001](https://doi.org/10.1016/j.tree.2009.11.001).
- Allen MM, Turnburke AC, Lagace EA, Steinback KE. 1983.** Effects of photosystem II herbicides on the photosynthetic membranes of the cyanobacterium *Aphanocapsa* 6308. *Plant Physiology* 71:388–392 DOI [10.1104/pp.71.2.388](https://doi.org/10.1104/pp.71.2.388).
- Alongi DM, McKinnon AD. 2005.** The cycling and fate of terrestrially-derived sediments and nutrients in the coastal zone of the Great Barrier Reef shelf. *Marine Pollution Bulletin* 51:239–252 DOI [10.1016/j.marpolbul.2004.10.033](https://doi.org/10.1016/j.marpolbul.2004.10.033).
- Angly FE, Dennis PG, Skarszewski A, Vanwongerghem I, Hugenholtz P, Tyson GW. 2014.** CopyRighter: a rapid tool for improving the accuracy of microbial community profiles through lineage-specific gene copy number correction. *Microbiome* 2:11 DOI [10.1186/2049-2618-2-11](https://doi.org/10.1186/2049-2618-2-11).
- Angly FE, Willner D, Prieto-Davó A, Edwards RA, Schmieder R, Vega-Thurber R, Antonopoulos DA, Barott K, Cottrell MT, Desnues C, Dinsdale EA, Furlan M, Haynes M, Henn MR, Hu Y, Kirchman DL, McDole T, McPherson JD, Meyer F, Miller RM, Mundt E, Naviaux RK, Rodriguez-Mueller B, Stevens R, Wegley L, Zhang L, Zhu B, Rohwer F. 2009.** The GAAS metagenomic tool and its estimations of viral and microbial average genome size in four major biomes. *PLoS Computational Biology* 5:e1000593 DOI [10.1371/journal.pcbi.1000593](https://doi.org/10.1371/journal.pcbi.1000593).
- Angly FE, Heath C, Morgan TC, Tonin H, Rich V, Schaffelke B, Bourne DG, Tyson GW. 2016.** Marine microbial communities of the Great Barrier Reef lagoon are influenced by riverine floodwaters and seasonal weather events. *PeerJ* 4:e1511 DOI [10.7717/peerj.1511](https://doi.org/10.7717/peerj.1511).

- Angly FE, Fields CJ, Tyson GW. 2014. The Bio-Community Perl toolkit for microbial ecology. *Bioinformatics* **30**:1926–1927 DOI [10.1093/bioinformatics/btu130](https://doi.org/10.1093/bioinformatics/btu130).
- Antunes A, Alam I, Bajic VB, Stingl U. 2011. Genome sequence of *Salinisphaera shabanensis*, a gammaproteobacterium from the harsh, variable environment of the brine-seawater interface of the Shaban Deep in the Red Sea. *Journal of Bacteriology* **193**:4555–4556 DOI [10.1128/JB.05459-11](https://doi.org/10.1128/JB.05459-11).
- Attaway HH, Paynter MJ, Camper ND. 1982. Degradation of selected phenylurea herbicides by anaerobic pond sediment. *Journal of Environmental Science and Health. Part. B, Pesticides, Food Contaminants, and Agricultural Wastes* **17**:683–699 DOI [10.1080/03601238209372350](https://doi.org/10.1080/03601238209372350).
- Azam F, Fenchel T, Field JG, Gray JS, Meyer-Reil LA, Thingstad F. 1983. The ecological role of water-column microbes in the sea. *Marine Ecology Progress Series* **10**:257–263 DOI [10.3354/meps010257](https://doi.org/10.3354/meps010257).
- Balakrishnan S, Takeda K, Sakugawa H. 2012. Occurrence of Diuron and Irgarol in seawater, sediments and planktons of Seto Inland Sea, Japan. *Geochemical Journal* **46**:169–177 DOI [10.2343/geochemj.1.0163](https://doi.org/10.2343/geochemj.1.0163).
- Bellas J, Beiras R, Mariño-Balsa JC, Fernández N. 2005. Toxicity of organic compounds to marine invertebrate embryos and larvae: a comparison between the sea urchin embryogenesis bioassay and alternative test species. *Ecotoxicology* **14**:337–353 DOI [10.1007/s10646-004-6370-y](https://doi.org/10.1007/s10646-004-6370-y).
- Bernal A, Ear U, Kyrpides N. 2001. Genomes OnLine Database (GOLD): a monitor of genome projects world-wide. *Nucleic Acids Research* **29**:126–127 DOI [10.1093/nar/29.1.126](https://doi.org/10.1093/nar/29.1.126).
- Bers K, Leroy B, Breugelmans P, Albers P, Lavigne R, Sørensen SR, Aamand J, Mot RD, Wattiez R, Springael D. 2011. A novel hydrolase identified by genomic-proteomic analysis of phenylurea herbicide mineralization by *Variovorax* sp. strain SRS16. *Applied and Environmental Microbiology* **77**:8754–8764 DOI [10.1128/AEM.06162-11](https://doi.org/10.1128/AEM.06162-11).
- Bers K, Batisson I, Proost P, Wattiez R, De Mot R, Springael D. 2013. HylA, an alternative hydrolase for initiation of catabolism of the phenylurea herbicide linuron in *Variovorax* sp. strains. *Applied and Environmental Microbiology* **79**:5258–5263 DOI [10.1128/AEM.01478-13](https://doi.org/10.1128/AEM.01478-13).
- Bettini P, McNally S, Sevignac M, Darmency H, Gasquez J, Dron M. 1987. Atrazine resistance in *Chenopodium album* low and high levels of resistance to the herbicide are related to the same chloroplast *psbA* gene mutation. *Plant Physiology* **84**:1442–1446 DOI [10.1104/pp.84.4.1442](https://doi.org/10.1104/pp.84.4.1442).
- Brodie JE, Kroon FJ, Schaffelke B, Wolanski EC, Lewis SE, Devlin MJ, Bohnet IC, Bainbridge ZT, Waterhouse J, Davis AM. 2012. Terrestrial pollutant runoff to the Great Barrier Reef: an update of issues, priorities and management responses. *Marine Pollution Bulletin* **65**:81–100 DOI [10.1016/j.marpolbul.2011.12.012](https://doi.org/10.1016/j.marpolbul.2011.12.012).
- Brusslan J, Haselkorn R. 1989. Resistance to the photosystem II herbicide diuron is dominant to sensitivity in the cyanobacterium *Synechococcus* sp. PCC7942. *The EMBO Journal* **8**:1237–1245.

- Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K, Madden TL. 2009. BLAST+: architecture and applications. *BMC Bioinformatics* 10:421 DOI 10.1186/1471-2105-10-421.
- Cole JR, Wang Q, Cardenas E, Fish J, Chai B, Farris RJ, Kulam-Syed-Mohideen AS, McGarrell DM, Marsh T, Garrity GM, Tiedje JM. 2009. The ribosomal database project: improved alignments and new tools for rRNA analysis. *Nucleic Acids Research* 37:D141–D145 DOI 10.1093/nar/gkn879.
- Cullington JE, Walker A. 1999. Rapid biodegradation of diuron and other phenylurea herbicides by a soil bacterium. *Soil Biology and Biochemistry* 31:677–686 DOI 10.1016/S0038-0717(98)00156-4.
- De'ath G, Fabricius KE, Sweatman H, Puotinen M. 2012. The 27-year decline of coral cover on the Great Barrier Reef and its causes. *Proceedings of the National Academy of Sciences of the United States of America* 109:17995–17999 DOI 10.1073/pnas.1208909109.
- El-Deeb BA, Soltan SM, Ali AM, Ali KA. 2000. Detoxication of the herbicide diuron by *Pseudomonas* sp. *Folia Microbiologica* 45:211–216 DOI 10.1007/BF02908946.
- Dellamatrice PM, Monteiro RTR. 2004. Isolation of diuron-degrading bacteria from treated soil. *Brazilian Archives of Biology and Technology* 47:999–1003 DOI 10.1590/S1516-89132004000600020.
- Deng X, Gao K, Sun J. 2012. Physiological and biochemical responses of *Synechococcus* sp. PCC7942 to Irgarol 1051 and diuron. *Aquatic Toxicology* 122–123:113–119.
- Dinsdale EA, Pantos O, Smriga S, Edwards RA, Angly F, Wegley L, Hatay M, Hall D, Brown E, Haynes M, Krause L, Sala E, Sandin SA, Thurber RV, Willis BL, Azam F, Knowlton N, Rohwer F. 2008. Microbial ecology of four coral atolls in the northern Line Islands. *PLoS ONE* 3:e1584 DOI 10.1371/journal.pone.0001584.
- Dixon P. 2003. VEGAN, a package of R functions for community ecology. *Journal of Vegetation Science* 14:927–930 DOI 10.1111/j.1654-1103.2003.tb02228.x.
- Dove SG, Kline DI, Pantos O, Angly FE, Tyson GW, Hoegh-Guldberg O. 2013. Future reef decalcification under a business-as-usual CO<sub>2</sub> emission scenario. *Proceedings of the National Academy of Sciences of the United States of America* 110:15342–15347 DOI 10.1073/pnas.1302701110.
- Eddy SR. 2011. Accelerated profile HMM searches. *PLOS Computational Biology* 7:e1002195 DOI 10.1371/journal.pcbi.1002195.
- Edgar RC. 2013. UPARSE: highly accurate OTU sequences from microbial amplicon reads. *Nature Methods* 10:996–998 DOI 10.1038/nmeth.2604.
- Ellis PA, Camper ND. 1982. Aerobic degradation of diuron by aquatic microorganisms *Journal of Environmental Science and Health, Part B* 17:277–289 DOI 10.1080/03601238209372319.
- Eriksson KM, Clarke AK, Franzen L-G, Kuylenstierna M, Martinez K, Blanck H. 2009. Community-level analysis of psbA gene sequences and Irgarol tolerance in marine periphyton. *Applied and Environmental Microbiology* 75:897–906 DOI 10.1128/AEM.01830-08.



- Fabricius KE. 2005.** Effects of terrestrial runoff on the ecology of corals and coral reefs: review and synthesis. *Marine Pollution Bulletin* **50**:125–146  
[DOI 10.1016/j.marpolbul.2004.11.028](https://doi.org/10.1016/j.marpolbul.2004.11.028).
- Fay F, Linossier I, Carteau D, Dheilily A, Silkina A, Vallée-Rééhel K. 2010.** Booster biocides and microfouling. *Biofouling* **26**:787–798 [DOI 10.1080/08927014.2010.518234](https://doi.org/10.1080/08927014.2010.518234).
- Fenchel T. 2008.** The microbial loop—25 years later. *Journal of Experimental Marine Biology and Ecology* **366**:99–103 [DOI 10.1016/j.jembe.2008.07.013](https://doi.org/10.1016/j.jembe.2008.07.013).
- Fierer N, Nemergut D, Knight R, Craine JM. 2010.** Changes through time: integrating microorganisms into the study of succession. *Research in Microbiology* **161**:635–642  
[DOI 10.1016/j.resmic.2010.06.002](https://doi.org/10.1016/j.resmic.2010.06.002).
- Flores F, Collier CJ, Mercurio P, Negri AP. 2013.** Phytotoxicity of four photosystem II herbicides to tropical seagrasses. *PLoS ONE* **8**:e75798  
[DOI 10.1371/journal.pone.0075798](https://doi.org/10.1371/journal.pone.0075798).
- Gadkari D. 1988.** Assessment of the effects of the photosynthesis-inhibiting herbicides Diuron, DCMU, metamitron and metribuzin on growth and nitrogenase activity of *Nostoc muscorum* and a new cyanobacterial isolate, strain G4. *Biology and Fertility of Soils* **6**(1):50–54 [DOI 10.1007/BF00257920](https://doi.org/10.1007/BF00257920).
- Haft DH, Selengut JD, White O. 2003.** The TIGRFAMs database of protein families. *Nucleic Acids Research* **31**:371–373 [DOI 10.1093/nar/gkg128](https://doi.org/10.1093/nar/gkg128).
- Haynes D, Ralph P, Prange J, Dennison B. 2000.** The impact of the herbicide diuron on photosynthesis in three species of tropical seagrass. *Marine Pollution Bulletin* **41**:288–293 [DOI 10.1016/S0025-326X\(00\)00127-2](https://doi.org/10.1016/S0025-326X(00)00127-2).
- Haynes D, Müller J, Carter S. 2000.** Pesticide and herbicide residues in sediments and seagrasses from the Great Barrier Reef world heritage area and Queensland coast. *Marine Pollution Bulletin* **41**:279–287 [DOI 10.1016/S0025-326X\(00\)00097-7](https://doi.org/10.1016/S0025-326X(00)00097-7).
- Hughes TP, Graham NAJ, Jackson JBC, Mumby PJ, Steneck RS. 2010.** Rising to the challenge of sustaining coral reef resilience. *Trends in Ecology & Evolution* **25**:633–642 [DOI 10.1016/j.tree.2010.07.011](https://doi.org/10.1016/j.tree.2010.07.011).
- Jones R, Müller JF, Haynes D, Schreiber U. 2003.** Effects of herbicides diuron and atrazine on corals of the Great Barrier Reef, Australia. *Marine Ecology Progress Series* **251**:153–167 [DOI 10.3354/meps251153](https://doi.org/10.3354/meps251153).
- Jones RJ. 2004.** Testing the photoinhibition model of coral bleaching using chemical inhibitors. *Marine Ecology. Progress Series* **284**:133–145 [DOI 10.3354/meps284133](https://doi.org/10.3354/meps284133).
- Kennedy K, Schroeder T, Shaw M, Haynes D, Lewis S, Bentley C, Paxman C, Carter S, Brando VE, Bartkow M, Hearn L, Müller JF. 2012a.** Long term monitoring of photosystem II herbicides—correlation with remotely sensed freshwater extent to monitor changes in the quality of water entering the Great Barrier Reef, Australia. *Marine Pollution Bulletin* **65**:292–305 [DOI 10.1016/j.marpolbul.2011.10.029](https://doi.org/10.1016/j.marpolbul.2011.10.029).
- Kennedy K, Devlin M, Bentley C, Lee-Chue K, Paxman C, Carter S, Lewis SE, Brodie J, Guy E, Vardy S, Martin KC, Jones A, Packett R, Müller JF. 2012b.** The influence of a season of extreme wet weather events on exposure of the World Heritage Area Great Barrier Reef to pesticides. *Marine Pollution Bulletin* **64**:1495–1507  
[DOI 10.1016/j.marpolbul.2012.05.014](https://doi.org/10.1016/j.marpolbul.2012.05.014).

- Khurana JL, Jackson CJ, Scott C, Pandey G, Horne I, Russell RJ, Herlt A, Easton CJ, Oakeshott JG. 2009.** Characterization of the phenylurea hydrolases A and B: founding members of a novel amidohydrolase subgroup. *Biochemical Journal* **418**:431–441 DOI [10.1042/BJ20081488](https://doi.org/10.1042/BJ20081488).
- Legrand H, Herlory O, Guarini J-M, Blanchard G, Richard P. 2006.** Inhibition of microphytobenthic photosynthesis by the herbicides atrazine and diuron. *Cahiers de Biologie Marine* **47**:39–45.
- Lema KA, Willis BL, Bourne DG. 2012.** Corals form characteristic associations with symbiotic nitrogen-fixing Bacteria. *Applied and Environmental Microbiology* **78**:3136–3144 DOI [10.1128/AEM.07800-11](https://doi.org/10.1128/AEM.07800-11).
- Lennon JT, Jones SE. 2011.** Microbial seed banks: the ecological and evolutionary implications of dormancy. *Nature Reviews Microbiology* **9**:119–130 DOI [10.1038/nrmicro2504](https://doi.org/10.1038/nrmicro2504).
- Lesser MP, Mazel CH, Gorbunov MY, Falkowski PG. 2004.** Discovery of symbiotic nitrogen-fixing Cyanobacteria in corals. *Science* **305**:997–1000 DOI [10.1126/science.1099128](https://doi.org/10.1126/science.1099128).
- Lewis SE, Brodie JE, Bainbridge ZT, Rohde KW, Davis AM, Masters BL, Maughan M, Devlin MJ, Müller JF, Schaffelke B. 2009.** Herbicides: a new threat to the Great Barrier Reef. *Environmental Pollution* **157**:2470–2484 DOI [10.1016/j.envpol.2009.03.006](https://doi.org/10.1016/j.envpol.2009.03.006).
- Lindell D, Sullivan MB, Johnson ZI, Tolonen AC, Rohwer F, Chisholm SW. 2004.** Transfer of photosynthesis genes to and from Prochlorococcus viruses. *Proceedings of the National Academy of Sciences of the United States of America* **101**:11013–11018 DOI [10.1073/pnas.0401526101](https://doi.org/10.1073/pnas.0401526101).
- Lüring M, Roessink I. 2006.** On the way to cyanobacterial blooms: impact of the herbicide metribuzin on the competition between a green alga (*Scenedesmus*) and a cyanobacterium (*Microcystis*). *Chemosphere* **65**:618–626 DOI [10.1016/j.chemosphere.2006.01.073](https://doi.org/10.1016/j.chemosphere.2006.01.073).
- Magnusson M, Heimann K, Ridd M, Negri AP. 2012.** Chronic herbicide exposures affect the sensitivity and community structure of tropical benthic microalgae. *Marine Pollution Bulletin* **65**:363–372 DOI [10.1016/j.marpolbul.2011.09.029](https://doi.org/10.1016/j.marpolbul.2011.09.029).
- Magnusson M, Heimann K, Negri AP. 2008.** Comparative effects of herbicides on photosynthesis and growth of tropical estuarine microalgae. *Marine Pollution Bulletin* **56**:1545–1552 DOI [10.1016/j.marpolbul.2008.05.023](https://doi.org/10.1016/j.marpolbul.2008.05.023).
- Mai H, Morin B, Pardon P, Gonzalez P, Budzinski H, Cachot J. 2013.** Environmental concentrations of irgarol, diuron and S-metolachlor induce deleterious effects on gametes and embryos of the Pacific oyster, *Crassostrea gigas*. *Marine Environmental Research* **89**:1–8 DOI [10.1016/j.marenvres.2013.04.003](https://doi.org/10.1016/j.marenvres.2013.04.003).
- McDonald D, Price MN, Goodrich J, Nawrocki EP, DeSantis TZ, Probst A, Andersen GL, Knight R, Hugenholtz P. 2012.** An improved Greengenes taxonomy with explicit ranks for ecological and evolutionary analyses of bacteria and archaea. *The ISME Journal* **6**:610–618 DOI [10.1038/ismej.2011.139](https://doi.org/10.1038/ismej.2011.139).

- McMahon K, Bengtson Nash S, Eaglesham G, Müller JF, Duke NC, Winderlich S. 2005.** Herbicide contamination and the potential impact to seagrass meadows in Hervey Bay, Queensland, Australia. *Marine Pollution Bulletin* **51**:325–334 DOI [10.1016/j.marpolbul.2004.10.045](https://doi.org/10.1016/j.marpolbul.2004.10.045).
- Mengistu LW, Mueller-Warrant GW, Liston A, Barker rE. 2000.** psbA Mutation (valine219 to isoleucine) in *Poa annua* resistant to metribuzin and diuron. *Pest Management Science* **56**:209–217 DOI [10.1002/\(SICI\)1526-4998\(200003\)56:3<209::AID-PS117>3.0.CO;2-8](https://doi.org/10.1002/(SICI)1526-4998(200003)56:3<209::AID-PS117>3.0.CO;2-8).
- Mercurio P, Müller JF, Eaglesham G, Flores F, Negri AP. 2015.** Herbicide persistence in seawater simulation experiments. *PLoS ONE* **10**:e0136391 DOI [10.1371/journal.pone.0136391](https://doi.org/10.1371/journal.pone.0136391).
- Metz JG, Pakrasi HB, Seibert M, Arntzer CJ. 1986.** Evidence for a dual function of the herbicide-binding D1 protein in photosystem II. *FEBS Letters* **205**:269–274 DOI [10.1016/0014-5793\(86\)80911-5](https://doi.org/10.1016/0014-5793(86)80911-5).
- Mhadhbi L, Beiras R. 2012.** Acute toxicity of seven selected pesticides (alachlor, atrazine, dieldrin, diuron, pirimiphos-methyl, chlorpyrifos, diazinon) to the marine fish (Turbot, *Psetta maxima*). *Water, Air, & Soil Pollution* **223**:5917–5930 DOI [10.1007/s11270-012-1328-9](https://doi.org/10.1007/s11270-012-1328-9).
- Mitchell C, Brodie J, White I. 2005.** Sediments, nutrients and pesticide residues in event flow conditions in streams of the Mackay Whitsunday Region, Australia. *Marine Pollution Bulletin* **51**:23–36 DOI [10.1016/j.marpolbul.2004.10.036](https://doi.org/10.1016/j.marpolbul.2004.10.036).
- Molander S, Blanck H. 1992.** Detection of pollution-induced community tolerance (PICT) in marine periphyton communities established under diuron exposure. *Aquatic Toxicology* **22**:129–143 DOI [10.1016/0166-445X\(92\)90028-L](https://doi.org/10.1016/0166-445X(92)90028-L).
- Moncada A. 2004.** Environmental fate of diuron. In: *Department of pesticide regulation report*. Sacramento: Environmental Monitoring Branch, Department of Pesticide Regulation Report.
- Ngigi A, Getenga Z, Boga H, Ndalut P. 2011.** Biodegradation of phenylurea herbicide diuron by microorganisms from long-term-treated sugarcane-cultivated soils in Kenya. *Toxicological & Environmental Chemistry* **93**:1623–1635 DOI [10.1080/02772248.2011.595718](https://doi.org/10.1080/02772248.2011.595718).
- Okamura H. 2002.** Photodegradation of the antifouling compounds Irgarol 1051 and Diuron released from a commercial antifouling paint. *Chemosphere* **48**:43–50 DOI [10.1016/S0045-6535\(02\)00025-5](https://doi.org/10.1016/S0045-6535(02)00025-5).
- Packett R, Dougall C, Rohde K, Noble R. 2009.** Agricultural lands are hot-spots for annual runoff polluting the southern Great Barrier Reef lagoon. *Marine Pollution Bulletin* **58**:976–986 DOI [10.1016/j.marpolbul.2009.02.017](https://doi.org/10.1016/j.marpolbul.2009.02.017).
- Pandolfi JM, Bradbury RH, Sala E, Hughes TP, Bjorndal KA, Cooke RG, McArdle D, McClenachan L, Newman MJH, Paredes G, Warner RR, Jackson JBC. 2003.** Global trajectories of the long-term decline of coral reef ecosystems. *Science* **301**:955–958 DOI [10.1126/science.1085706](https://doi.org/10.1126/science.1085706).

- Peng Y, Leung HCM, Yiu SM, Chin FYL. 2012. IDBA-UD: a de novo assembler for single-cell and metagenomic sequencing data with highly uneven depth. *Bioinformatics* 28:1420–1428 DOI 10.1093/bioinformatics/bts174.
- Pesce S, Martin-Laurent F, Rouard N, Robin A, Montuelle B. 2010. Evidence for adaptation of riverine sediment microbial communities to diuron mineralization: incidence of runoff and soil erosion. *Journal of Soils and Sediments* 10:698–707 DOI 10.1007/s11368-010-0202-x.
- Pesce S, Beguet J, Rouard N, Devers-Lamrani M, Martin-Laurent F. 2012. Response of a diuron-degrading community to diuron exposure assessed by real-time quantitative PCR monitoring of phenylurea hydrolase A and B encoding genes. *Applied Microbiology and Biotechnology* 97:1661–1668 DOI 10.1007/s00253-012-4318-3.
- Proia L, Morin S, Peipoch M, Romani AM, Sabater S. 2011. Resistance and recovery of river biofilms receiving short pulses of Triclosan and Diuron. *Science of The Total Environment* 409:3129–3137 DOI 10.1016/j.scitotenv.2011.05.013.
- Quast C, Pruesse E, Yilmaz P, Gerken J, Schweer T, Yarza P, Peplies J, Glockner FO. 2012. The SILVA ribosomal RNA gene database project: improved data processing and web-based tools. *Nucleic Acids Research* 41:D590–D596 DOI 10.1093/nar/gks1219.
- R Foundation for Statistical Computing, Vienna, Austria. *R: a language and environment for statistical computing*.
- Ricart M, Barceló D, Geislinger A, Guasch H, Alda ML de, Romani AM, Vidal G, Villagrasa M, Sabater S. 2009. Effects of low concentrations of the phenylurea herbicide diuron on biofilm algae and bacteria. *Chemosphere* 76:1392–1401 DOI 10.1016/j.chemosphere.2009.06.017.
- Rohwer F, Seguritan V, Azam F, Knowlton N. 2002. Diversity and distribution of coral-associated bacteria. *Marine Ecology. Progress Series* 243:1–10 DOI 10.3354/meps243001.
- Rost B. 1999. Twilight zone of protein sequence alignments. *Protein Engineering* 12:85–94 DOI 10.1093/protein/12.2.85.
- Rumble SM, Lacroute P, Dalca AV, Fiume M, Sidow A, Brudno M. 2009. SHRiMP: accurate mapping of short color-space reads. *PLOS Computational Biology* 5:e1000386 DOI 10.1371/journal.pcbi.1000386.
- Safi J, Awad Y, El-Nahhal Y. 2014. Bioremediation of diuron in soil environment: influence of cyanobacterial mat. *American Journal of Plant Sciences* 05:1081–1089 DOI 10.4236/ajps.2014.58120.
- Sandin SA, Smith JE, DeMartini EE, Dinsdale EA, Donner SD, Friedlander AM, Konotchick T, Malay M, Maragos JE, Obura D, Pantos O, Paulay G, Richie M, Rohwer F, Schroeder RE, Walsh S, Jackson JBC, Knowlton N, Sala E, Ahmed N. 2008. Baselines and degradation of coral reefs in the Northern Line Islands. *PLoS ONE* 3:e1548 DOI 10.1371/journal.pone.0001548.
- Schaffelke B, Carleton J, Skuza M, Zagorskis I, Furnas MJ. 2012. Water quality in the in-shore Great Barrier Reef lagoon: implications for long-term monitoring and management. *Marine Pollution Bulletin* 65:249–260 DOI 10.1016/j.marpolbul.2011.10.031.

- Segata N, Izard J, Waldron L, Gevers D, Miropolsky L, Garrett WS, Huttenhower C. 2011.** Metagenomic biomarker discovery and explanation. *Genome Biology* 12:R60 DOI [10.1186/gb-2011-12-6-r60](https://doi.org/10.1186/gb-2011-12-6-r60).
- Selenska-Pobell S, Radeva G. 2004.** Bacterial diversity in water samples from two Uranium mill tailings as revealed by 16S rDNA retrieval. *Comptes Rendus de l'Academie Bulgare des Sciences (1991)* 57:85–90.
- Sharon I, Tzahor S, Williamson S, Shmoish M, Man-Aharonovich D, Rusch DB, Yooseph S, Zeidner G, Golden SS, Mackey SR, Adir N, Weingart U, Horn D, Venter JC, Mandel-Gutfreund Y, Béjà O. 2007.** Viral photosynthetic reaction center genes and transcripts in the marine environment. *The ISME Journal* 1:492–501 DOI [10.1038/ismej.2007.67](https://doi.org/10.1038/ismej.2007.67).
- Shaw M, Furnas MJ, Fabricius K, Haynes D, Carter S, Eaglesham G, Müller JF. 2010.** Monitoring pesticides in the Great Barrier Reef. *Marine Pollution Bulletin* 60:113–122 DOI [10.1016/j.marpolbul.2009.08.026](https://doi.org/10.1016/j.marpolbul.2009.08.026).
- Shaw CM, Brodie J, Müller JF. 2012.** Phytotoxicity induced in isolated zooxanthellae by herbicides extracted from Great Barrier Reef flood waters. *Marine Pollution Bulletin* 65:355–362 DOI [10.1016/j.marpolbul.2012.01.037](https://doi.org/10.1016/j.marpolbul.2012.01.037).
- Shaw M, Müller JF. 2005.** Preliminary evaluation of the occurrence of herbicides and PAHs in the wet tropics region of the Great Barrier Reef, Australia, using passive samplers. *Marine Pollution Bulletin* 51:876–881 DOI [10.1016/j.marpolbul.2005.04.015](https://doi.org/10.1016/j.marpolbul.2005.04.015).
- Sinning I. 1992.** Herbicide binding in the bacterial photosynthetic reaction center. *Trends in Biochemical Sciences* 17:150–154 DOI [10.1016/0968-0004\(92\)90324-3](https://doi.org/10.1016/0968-0004(92)90324-3).
- Smith R, Middlebrook R, Turner R, Huggins R, Vardy S, Warne M. 2012.** Large-scale pesticide monitoring across Great Barrier Reef catchments—paddock to reef integrated monitoring, modelling and reporting program. *Marine Pollution Bulletin* 65:117–127 DOI [10.1016/j.marpolbul.2011.08.010](https://doi.org/10.1016/j.marpolbul.2011.08.010).
- Sørensen SR, Albers CN, Aamand J. 2008.** Rapid mineralization of the phenylurea herbicide diuron by *Variovorax* sp. strain SRS16 in pure culture and within a two-member consortium. *Applied and Environmental Microbiology* 74:2332–2340 DOI [10.1128/AEM.02687-07](https://doi.org/10.1128/AEM.02687-07).
- Stasinakis AS, Kotsifa S, Gatidou G, Mamais D. 2009.** Diuron biodegradation in activated sludge batch reactors under aerobic and anoxic conditions. *Water Research* 43:1471–1479 DOI [10.1016/j.watres.2008.12.040](https://doi.org/10.1016/j.watres.2008.12.040).
- Sugrue E, Fraser NJ, Hopkins DH, Carr PD, Khurana JL, Oakeshott JG, Scott C, Jackson CJ. 2015.** Evolutionary expansion of the amidohydrolase superfamily in bacteria in response to the synthetic compounds molinate and diuron. *Applied and Environmental Microbiology* 81(7):2612–2624 DOI [10.1128/AEM.04016-14](https://doi.org/10.1128/AEM.04016-14).
- Sumpono, Perotti P, Belan A, Forestier C, Lavedrine B, Bohatier J. 2003.** Effect of diuron on aquatic bacteria in laboratory-scale wastewater treatment ponds with special reference to *Aeromonas* species studied by colony hybridization. *Chemosphere* 50:445–455 DOI [10.1016/S0045-6535\(02\)00621-5](https://doi.org/10.1016/S0045-6535(02)00621-5).

- Suzuki M, Preston C, Béjà O, De la Torre J, Steward G, DeLong E. 2004.** Phylogenetic screening of ribosomal RNA gene-containing clones in bacterial artificial chromosome (BAC) libraries from different depths in Monterey Bay. *Microbial Ecology* 48:473–488 DOI 10.1007/s00248-004-0213-5.
- Thomas KV, McHugh M, Waldo M. 2002.** Antifouling paint booster biocides in UK coastal waters: inputs, occurrence and environmental fate. *Science of The Total Environment* 293:117–127 DOI 10.1016/S0048-9697(01)01153-6.
- Thurber RV, Willner-Hall D, Rodriguez-Mueller B, Desnues C, Edwards RA, Angly F, Dinsdale E, Kelly L, Rohwer F. 2009.** Metagenomic analysis of stressed coral holobionts. *Environmental Microbiology* 11:2148–2163 DOI 10.1111/j.1462-2920.2009.01935.x.
- Tlili A, Dorigo U, Montuelle B, Margoum C, Carluer N, Gouy V, Bouchez A, Bérard A. 2008.** Responses of chronically contaminated biofilms to short pulses of diuron: an experimental study simulating flooding events in a small river. *Aquatic Toxicology* 87:252–263 DOI 10.1016/j.aquatox.2008.02.004.
- Tlili A, Montuelle B, Bérard A, Bouchez A. 2011.** Impact of chronic and acute pesticide exposures on periphyton communities. *Science of The Total Environment* 409:2102–2113 DOI 10.1016/j.scitotenv.2011.01.056.
- Turnbull GA, Ousley M, Walker A, Shaw E, Morgan JA. 2001.** Degradation of substituted phenylurea herbicides by *Arthrobacter globiformis* strain D47 and characterization of a plasmid-associated hydrolase gene, *puhA*. *Applied and Environmental Microbiology* 67:2270–2275 DOI 10.1128/AEM.67.5.2270-2275.2001.
- Villaverde J, Posada-Baquero R, Rubio-Bellido M, Laiz L, Saiz-Jimenez C, Sanchez-Trujillo MA, Morillo E. 2012.** Enhanced mineralization of diuron using a cyclodextrin-based bioremediation technology. *Journal of Agricultural and Food Chemistry* 60:9941–9947 DOI 10.1021/jf3021909.
- Voth DE, Heinzen RA. 2007.** Lounging in a lysosome: the intracellular lifestyle of *Coxiella burnetii*. *Cellular Microbiology* 9:829–840 DOI 10.1111/j.1462-5822.2007.00901.x.
- Waterhouse AM, Procter JB, Martin DMA, Clamp M, Barton GJ. 2009.** Jalview Version 2—a multiple sequence alignment editor and analysis workbench. *Bioinformatics* 25:1189–1191 DOI 10.1093/bioinformatics/btp033.
- Webster NS, Soo R, Cobb R, Negri AP. 2011.** Elevated seawater temperature causes a microbial shift on crustose coralline algae with implications for the recruitment of coral larvae. *The ISME Journal* 5:759–770 DOI 10.1038/ismej.2010.152.
- Widehem P, Ait-Aïssa S, Tixier C, Sancelme M, Veschambre H, Truffaut N. 2002.** Isolation, characterization and diuron transformation capacities of a bacterial strain *Arthrobacter* sp. N2. *Chemosphere* 46:527–534 DOI 10.1016/S0045-6535(01)00192-8.
- Worden AZ, Lee J-H, Mock T, Rouzé P, Simmons MP, Aerts AL, Allen AE, Cuvelier ML, Derelle E, Everett MV, Foulon E, Grimwood J, Gundlach H, Henrissat B, Napoli C, McDonald SM, Parker MS, Rombauts S, Salamov A, Dassow PV, Badger JH, Coutinho PM, Demir E, Dubchak I, Gentemann C, Eikrem W, Gready JE, John U, Lanier W, Lindquist EA, Lucas S, Mayer KFX, Moreau H, Not F, Otilar R, Panaud O, Pangilinan J, Paulsen I, Piegu B, Poliakov A, Robbens S, Schmutz J, Toulza E,**

Wyss T, Zelensky A, Zhou K, Armbrust EV, Bhattacharya D, Goodenough UW, Peer YV, De Grigoriev IV. 2009. Green evolution and dynamic adaptations revealed by genomes of the marine picoeukaryotes *Micromonas*. *Science* 324:268–272  
[DOI 10.1126/science.1167222](https://doi.org/10.1126/science.1167222).

Xu H, Lu A, Yu H, Sun J, Shen M. 2013. Distribution of diuron in coastal seawater and sediments from west sea area of Zhoushan island. *Open Journal of Marine Science* 03:140–147 [DOI 10.4236/ojms.2013.33015](https://doi.org/10.4236/ojms.2013.33015).

Zeidner G, Bielawski JP, Shmoish M, Scanlan DJ, Sabehi G, Béjà O. 2005. Potential photosynthesis gene recombination between *Prochlorococcus* and *Synechococcus* via viral intermediates. *Environmental Microbiology* 7:1505–1513  
[DOI 10.1111/j.1462-2920.2005.00833.x](https://doi.org/10.1111/j.1462-2920.2005.00833.x).