

RESEARCH

Open Access

Elliptic problems with nonhomogeneous boundary condition and derivatives of nonlinear terms

Dumitru Motreanu¹ and Viorica V Motreanu^{2*}

*Correspondence:

motreanu@post.bgu.ac.il

²Department of Mathematics, Ben Gurion University of the Negev, Be'er Sheva, 84105, Israel

Full list of author information is available at the end of the article

Abstract

The paper presents existence results for nonlinear elliptic problems under a nonhomogeneous Dirichlet boundary condition. The considered elliptic equations exhibit nonlinearities containing derivatives of the solution.

MSC: 35H30; 35A16**Keywords:** quasilinear elliptic problem; nonhomogeneous Dirichlet boundary conditions; existence result

1 Introduction

The aim of the paper is two-fold: first, to study nonlinear elliptic problems under nonhomogeneous Dirichlet boundary condition; second, to incorporate in the problem statement nonlinearities exhibiting derivatives of the solution. These requirements need to develop a nonstandard approach, in particular prevent the use of variational methods.

Specifically, we study two problems on a bounded domain $\Omega \subset \mathbb{R}^N$ ($N \geq 1$) with Lipschitz boundary $\partial\Omega$. We first consider the problem

$$\begin{cases} \operatorname{div}(a(x, u)\nabla u) = \operatorname{div} b(x, u, \nabla u) + f(x, u) & \text{in } \Omega, \\ u = g & \text{on } \partial\Omega, \end{cases} \quad (1)$$

where $a : \Omega \times \mathbb{R} \rightarrow \mathcal{S}_N(\mathbb{R})$, $b : \Omega \times \mathbb{R} \times \mathbb{R}^N \rightarrow \mathbb{R}^N$, $f : \Omega \times \mathbb{R} \rightarrow \mathbb{R}$ are Carathéodory functions (that is, they are measurable in $x \in \Omega$ and continuous in the other variables), $g \in H^1(\Omega)$, and $\mathcal{S}_N(\mathbb{R})$ denotes the space of $N \times N$ sized symmetric matrices. In the following definition we make clear what we understand by solution to problem (1).

Definition 1 A (weak) solution of problem (1) is an element $u \in H^1(\Omega)$ such that $u - g \in H_0^1(\Omega)$, $a(\cdot, u)\nabla u \in L^2(\Omega)^N$, $b(\cdot, u) \in L^2(\Omega)^N$, $f(\cdot, u) \in L^2(\Omega)$, and

$$\int_{\Omega} (a(x, u)\nabla u) \cdot \nabla v \, dx = \int_{\Omega} b(x, u, \nabla u) \cdot \nabla v \, dx - \int_{\Omega} f(x, u)v \, dx \quad \text{for all } v \in H_0^1(\Omega).$$

Next we focus on nonhomogeneous Dirichlet problems where, contrary to problem (1), the dependence with respect to the gradient ∇u of the solution u is not expressed in a

divergence form, namely

$$\begin{cases} \operatorname{div}(a(x, u)\nabla u) = \operatorname{div} b_0(x, u) + \sum_{i=1}^N b_i(x, u) \frac{\partial u}{\partial x_i} + f(x, u) & \text{in } \Omega, \\ u = g & \text{on } \partial\Omega. \end{cases} \quad (2)$$

Here $a : \Omega \times \mathbb{R} \rightarrow \mathcal{S}_N(\mathbb{R}), f : \Omega \times \mathbb{R} \rightarrow \mathbb{R}$ and g are as in problem (1), while $b_0 : \Omega \times \mathbb{R} \rightarrow \mathbb{R}^N$ and $b_i : \Omega \times \mathbb{R} \rightarrow \mathbb{R}$ ($i \in \{1, \dots, N\}$) are Carathéodory functions. The meaning of solution of problem (2) is as follows.

Definition 2 A (weak) solution of problem (2) is an element $u \in H^1(\Omega)$ such that $u - g \in H_0^1(\Omega)$, $a(\cdot, u)\nabla u \in L^2(\Omega)^N$, $b_0(\cdot, u) \in L^2(\Omega)^N$, $b_i(\cdot, u) \frac{\partial u}{\partial x_i} \in L^2(\Omega)$ for all $i \in \{1, \dots, N\}$, $f(\cdot, u) \in L^2(\Omega)$, and

$$\int_{\Omega} (a(x, u)\nabla u) \cdot \nabla v \, dx = \int_{\Omega} b_0(x, u) \cdot \nabla v \, dx - \sum_{i=1}^N \int_{\Omega} b_i(x, u) \frac{\partial u}{\partial x_i} v \, dx - \int_{\Omega} f(x, u) v \, dx$$

for all $v \in H_0^1(\Omega)$.

Problems of type (1) and (2) have been investigated in settings that are different from ours (see, e.g., [1–6]). For instance, problem (1) is studied in [3] when $N = 1$ and $N = 2$ with functions $a(x, s)$ and $b(x, s, \xi) = b(x, s)$ corresponding to certain physical models, as described by Reynolds equation where $b(x, s) = h(x)\rho(s)V$ with $h(x) \in \mathbb{R}, \rho(s) \geq 0, \rho(0) = 0$, and $V \in \mathbb{R}^N$. Whereas many of the previous results on problems (1) and (2) involve technical and somewhat restrictive assumptions on the data, the purpose of the present paper is to provide an elementary resolution of problems (1) and (2) in geometrically relevant situation. As an example of such a geometrically relevant situation, we mention the assumption on the term $b(x, \cdot, \xi)$ in problem (1) to vanish at two points.

Our results are stated as Theorems 1 and 2. They are existence and location theorems on problems (1) and (2), respectively, guaranteeing solutions in the sense of Definitions 1 and 2 that fulfill an estimate $\gamma_* \leq u \leq \gamma^*$ with given constants $\gamma_* \leq \gamma^*$. This *a priori* estimate of the solution is derived through natural geometric hypotheses that can be directly checked. It is also worthwhile to remark that we cannot drop by translation the nonhomogeneous boundary conditions to become homogeneous because our hypotheses would be no longer verified. The arguments used in the proof are based on truncation techniques and Schauder’s fixed point theorem. We emphasize that, due to the type of assumptions we impose, it is essential in our approach to keep separate the two terms in divergence form appearing in the statement of (1) and (2). A careful inspection of our proofs shows that we rely on the linearity with respect to the gradient ∇u in the first divergence term and on the vanishing at suitable points in the second divergence term.

The rest of the paper is organized as follows. Section 2 is devoted to problem (1). Section 3 studies problem (2).

2 Result on problem (1)

Throughout the paper the notation $\|\cdot\|_{L^2}$ and $\|\cdot\|_{H^1}$ stands for the usual norms on $L^2(\Omega)$ (or $L^2(\Omega, \mathbb{R}^N)$) and $H^1(\Omega)$, respectively. By $|\cdot|$ we denote the Euclidean norm of \mathbb{R}^N .

Let λ_1 be the first eigenvalue of the negative Laplacian differential operator on $H_0^1(\Omega)$, which is known to be positive and characterized by

$$\lambda_1 = \inf \left\{ \frac{\|\nabla v\|_{L^2}^2}{\|v\|_{L^2}^2} : v \in H_0^1(\Omega), v \neq 0 \right\}. \quad (3)$$

We suppose the following hypotheses on the data a, b, f , and g in problem (1):

(H₁) There is a Carathéodory function $\lambda : \Omega \times \mathbb{R} \rightarrow (0, +\infty)$ such that

$$(a(x, s)\xi) \cdot \xi \geq \lambda(x, s)|\xi|^2 \quad \text{for a.a. } x \in \Omega, \text{ all } s \in \mathbb{R}, \text{ all } \xi \in \mathbb{R}^N.$$

(H₂) There are constants $\gamma_*, \gamma^* \in \mathbb{R}$ with $\gamma_* \leq g(x) \leq \gamma^*$ on $\partial\Omega$ such that

$$\begin{aligned} b(x, \gamma_*, \xi) &= b(x, \gamma^*, \xi) = 0 \quad \text{for a.a. } x \in \Omega, \text{ all } \xi \in \mathbb{R}^N, \\ f(x, \gamma_*) &= f(x, \gamma^*) = 0 \quad \text{for a.a. } x \in \Omega. \end{aligned}$$

(H₃) The functions a, f are bounded on the set $M := \{(x, s) \in \Omega \times \mathbb{R} : \gamma_* \leq s \leq \gamma^*\}$ and

$$|b(x, s, \xi)| \leq C(1 + |\xi|) \quad \text{for a.a. } x \in \Omega, \text{ all } s \in [\gamma_*, \gamma^*], \text{ all } \xi \in \mathbb{R}^N.$$

(H₄) There is a Carathéodory function $v : \Omega \times \mathbb{R} \rightarrow (0, +\infty)$ such that

$$\rho := \inf_{(x,s) \in M} (\lambda(x, s) - v(x, s)) > 0$$

and

$$\begin{aligned} (b(x, s, \xi) - b(x, s, \eta)) \cdot (\xi - \eta) &\leq v(x, s)|\xi - \eta|^2 \\ \text{for a.a. } x \in \Omega, \text{ all } s \in \mathbb{R}, \xi, \eta \in \mathbb{R}^N. \end{aligned}$$

Remark 1 The constants γ_* and γ^* are not solutions of problem (1), unless $g \equiv \gamma_*$ or $g \equiv \gamma^*$ on $\partial\Omega$. Thus, in general, problem (1) has no evident solution.

Remark 2 Due to their different structure and requirements, the two terms in (1) that are in divergence form cannot be combined.

Remark 3 The last part of hypothesis (H₄) incorporates the monotonicity condition

$$(b(x, s, \xi) - b(x, s, \eta)) \cdot (\xi - \eta) \leq 0 \quad \text{for a.a. } x \in \Omega, \text{ all } s \in \mathbb{R}, \text{ all } \xi, \eta \in \mathbb{R}^N,$$

as well as the Lipschitz condition

$$|b(x, s, \xi) - b(x, s, \eta)| \leq v(x, s)|\xi - \eta| \quad \text{for a.a. } x \in \Omega, \text{ all } s \in \mathbb{R}, \text{ all } \xi, \eta \in \mathbb{R}^N,$$

and it is more general than both of them.

The result that we set forth in this section is the following theorem ensuring existence and location of solution for problem (1).

Theorem 1 Assume that hypotheses (H₁)-(H₄) are satisfied. Then problem (1) has at least one solution $u \in H^1(\Omega)$ in the sense of Definition 1 satisfying

$$\gamma_* \leq u(x) \leq \gamma^* \quad \text{for a.a. } x \in \Omega,$$

with γ_* and γ^* given in (H₂).

Proof Consider the set

$$C = \{u \in L^2(\Omega) : \gamma_* \leq u(x) \leq \gamma^* \text{ a.e. } x \in \Omega\},$$

which is a nonempty, bounded, closed, convex subset in $L^2(\Omega)$.

Claim 1: Given $u \in C$, there is a unique solution $w_u \in H^1(\Omega)$ of the problem

$$\begin{cases} \int_{\Omega} (a(x, u) \nabla w_u) \cdot \nabla v \, dx = \int_{\Omega} b(x, u, \nabla w_u) \cdot \nabla v \, dx - \int_{\Omega} f(x, u) v \, dx & \text{for all } v \in H_0^1(\Omega), \\ w_u - g \in H_0^1(\Omega). \end{cases}$$

Note that Claim 1 is equivalent to solving uniquely the problem

$$\begin{cases} \langle A(w), v \rangle = B(v) & \text{for all } v \in H_0^1(\Omega), \\ w \in H_0^1(\Omega). \end{cases} \tag{4}$$

Here $A : H_0^1(\Omega) \rightarrow H^{-1}(\Omega)$ and $B \in H^{-1}(\Omega)$ in (4) are expressed by

$$\langle A(w), v \rangle = \int_{\Omega} (a(x, u) \nabla w - b(x, u, \nabla(w + g))) \cdot \nabla v \, dx$$

and

$$B(v) = - \int_{\Omega} (a(x, u) \nabla g) \cdot \nabla v \, dx - \int_{\Omega} f(x, u) v \, dx$$

for all $w, v \in H_0^1(\Omega)$. Notice that the operators A and B are well defined due to our hypotheses.

With the fixed element $u \in C$, let us introduce the Carathéodory map $\tilde{a} : \Omega \times \mathbb{R}^N \rightarrow \mathbb{R}^N$ by

$$\tilde{a}(x, \xi) = a(x, u(x)) \xi - b(x, u(x), \xi + \nabla g(x)) \quad \text{for a.a. } x \in \Omega, \text{ all } \xi \in \mathbb{R}^N.$$

From hypotheses (H₁), (H₃), (H₄), and because $u \in C$, it follows that \tilde{a} satisfies the properties: there is a constant $c_0 > 0$ such that

$$|\tilde{a}(x, \xi)| \leq c_0(1 + |\xi|) \quad \text{for a.a. } x \in \Omega, \text{ all } \xi \in \mathbb{R}^N \tag{5}$$

and

$$(\tilde{a}(x, \xi) - \tilde{a}(x, \eta)) \cdot (\xi - \eta) \geq \rho |\xi - \eta|^2 \quad \text{for a.a. } x \in \Omega, \text{ all } \xi, \eta \in \mathbb{R}^N. \tag{6}$$

Estimate (5) guarantees that the operator A is bounded (in the sense to be bounded on bounded sets). It is easily seen that (6) implies that A is coercive, that is,

$$\lim_{\|\nabla v\|_{L^2} \rightarrow +\infty} \frac{\langle A(v), v \rangle}{\|\nabla v\|_{L^2}} = +\infty.$$

Moreover, relations (5)-(6) ensure that the operator A is maximal monotone, so pseudomonotone (see, e.g., [7, §2.3.1]). Since A is bounded, coercive, and pseudomonotone, it is surjective (see, e.g., [7, Theorem 2.99]), whence the existence of w_u in Claim 1. The uniqueness of w_u is a direct consequence of (6) (notice that $\rho > 0$). This establishes Claim 1.

Now, taking advantage of Claim 1, we define the operator $T : C \rightarrow H^1(\Omega)$ by $T(u) = w_u$ for all $u \in C$, where w_u is the unique element corresponding to $u \in C$ as proved in Claim 1.

Claim 2: The mapping $T : C \rightarrow H^1(\Omega)$ is continuous.

Let $u \in C$ and let $\{u_n\}_{n \geq 1} \subset C$ be a sequence such that $u_n \rightarrow u$ in $L^2(\Omega)$. Denote $w_n = T(u_n)$ and $w = T(u)$. Using the definition of T and choosing $v = w_n - w \in H_0^1(\Omega)$ as a test function in Claim 1 (written with u_n and u), we have

$$\begin{aligned} & \int_{\Omega} (a(x, u_n) \nabla(w - w_n)) \cdot \nabla(w - w_n) \, dx \\ &= \int_{\Omega} ((a(x, u_n) - a(x, u)) \nabla w) \cdot \nabla(w - w_n) \, dx + \int_{\Omega} (a(x, u) \nabla w) \cdot \nabla(w - w_n) \, dx \\ & \quad - \int_{\Omega} (a(x, u_n) \nabla w_n) \cdot \nabla(w - w_n) \, dx \\ &= \int_{\Omega} ((a(x, u_n) - a(x, u)) \nabla w + b(x, u, \nabla w) - b(x, u_n, \nabla w_n)) \cdot \nabla(w - w_n) \, dx \\ & \quad - \int_{\Omega} (f(x, u) - f(x, u_n))(w - w_n) \, dx. \end{aligned}$$

Combining this formula with (H_1) , (H_3) , (H_4) , (3) and the Cauchy-Schwarz inequality, we obtain

$$\begin{aligned} & \int_{\Omega} \lambda(x, u_n) |\nabla(w - w_n)|^2 \, dx \\ & \leq \| (a(x, u_n) - a(x, u)) \nabla w + b(x, u, \nabla w) - b(x, u_n, \nabla w) \|_{L^2} \| \nabla(w - w_n) \|_{L^2} \\ & \quad + \frac{1}{\sqrt{\lambda_1}} \| f(x, u_n) - f(x, u) \|_{L^2} \| \nabla(w - w_n) \|_{L^2} + \int_{\Omega} v(x, u_n) |\nabla(w - w_n)|^2 \, dx. \end{aligned}$$

Taking into account hypothesis (H_4) leads to

$$\begin{aligned} \rho \| \nabla(w - w_n) \|_{L^2} & \leq \| (a(x, u_n) - a(x, u)) \nabla w + b(x, u, \nabla w) - b(x, u_n, \nabla w) \|_{L^2} \\ & \quad + \frac{1}{\sqrt{\lambda_1}} \| f(x, u_n) - f(x, u) \|_{L^2}. \end{aligned} \tag{7}$$

Set

$$h_n(x) = (a(x, u_n) - a(x, u)) \nabla w + b(x, u, \nabla w) - b(x, u_n, \nabla w).$$

We claim that

$$h_n \rightarrow 0 \quad \text{in } L^2(\Omega). \tag{8}$$

To this end, we show that any subsequence of $\{h_n\}_{n \geq 1}$ possesses a subsequence converging to 0 in $L^2(\Omega)$. Since $u_n \rightarrow u$ in $L^2(\Omega)$, we have that, along a relabeled subsequence, $h_n(x) \rightarrow 0$ for a.a. $x \in \Omega$. Invoking (H₃), we have that $|h_n(x)|^2 \leq c_1(|\nabla w(x)|^2 + 1)$, with some constant $c_1 > 0$. Through Lebesgue's dominated convergence theorem, we conclude that $\|h_n\|_{L^2} \rightarrow 0$ as $n \rightarrow \infty$, so (8) holds true.

Similarly, we have

$$f(x, u_n) - f(x, u) \rightarrow 0 \quad \text{in } L^2(\Omega).$$

Then, in view of (7), we infer that $\|\nabla(w_n - w)\|_{L^2} \rightarrow 0$. Since the domain Ω is bounded and $w_n - w \in H_0^1(\Omega)$, we can make use of the Poincaré inequality for $w_n - w$, which yields $\|w_n - w\|_{L^2} \rightarrow 0$, whence $w_n \rightarrow w$ in $H^1(\Omega)$. This establishes Claim 2.

With the truncation function $\tau : \mathbb{R} \rightarrow \mathbb{R}$ defined by

$$\tau(s) = \begin{cases} \gamma_* & \text{if } s < \gamma_*, \\ s & \text{if } \gamma_* \leq s \leq \gamma^*, \\ \gamma^* & \text{if } s > \gamma^*, \end{cases} \tag{9}$$

consider the operator $S : C \rightarrow C$ introduced as follows

$$S(u)(x) = \tau(T(u)(x)) \quad \text{for a.a. } x \in \Omega. \tag{10}$$

Note that S takes values in $C \cap H^1(\Omega)$.

Claim 3: The mapping $S : C \rightarrow C$ has a fixed point.

Since $T : C \rightarrow H^1(\Omega)$ is continuous by Claim 2 (thus *a fortiori* $T : C \rightarrow L^2(\Omega)$ is continuous) and τ is a bounded continuous function, we infer that $S : C \rightarrow C$ is continuous. We claim that $S : C \rightarrow C$ is a compact operator. To this end, it suffices to check that $S(C)$ is relatively compact in $L^2(\Omega)$. Because of the compact embedding of $H^1(\Omega)$ in $L^2(\Omega)$, it is sufficient to prove that $S(C)$ is bounded in $H^1(\Omega)$.

Let $u \in C$ and denote $w = T(u)$. By the definition of T and inserting therein the test function $v = w - g \in H_0^1(\Omega)$, we see that

$$\begin{aligned} & \int_{\Omega} (a(x, u) \nabla(w - g)) \cdot \nabla(w - g) \, dx \\ &= \int_{\Omega} (-a(x, u) \nabla g + b(x, u, \nabla w)) \cdot \nabla(w - g) \, dx - \int_{\Omega} f(x, u)(w - g) \, dx. \end{aligned}$$

Then, as in the proof of Claim 2, from assumptions (H₁) and (H₄) we obtain that

$$\begin{aligned} \int_{\Omega} \lambda(x, u) |\nabla(w - g)|^2 \, dx &\leq \| -a(x, u) \nabla g + b(x, u, \nabla g) \|_{L^2} \| \nabla(w - g) \|_{L^2} \\ &\quad + \frac{1}{\sqrt{\lambda_1}} \| f(x, u) \|_{L^2} \| \nabla(w - g) \|_{L^2} + \int_{\Omega} v(x, u) |\nabla(w - g)|^2 \, dx, \end{aligned}$$

whence, by (H₄),

$$\rho \| \nabla(w - g) \|_{L^2} \leq c_2 \tag{11}$$

with a constant $c_2 > 0$ independent of u .

Using (11), we derive

$$\begin{aligned} \|S(u)\|_{H^1}^2 &= \int_{\Omega} |\tau(w(x))|^2 dx + \int_{\Omega} |\nabla(\tau \circ w)(x)|^2 dx \\ &\leq \int_{\Omega} (|\gamma_*| + |\gamma^*|)^2 dx + \int_{\{\gamma_* \leq w \leq \gamma^*\}} |\nabla w|^2 dx \\ &\leq c_3(1 + \|\nabla(w - g)\|_{L^2}^2) \leq c_4 \end{aligned}$$

with constants $c_3, c_4 > 0$ independent of u . It follows that the set $S(C)$ is bounded in $H^1(\Omega)$, so according to what was said before, the map $S : C \rightarrow C$ is compact. Consequently, Schauder's fixed point theorem can be applied (see, e.g., [8, p.452]), through which it follows that S admits a fixed point in C . This shows Claim 3.

Claim 4: Let $u \in C$ be a fixed point of S . Then there holds $T(u) = u$.

The existence of a point $u \in C$ such that $S(u) = u$ is ensured by Claim 3. Fix such a point u and set $w = T(u)$. In order to deduce the desired conclusion from $S(u) = u$, it suffices to check that $\gamma_* \leq w \leq \gamma^*$ a.e. in Ω . We only verify the inequality $\gamma_* \leq w$ a.e. in Ω because the proof of the other inequality is similar. By virtue of hypothesis (H_2) , we have $w = g \geq \gamma_*$ on $\partial\Omega$ (in the sense of traces), hence $(w - \gamma_*)^- = 0$ on $\partial\Omega$ and so the function $(w - \gamma_*)^-$ belongs to $H_0^1(\Omega)$ (see, e.g., [7, p.35]). Using $v = (w - \gamma_*)^-$ as a test function in the definition of T gives

$$\begin{aligned} \int_{\Omega} (a(x, u)\nabla w) \cdot \nabla(w - \gamma_*)^- dx &= \int_{\Omega} b(x, u, \nabla w) \cdot \nabla(w - \gamma_*)^- dx \\ &\quad - \int_{\Omega} f(x, u)(w - \gamma_*)^- dx, \end{aligned}$$

which reads as

$$\begin{aligned} \int_{\{\gamma_* \geq w\}} (a(x, u)\nabla w) \cdot \nabla w dx &= \int_{\{\gamma_* \geq w\}} b(x, u, \nabla w) \cdot \nabla w dx \\ &\quad - \int_{\{\gamma_* \geq w\}} f(x, u)(w - \gamma_*) dx. \end{aligned} \tag{12}$$

By the assumption that $S(u) = u$ and from (10) we know that $u(x) = S(u)(x) = \tau(w(x))$ for a.a. $x \in \Omega$, hence $u = \gamma_*$ a.e. in $\{\gamma_* \geq w\}$. Then hypothesis (H_2) implies that $b(x, u, \nabla w) = 0$ and $f(x, u) = 0$ a.e. in $\{\gamma_* \geq w\}$. Consequently, (12), (H_1) , and (H_4) entail

$$\int_{\{\gamma_* \geq w\}} |\nabla w|^2 dx \leq 0,$$

whence $\nabla(w - \gamma_*)^- = -\nabla w = 0$ a.e. in $\{\gamma_* \geq w\}$. On the other hand, we have $\nabla(w - \gamma_*)^- = 0$ in $\{\gamma_* < w\}$. Altogether, we obtain that $\nabla(w - \gamma_*)^- = 0$ in Ω . Since $(w - \gamma_*)^- \in H_0^1(\Omega)$, we conclude that $(w - \gamma_*)^- = 0$ a.e. in Ω , thus $w \geq \gamma_*$ a.e. in Ω . This proves Claim 4.

By Claims 3 and 4, the operator T admits a fixed point $u \in C$. Then the definition of T implies that $u = T(u) \in g + H_0^1(\Omega)$, so u is a solution of problem (1). In addition, the fact that $u \in C$ guarantees that $\gamma_* \leq u \leq \gamma^*$ a.e. in Ω . The proof of Theorem 1 is complete. \square

3 Result on problem (2)

The hypotheses on the data a, b_i ($i \in \{0, 1, \dots, N\}$), f , and g in problem (2) that we suppose are as follows: (H_1) in Section 2,

(H'_2) There exist constants $\gamma_*, \gamma^* \in \mathbb{R}$ such that $\gamma_* \leq g(x) \leq \gamma^*$ on $\partial\Omega$ and

$$b_i(x, \gamma_*) = b_i(x, \gamma^*) = 0 \quad \text{for a.a. } x \in \Omega, \text{ all } i \in \{0, 1, \dots, N\},$$

$$f(x, \gamma_*) = f(x, \gamma^*) = 0 \quad \text{for a.a. } x \in \Omega.$$

(H'_3) There exist constants $m > 0$ and $\tilde{m}, m_0 \geq 0$ such that

$$|a(x, s)| \leq m, \quad |f(x, s)| \leq \tilde{m}, \quad \text{and} \quad |b_0(x, s)| \leq m_0$$

$$\text{for a.a. } x \in \Omega, \text{ all } s \in [\gamma_*, \gamma^*].$$

(H'_4) $\mu := \inf_{(x,s) \in \Omega \times [\gamma_*, \gamma^*]} \lambda(x, s) > 0$ and there exist constants $m_i \geq 0, i \in \{1, \dots, N\}$, with

$$\bar{m} := \sqrt{\sum_{i=1}^N m_i^2} < \mu \sqrt{\lambda_1}, \text{ such that}$$

$$|b_i(x, s)| \leq m_i \quad \text{for a.a. } x \in \Omega, \text{ all } s \in [\gamma_*, \gamma^*], i \in \{1, \dots, N\}.$$

Remark 4 As in the case of problem (1), we note that the constant functions $u \equiv \gamma_*$ and $u \equiv \gamma^*$ are not solutions of problem (2), unless $g \equiv \gamma_*$ or $g \equiv \gamma^*$ on $\partial\Omega$.

Now we state our result of existence and location of solutions for problem (2).

Theorem 2 Assume that (H_1) , (H'_2) , (H'_3) , and (H'_4) are satisfied. Then problem (2) has at least one solution $u \in H^1(\Omega)$ in the sense of Definition 2 satisfying

$$\gamma_* \leq u(x) \leq \gamma^* \quad \text{for a.a. } x \in \Omega,$$

with γ_* and γ^* as in (H'_2) .

Proof We follow the pattern of proof of Theorem 1. Hence, using the constants γ_* and γ^* prescribed in (H'_2) , we consider

$$C = \{u \in L^2(\Omega) : \gamma_* \leq u(x) \leq \gamma^* \text{ for a.a. } x \in \Omega\}$$

which is a nonempty, bounded, closed, convex subset of $L^2(\Omega)$. We proceed by proving four claims regarding problem (2) that correspond to those in the proof of Theorem 1 for problem (1). We provide the proof since there are some differences with respect to the proof of Theorem 1.

Claim 1: For every $u \in C$, there is a unique solution $w_u \in H^1(\Omega)$ of the problem

$$\begin{cases} \int_{\Omega} (a(x, u) \nabla w_u) \cdot \nabla v \, dx \\ = \int_{\Omega} b_0(x, u) \cdot \nabla v \, dx - \sum_{i=1}^N \int_{\Omega} b_i(x, u) \frac{\partial w_u}{\partial x_i} v \, dx - \int_{\Omega} f(x, u) v \, dx & \text{for all } v \in H_0^1(\Omega), \\ w_u - g \in H_0^1(\Omega). \end{cases}$$

As in the proof of Theorem 1, first we note that Claim 1 is equivalent to proving that the problem

$$\begin{cases} A(w, v) = B(v) & \text{for all } v \in H_0^1(\Omega), \\ w \in H_0^1(\Omega) \end{cases} \quad (13)$$

admits a unique solution, where

$$A(w, v) = \int_{\Omega} (a(x, u) \nabla w) \cdot \nabla v \, dx + \sum_{i=1}^N \int_{\Omega} b_i(x, u) \frac{\partial w}{\partial x_i} v \, dx$$

and

$$B(v) = \int_{\Omega} (b_0(x, u) - a(x, u) \nabla g) \cdot \nabla v \, dx - \sum_{i=1}^N \int_{\Omega} b_i(x, u) \frac{\partial g}{\partial x_i} v \, dx - \int_{\Omega} f(x, u) v \, dx.$$

For $u \in C$, by the Cauchy-Schwarz inequalities in $L^2(\Omega)$ and in \mathbb{R}^N , as well as (H'_4) and (3), we derive the estimate

$$\begin{aligned} \left| \sum_{i=1}^N \int_{\Omega} b_i(x, u) \frac{\partial w}{\partial x_i} v \, dx \right| &\leq \sum_{i=1}^N \int_{\Omega} \left| b_i(x, u) \frac{\partial w}{\partial x_i} \right| |v| \, dx \\ &\leq \|v\|_{L^2} \sum_{i=1}^N \left(\int_{\Omega} |b_i(x, u)|^2 \left| \frac{\partial w}{\partial x_i} \right|^2 \, dx \right)^{\frac{1}{2}} \\ &\leq \|v\|_{L^2} \sum_{i=1}^N m_i \left(\int_{\Omega} \left| \frac{\partial w}{\partial x_i} \right|^2 \, dx \right)^{\frac{1}{2}} \\ &\leq \|v\|_{L^2} \left(\sum_{i=1}^N m_i^2 \right)^{\frac{1}{2}} \left(\sum_{i=1}^N \int_{\Omega} \left| \frac{\partial w}{\partial x_i} \right|^2 \, dx \right)^{\frac{1}{2}} \\ &= \|v\|_{L^2} \bar{m} \left(\int_{\Omega} |\nabla w|^2 \, dx \right)^{\frac{1}{2}} \\ &\leq \frac{\bar{m}}{\sqrt{\lambda_1}} \|\nabla v\|_{L^2} \|\nabla w\|_{L^2} \end{aligned} \quad (14)$$

for all $w, v \in H_0^1(\Omega)$. Using the Cauchy-Schwarz inequality in $L^2(\Omega)$, the fact that $u \in C$, (H'_3) and (14), we get

$$\begin{aligned} |A(w, v)| &\leq \|a(x, u) \nabla w\|_{L^2} \|\nabla v\|_{L^2} + \left| \sum_{i=1}^N \int_{\Omega} b_i(x, u) \frac{\partial w}{\partial x_i} v \, dx \right| \\ &\leq \left(m + \frac{\bar{m}}{\sqrt{\lambda_1}} \right) \|\nabla w\|_{L^2} \|\nabla v\|_{L^2} \quad \text{for all } w, v \in H_0^1(\Omega), \end{aligned}$$

which ensures that $A : H_0^1(\Omega) \times H_0^1(\Omega) \rightarrow \mathbb{R}$ is a continuous bilinear form. From (H_1) , the fact that $u \in C$, (H'_4) and (14), we have

$$A(v, v) \geq \left(\mu - \frac{\bar{m}}{\sqrt{\lambda_1}} \right) \|\nabla v\|_{L^2}^2 \quad \text{for all } v \in H_0^1(\Omega).$$

Since $\bar{m} < \mu\sqrt{\lambda_1}$ (as postulated in (H'_4)), we infer that $A : H_0^1(\Omega) \times H_0^1(\Omega) \rightarrow \mathbb{R}$ is also coercive.

On the basis of the reasoning in (14), the following estimate holds

$$\left| \sum_{i=1}^N \int_{\Omega} b_i(x, u) \frac{\partial g}{\partial x_i} v \, dx \right| \leq \frac{\bar{m}}{\sqrt{\lambda_1}} \|\nabla g\|_{L^2} \|\nabla v\|_{L^2} \tag{15}$$

for all $v \in H_0^1(\Omega)$. Taking into account that $u \in C$, (H'_3) , (15) and (3), we see that

$$|B(v)| \leq \left(\left(m + \frac{\bar{m}}{\sqrt{\lambda_1}} \right) \|\nabla g\|_{L^2} + \left(m_0 + \frac{\tilde{m}}{\sqrt{\lambda_1}} \right) |\Omega|^{\frac{1}{2}} \right) \|\nabla v\|_{L^2} \quad \text{for all } v \in H_0^1(\Omega),$$

where $|\Omega|$ stands for the Lebesgue measure of Ω . Therefore $B : H_0^1(\Omega) \rightarrow \mathbb{R}$ is linear and continuous. The properties of the mappings A and B permit to apply the Lax-Milgram theorem, through which we conclude that problem (13) admits a unique solution. This establishes Claim 1.

As in the proof of Theorem 1, we introduce the operator $T : C \rightarrow H^1(\Omega)$ defined by $T(u) = w_u$ for all $u \in C$, with w_u given in Claim 1.

Claim 2: The mapping $T : C \rightarrow H^1(\Omega)$ is continuous.

In order to prove this assertion, we proceed as in the proof of Claim 2 in Theorem 1. Fix $u \in C$ and consider a sequence $\{u_n\}_{n \geq 1} \subset C$ such that $u_n \rightarrow u$ in $L^2(\Omega)$. Denoting $w_n = T(u_n)$ and $w = T(u)$, we find that

$$\begin{aligned} & \int_{\Omega} (a(x, u_n) \nabla(w - w_n)) \cdot \nabla(w - w_n) \, dx \\ &= \int_{\Omega} ((a(x, u_n) - a(x, u)) \nabla w + b_0(x, u) - b_0(x, u_n)) \cdot \nabla(w - w_n) \, dx \\ & \quad - \sum_{i=1}^N \int_{\Omega} \left[(b_i(x, u) - b_i(x, u_n)) \frac{\partial w}{\partial x_i} + b_i(x, u_n) \frac{\partial(w - w_n)}{\partial x_i} \right] (w - w_n) \, dx \\ & \quad - \int_{\Omega} (f(x, u) - f(x, u_n))(w - w_n) \, dx. \end{aligned} \tag{16}$$

A straightforward calculation entails

$$\left| \sum_{i=1}^N \int_{\Omega} b_i(x, u_n) \frac{\partial(w - w_n)}{\partial x_i} (w - w_n) \, dx \right| \leq \frac{\bar{m}}{\sqrt{\lambda_1}} \|\nabla(w - w_n)\|_{L^2}^2. \tag{17}$$

Combining (H_1) , (H'_3) , (H'_4) , (16), (17), (3), and the Cauchy-Schwarz inequality yields

$$\begin{aligned} \mu \|\nabla(w - w_n)\|_{L^2} &\leq \|(a(x, u_n) - a(x, u)) \nabla w + b_0(x, u) - b_0(x, u_n)\|_{L^2} \\ & \quad + \frac{1}{\sqrt{\lambda_1}} \left\| \sum_{i=1}^N (b_i(x, u) - b_i(x, u_n)) \frac{\partial w}{\partial x_i} \right\|_{L^2} \\ & \quad + \frac{1}{\sqrt{\lambda_1}} \|f(x, u) - f(x, u_n)\|_{L^2} + \frac{\bar{m}}{\sqrt{\lambda_1}} \|\nabla(w - w_n)\|_{L^2}, \end{aligned} \tag{18}$$

with μ and \bar{m} in (H'_4) .

Proceeding as in (8) we show that

$$(a(x, u_n) - a(x, u)) \nabla w + b_0(x, u) - b_0(x, u_n) \rightarrow 0 \quad \text{in } L^2(\Omega), \tag{19}$$

$$\sum_{i=1}^N (b_i(x, u) - b_i(x, u_n)) \frac{\partial w}{\partial x_i} \rightarrow 0 \quad \text{and} \quad f(x, u) - f(x, u_n) \rightarrow 0 \quad \text{in } L^2(\Omega). \tag{20}$$

Now it suffices to combine (18), (19), (20) and recall that $\bar{m} < \mu \sqrt{\lambda_1}$ (see (H'_4)) to conclude that $\|\nabla(w_n - w)\|_{L^2} \rightarrow 0$. Then, because $w_n - w \in H_0^1(\Omega)$ and Ω is bounded, by the Poincaré inequality, we also deduce that $\|w_n - w\|_{L^2} \rightarrow 0$. This amounts to saying that $w_n \rightarrow w$ in $H^1(\Omega)$, which proves Claim 2.

Following the approach developed in the proof of Theorem 1, we introduce the operator $S : C \rightarrow C$ given by (10), with the truncation function $\tau : \mathbb{R} \rightarrow \mathbb{R}$ defined in (9) corresponding to the constants γ_* and γ^* in (H'_2) .

Claim 3: The mapping $S : C \rightarrow C$ has a fixed point.

Claim 2 readily implies that the mapping $S : C \rightarrow C$ is continuous. Let us check that $S : C \rightarrow C$ is a compact operator. To see this, it suffices to check that $S(C)$ is relatively compact in $L^2(\Omega)$. Thanks to the compactness of the embedding of $H^1(\Omega)$ into $L^2(\Omega)$, this reduces to show that $S(C)$ is bounded in $H^1(\Omega)$. To this end, let $u \in C$ and denote $w = T(u)$. We can argue as in the proof of Theorem 1 by relying now on the present hypotheses. We obtain from $w = T(u)$ with the test function $v = w - g \in H_0^1(\Omega)$, in conjunction with (H_1) , (H'_3) , (H'_4) , that

$$\begin{aligned} \mu \|\nabla(w - g)\|_{L^2} &\leq \|b_0(x, u) - a(x, u) \nabla g\|_{L^2} + \frac{1}{\sqrt{\lambda_1}} (\bar{m} \|\nabla w\|_{L^2} + \tilde{m}) \\ &\leq \frac{\bar{m}}{\sqrt{\lambda_1}} \|\nabla(w - g)\|_{L^2} + c_0, \end{aligned}$$

where $c_0 > 0$ is a constant independent of u . In view of hypothesis (H'_4) , it follows that

$$\|\nabla w\|_{L^2} \leq c_1, \tag{21}$$

with a constant $c_1 > 0$ independent of u . Using (21) and the definition of S , we get the estimate

$$\begin{aligned} \|S(u)\|_{H^1}^2 &= \int_{\Omega} |\tau(w(x))|^2 dx + \int_{\Omega} |\nabla(\tau \circ w)(x)|^2 dx \\ &\leq \int_{\Omega} (|\gamma_*| + |\gamma^*|)^2 dx + \int_{\{\gamma_* \leq w \leq \gamma^*\}} |\nabla w|^2 dx \leq c_2, \end{aligned}$$

with $c_2 > 0$ independent of u . We conclude that the set $S(C)$ is bounded in $H^1(\Omega)$, so relatively compact in $L^2(\Omega)$. Therefore the map $S : C \rightarrow C$ is compact. This enables us to apply Schauder's fixed point theorem (see, e.g., [8, p.452]), which implies that S possesses a fixed point in C . Claim 3 is thus shown.

Claim 4: If $u \in C$ is a fixed point of S , then $T(u) = u$.

Let $u \in C$ be a fixed point of S and set $w = T(u)$. In order to show that u is a fixed point of T , it is needed to be fulfilled $\gamma_* \leq w \leq \gamma^*$ a.e. in Ω . The proof is done following the

pattern of the corresponding part in the proof of Theorem 1. We outline the proof of $w \geq \gamma_*$ a.e. in Ω (the proof of the other inequality is similar).

Testing in $w = T(u)$ with $v = (w - \gamma_*)^- \in H_0^1(\Omega)$ yields

$$\int_{\{\gamma_* \geq w\}} (a(x, u) \nabla w) \cdot \nabla w \, dx = \int_{\{\gamma_* \geq w\}} b_0(x, u) \cdot \nabla w \, dx - \int_{\{\gamma_* \geq w\}} \left(\sum_{i=1}^N b_i(x, u) \frac{\partial w}{\partial x_i} + f(x, u) \right) (w - \gamma_*) \, dx. \quad (22)$$

In $\{\gamma_* \geq w\}$ it is true that

$$u(x) = S(u)(x) = \tau(w(x)) = \gamma_* \quad \text{for a.e. } x \in \Omega.$$

Then hypothesis (H_2') implies that

$$b_i(x, u) = f(x, u) = 0 \quad \text{a.e. in } \{\gamma_* \geq w\}, i \in \{0, 1, \dots, N\}.$$

Combining with (22), (H_1) , and (H_4') entails

$$\int_{\{\gamma_* \geq w\}} |\nabla w|^2 \, dx \leq 0.$$

It turns out that $\nabla(w - \gamma_*)^- = -\nabla w = 0$ a.e. in $\{\gamma_* \geq w\}$. Also, it is clear that $\nabla(w - \gamma_*)^- = 0$ in $\{\gamma_* < w\}$. Consequently, the equality $\nabla(w - \gamma_*)^- = 0$ in Ω is valid, which results in $(w - \gamma_*)^- = 0$ a.e. in Ω because $(w - \gamma_*)^- \in H_0^1(\Omega)$. This reads as $w \geq \gamma_*$ a.e. in Ω , so Claim 4 is fulfilled.

Now we can conclude the proof. Claims 3 and 4 ensure that there exists a fixed point $u \in C$ of the operator T . This means that $u = T(u) \in g + H_0^1(\Omega)$ and u is a solution of problem (2). Moreover, since $u \in C$, we also have $\gamma_* \leq u \leq \gamma^*$ a.e. in Ω . The desired conclusion is achieved. \square

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

DM and WM jointly worked and obtained all the results presented in the paper and participated equally in the preparation of the paper. Both authors read and approved the final manuscript.

Author details

¹Département de Mathématiques, Université de Perpignan, Perpignan, 66860, France. ²Department of Mathematics, Ben Gurion University of the Negev, Be'er Sheva, 84105, Israel.

Acknowledgements

The second author is supported by the Marie Curie Intra-European Fellowship for Career Development within the European Community's 7th Framework Program (Grant Agreement No. PIEF-GA-2010-274519).

Received: 22 October 2013 Accepted: 6 December 2013 Published: 07 Jan 2014

References

1. Bensoussan, A, Boccardo, L, Murat, F: On a nonlinear partial differential equation having natural growth terms and unbounded solution. *Ann. Inst. Henri Poincaré, Anal. Non Linéaire* **5**, 347-364 (1988)
2. Boccardo, L, Giachetti, D, Diaz, JI, Murat, F: Existence and regularity of renormalized solutions for some elliptic problems involving derivatives of nonlinear terms. *J. Differ. Equ.* **106**, 215-237 (1993)

3. Buscaglia, G, Ciuperca, I, Jai, M: Existence and uniqueness for several non-linear elliptic problems arising in lubrication theory. *J. Differ. Equ.* **218**, 187-215 (2005)
4. Ferone, V, Murat, F: Nonlinear problems having natural growth in the gradient: an existence result when the source terms are small. *Nonlinear Anal.* **42**, 1309-1326 (2000)
5. Gilbarg, D, Trudinger, NS: *Elliptic Partial Differential Equations of Second Order*. Springer, Berlin (2001)
6. Ladyzhenskaya, OA, Ural'tseva, NN: *Linear and Quasilinear Elliptic Equations*. Academic Press, New York (1968)
7. Carl, S, Le, VK, Motreanu, D: *Nonsmooth Variational Problems and Their Inequalities. Comparison Principles and Applications*. Springer, New York (2007)
8. Zeidler, E: *Nonlinear Functional Analysis and Its Applications. I. Fixed-Point Theorems*. Springer, New York (1986)

10.1186/1687-2770-2014-6

Cite this article as: Motreanu and Motreanu: **Elliptic problems with nonhomogeneous boundary condition and derivatives of nonlinear terms.** *Boundary Value Problems* 2014, **2014**:6

Submit your manuscript to a SpringerOpen[®] journal and benefit from:

- ▶ Convenient online submission
- ▶ Rigorous peer review
- ▶ Immediate publication on acceptance
- ▶ Open access: articles freely available online
- ▶ High visibility within the field
- ▶ Retaining the copyright to your article

Submit your next manuscript at ▶ springeropen.com
