

Hindawi Publishing Corporation  
Discrete Dynamics in Nature and Society  
Volume 2011, Article ID 527574, 18 pages  
doi:10.1155/2011/527574

## Research Article

# On Solution and Stability of a Two-Variable Functional Equations

Y. J. Cho,<sup>1</sup> R. Saadati,<sup>2</sup> S. Shabanian,<sup>3</sup> and S. M. Vaezpour<sup>3</sup>

<sup>1</sup> Department of Mathematics Education and RINS, Gyeongsang National University, Chinju 660-701, Republic of Korea

<sup>2</sup> Department of Mathematics, Science and Research Branch, Islamic Azad University, Tehran 14778, Iran

<sup>3</sup> Department of Mathematics and Computer Science, Amirkabir University of Technology, 424 Hafez Avenue, Tehran 15914, Iran

Correspondence should be addressed to R. Saadati, [rsaadati@eml.cc](mailto:rsaadati@eml.cc)

Received 5 June 2011; Accepted 30 August 2011

Academic Editor: Yihong Song

Copyright © 2011 Y. J. Cho et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

The main purpose of this paper is to investigate the stability of the functional equation  $f(x + y, y + z) = 2f(x/2, y/2) + 2f(y/2, z/2)$  in normed spaces. The solutions of such functional equations are considered.

## 1. Introduction

The stability of functional equations was originated from a question of Ulam in 1940, concerning the stability of group homomorphism [1]. Since then, the stability problems of various functional equations have been extensively investigated by a number of authors and there are many investigating results concerning this problem (see [2–12]).

Throughout this paper, we study the stability of the functional equation

$$f(x + y, y + z) = 2f\left(\frac{x}{2}, \frac{y}{2}\right) + 2f\left(\frac{y}{2}, \frac{z}{2}\right), \quad (1.1)$$

where  $X$  is a normed space,  $Y$  is a Banach space, and  $f : X \times X \rightarrow Y$  is a mapping. The solutions of such functions are considered.

We recall that a mapping  $f : X \times X \rightarrow Y$  is called *additive* if

$$f(x + z, y + w) = f(x, y) + f(z, w). \quad (1.2)$$

Throughout this paper  $X$  is a normed space and  $Y$  is a Banach space.

## 2. The General Solution of the Functional Equation (1.1)

Before proceeding the proof of the main result, we shall need the following two lemmas.

**Lemma 2.1.** *Let  $f : X \times X \rightarrow Y$  be a mapping satisfying (1.1) and  $f(-x, -y) = f(x, y)$  for all  $x, y \in X$ . Then  $f$  is zero.*

*Proof.* Suppose that  $f : X \times X \rightarrow Y$  is a mapping satisfying (1.1) and  $f(-x, -y) = f(x, y)$ . It is clear that  $f(0, 0) = 0$ . Letting  $y = z = 0$  and replacing  $x$  by  $2x$  in (1.1), we obtain

$$f(2x, 0) = 2f(x, 0). \quad (2.1)$$

Similarly, letting  $x = y = 0$  and replacing  $z$  by  $2x$  in (1.1), we obtain

$$f(0, 2x) = 2f(0, x). \quad (2.2)$$

Putting  $y = 0$  and replacing  $z$  by  $y$  in (1.1), we get

$$f(x, y) = 2f\left(\frac{x}{2}, 0\right) + 2f\left(0, \frac{y}{2}\right) = f(x, 0) + f(0, y), \quad (2.3)$$

for all  $x, y \in X$ , where the last equality follows from (2.1) and (2.2). It follows from (2.3) and (1.1) that

$$\begin{aligned} f(x + y, 0) + f(0, x - y) &= f(x + y, x - y) \\ &= 2f\left(\frac{y}{2}, \frac{x}{2}\right) + 2f\left(\frac{x}{2}, \frac{-y}{2}\right) \\ &= 2f\left(\frac{y}{2}, 0\right) + 2f\left(0, \frac{x}{2}\right) + 2f\left(\frac{x}{2}, 0\right) + 2f\left(0, \frac{-y}{2}\right) \\ &= f(y, 0) + f(0, x) + f(x, 0) + f(0, -y). \end{aligned} \quad (2.4)$$

Replacing  $x$  and  $y$  by  $x/2$  and  $x/2$  in (2.4), respectively, we obtain

$$f(x, 0) = 2f\left(0, \frac{x}{2}\right) + 2f\left(\frac{x}{2}, 0\right). \quad (2.5)$$

Also, replacing  $y$  and  $x$  by  $-x/2$  and  $x/2$  in (2.4), respectively, we get

$$f(0, x) = 2f\left(\frac{x}{2}, 0\right) + 2f\left(0, \frac{x}{2}\right). \quad (2.6)$$

It follows from (2.5) and (2.6) that

$$f(x, 0) = f(0, x). \quad (2.7)$$

Consequently, by (2.4), we obtain

$$f(x + y, 0) + f(x - y, 0) = 2f(x, 0) + 2f(y, 0). \quad (2.8)$$

Replacing  $y$  by  $x$  in (2.8), we get

$$f(2x, 0) = 4f(x, 0). \quad (2.9)$$

By (2.1) and (2.9), we have

$$f(x, 0) = 0. \quad (2.10)$$

The equalities (2.7) and (2.10) imply that

$$f(0, x) = 0. \quad (2.11)$$

Now, it follows from (2.3), (2.10), and (2.11) that

$$f(x, y) = 0. \quad (2.12)$$

This completes the proof.  $\square$

**Lemma 2.2.** *Let  $f : X \times X \rightarrow Y$  be a mapping satisfying (1.1) and  $f(-x, -y) = -f(x, y)$  for all  $x, y \in X$ . Then  $f$  is additive.*

*Proof.* It is easy to show that  $f(0, 0) = 0$ . Putting  $y = z = 0$  and replacing  $x$  by  $2x$  in (1.1), we obtain

$$f(2x, 0) = 2f(x, 0). \quad (2.13)$$

Again, putting  $x = y = 0$  and replacing  $z$  by  $2x$  in (1.1), we obtain

$$f(0, 2x) = 2f(0, x). \quad (2.14)$$

Putting  $y = 0$  and replacing  $z$  by  $y$  in (1.1), we get

$$f(x, y) = 2f\left(\frac{x}{2}, 0\right) + 2f\left(0, \frac{y}{2}\right) = f(x, 0) + f(0, y), \quad (2.15)$$

for all  $x, y \in X$ , where the last equality follows from (2.13) and (2.14). It follows from (2.13), (2.14), and (2.15) that

$$\begin{aligned} f(x+y, 0) - f(0, x-y) &= f(x+y, 0) + f(0, y-x) \\ &= f(x+y, y-x) \\ &= f(x, y) + f(y, -x). \end{aligned} \tag{2.16}$$

Replacing  $y$  by  $-y$  in (2.16), we obtain

$$f(x-y, 0) - f(0, x+y) = f(x, -y) + f(-y, -x). \tag{2.17}$$

Also, we have

$$\begin{aligned} f(x+y, 0) + f(0, x-y) &= f(x+y, x-y) \\ &= f(x, -y) + f(y, x). \end{aligned} \tag{2.18}$$

Replacing  $y$  by  $-y$  in (2.18), we get

$$f(x-y, 0) + f(0, x+y) = f(x, y) + f(-y, x). \tag{2.19}$$

Now, by (2.16), (2.17), (2.18), and (2.19), we have

$$f(x+y, 0) - f(x-y, 0) = 2f(y, 0). \tag{2.20}$$

Replacing  $x$  with  $x+y$  in the above equality, we get

$$f(x+2y, 0) - f(x, 0) = 2f(y, 0). \tag{2.21}$$

Now, replacing  $y$  by  $y/2$  in the previous equality, we obtain

$$f(x+y, 0) = f(x, 0) + f(y, 0). \tag{2.22}$$

Similarly, one can prove that

$$f(0, z+w) = f(0, z) + f(0, w). \tag{2.23}$$

By (2.22) and (2.23), we conclude that

$$f(x+y, z+w) = f(x, z) + f(y, w), \tag{2.24}$$

which shows that  $f$  is additive. This completes the proof.  $\square$

Now, we are ready to present the general solution of (1.1).

**Theorem 2.3.** *Every mapping  $f : X \times X \rightarrow Y$  satisfying (1.1) is additive.*

*Proof.* One can write

$$f(x, y) = f^e(x, y) + f^o(x, y), \quad (2.25)$$

where  $f^e(x, y) := (f(x, y) + f(-x, -y))/2$  and  $f^o := (f(x, y) - f(-x, -y))/2$ . Since  $f$  satisfies (1.1),  $f^e$  and  $f^o$  satisfy (1.1). Further, we have  $f^e(-x, -y) = f^e(x, y)$  and  $f^o(-x, -y) = -f^o(x, y)$  for all  $x, y \in X$ . By Lemmas 2.1 and 2.2,  $f^e(x, y)$  is zero and  $f^o(x, y)$  is additive, respectively. It follows that  $f$  is additive. This completes the proof.  $\square$

### 3. The Stability of the Functional Equation (1.1)

Throughout this section, we prove the stability of the functional equation (1.1).

**Theorem 3.1.** *Let  $f : X \times X \rightarrow Y$  be a mapping such that*

$$\left\| f(x + y, y + z) - 2f\left(\frac{x}{2}, \frac{y}{2}\right) - 2f\left(\frac{y}{2}, \frac{z}{2}\right) \right\| \leq \phi(x, y, z), \quad (3.1)$$

where  $\phi : X \times X \times X \rightarrow [0, \infty)$  satisfies  $\sum_{i=0}^{\infty} 2^i \phi(x/2^i, y/2^i, z/2^i) < \infty$  for all  $x, y, z \in X$ . Then there exists a mapping  $F : X \times X \rightarrow Y$  satisfying (1.1),

$$\begin{aligned} \|f(x, y) - F(x, y)\| &\leq \sum_{i=1}^{\infty} 2^i \left( \phi\left(\frac{x}{2^i}, 0, 0\right) + \phi\left(0, 0, \frac{y}{2^i}\right) \right) + \phi(x, 0, y), \\ \lim_{n \rightarrow \infty} 2^n f\left(\frac{x}{2^n}, \frac{y}{2^n}\right) &= F(x, y), \end{aligned} \quad (3.2)$$

for all  $x, y \in X$ .

*Proof.* Letting  $x = y = z = 0$ , we have  $f(0, 0) = 0$ . Putting  $y = z = 0$ , we obtain

$$\left\| f(x, 0) - 2f\left(\frac{x}{2}, 0\right) \right\| \leq \phi(x, 0, 0). \quad (3.3)$$

By induction, we conclude that

$$\left\| f(x, 0) - 2^n f\left(\frac{x}{2^n}, 0\right) \right\| \leq \sum_{i=0}^{n-1} 2^i \phi\left(\frac{x}{2^i}, 0, 0\right). \quad (3.4)$$

Put  $K_{m,n} = \|2^m f(x/2^m, 0) - 2^n f(x/2^n, 0)\|$ . Then

$$\begin{aligned} K_{m,n} &\leq K_{m,n+1} + K_{n+1,n} \\ &= K_{m,n+1} + 2^n \left\| 2f\left(\frac{x}{2^{n+1}}, 0\right) - f\left(\frac{x}{2^n}, 0\right) \right\| \\ &\leq K_{m,n+1} + 2^n \phi\left(\frac{x}{2^n}, 0, 0\right), \end{aligned} \quad (3.5)$$

for all  $m, n \in \mathbb{N}$  with  $m > n$ . By induction, we get

$$\left\| 2^m f\left(\frac{x}{2^m}, 0\right) - 2^n f\left(\frac{x}{2^n}, 0\right) \right\| \leq \sum_{i=n}^{m-1} 2^i \phi\left(\frac{x}{2^i}, 0, 0\right) < \epsilon, \quad (3.6)$$

which implies that  $\{2^n f(x/2^n, 0)\}$  is a Cauchy sequence in  $Y$  for all  $x \in X$ . Since  $Y$  is complete, there exists  $h(x)$  such that

$$\left\| 2^n f\left(\frac{x}{2^n}, 0\right) - h(x) \right\| \rightarrow 0, \quad (3.7)$$

as  $n \rightarrow \infty$  for all  $x \in X$ . Since

$$\left\| h(x) - 2h\left(\frac{x}{2}\right) \right\| \leq \left\| h(x) - 2^n f\left(\frac{x}{2^n}, 0\right) \right\| + \left\| 2^n f\left(\frac{x}{2^n}, 0\right) - 2h\left(\frac{x}{2}\right) \right\| \rightarrow 0, \quad (3.8)$$

as  $n \rightarrow \infty$ ,  $h(x) = 2h(x/2)$  for all  $x \in X$ .

Similarly, we can prove that there exists a mapping  $g : X \rightarrow Y$  such that

$$\left\| 2^n f\left(0, \frac{x}{2^n}\right) - g(x) \right\| \rightarrow 0, \quad (3.9)$$

as  $n \rightarrow \infty$  and  $g(x) = 2g(x/2)$  for all  $x \in X$ .

Now, if

$$d_n(x, y, z) = f\left(\frac{x+y}{2^n}, \frac{y+z}{2^n}\right) - 2f\left(\frac{x}{2^{n+1}}, \frac{y}{2^{n+1}}\right) - 2f\left(\frac{y}{2^{n+1}}, \frac{z}{2^{n+1}}\right), \quad (3.10)$$

and  $F_n = \|2^n f((x+y)/2^n, y/2^n) - h(x+y) - g(y)\|$ , then we get

$$\begin{aligned} F_n &= \left\| 2^n d_n(x+y, 0, y) + 2^{n+1} f\left(\frac{x+y}{2^{n+1}}, 0\right) + 2^{n+1} f\left(0, \frac{y}{2^{n+1}}\right) - h(x+y) - g(y) \right\| \\ &\leq \left\| 2^n d_n(x+y, 0, y) \right\| + \left\| 2^{n+1} f\left(\frac{x+y}{2^{n+1}}, 0\right) - h(x+y) \right\| + \left\| 2^{n+1} f\left(0, \frac{y}{2^{n+1}}\right) - g(y) \right\| \end{aligned}$$

$$\begin{aligned}
&\leq 2^n \|d_n(x+y, 0, y)\| + \left\| 2^{n+1} f\left(\frac{x+y}{2^{n+1}}, 0\right) - h(x+y) \right\| + \left\| 2^{n+1} f\left(0, \frac{y}{2^{n+1}}\right) - g(y) \right\| \\
&\leq 2^n \phi\left(\frac{x+y}{2^n}, 0, \frac{y}{2^n}\right) + \left\| 2^{n+1} f\left(\frac{x+y}{2^{n+1}}, 0\right) - h(x+y) \right\| + \left\| 2^{n+1} f\left(0, \frac{y}{2^{n+1}}\right) - g(y) \right\|,
\end{aligned} \tag{3.11}$$

for all  $x, y \in X$ . Thus we obtain

$$\lim_{n \rightarrow \infty} 2^n f\left(\frac{x+y}{2^n}, \frac{y}{2^n}\right) = h(x+y) + g(y), \tag{3.12}$$

for all  $x, y \in X$ . Let  $G_n = \|2^n f(x/2^n, y/2^n) - h(x+y) - g(y) + h(y)\|$ . Then we conclude that

$$\begin{aligned}
G_n &= \left\| -2^{n-1} d_{n-1}(x, y, 0) + 2^{n-1} f\left(\frac{x+y}{2^{n-1}}, \frac{y}{2^{n-1}}\right) - 2^n f\left(\frac{y}{2^n}, 0\right) - h(x+y) - g(y) + h(y) \right\| \\
&\leq \left\| 2^{n-1} d_{n-1}(x, y, 0) \right\| + \left\| 2^{n-1} f\left(\frac{x+y}{2^{n-1}}, \frac{y}{2^{n-1}}\right) - h(x+y) - g(y) \right\| \\
&\quad + \left\| 2^n f\left(\frac{y}{2^n}, 0\right) - h(y) \right\| \\
&\leq 2^{n-1} \phi\left(\frac{x}{2^{n-1}}, \frac{y}{2^{n-1}}, 0\right) + \left\| 2^{n-1} f\left(\frac{x+y}{2^{n-1}}, \frac{y}{2^{n-1}}\right) - h(x+y) - g(y) \right\| \\
&\quad + \left\| 2^n f\left(\frac{y}{2^n}, 0\right) - h(y) \right\|,
\end{aligned} \tag{3.13}$$

for all  $x, y \in X$ . It follows from (3.12) that

$$\lim_{n \rightarrow \infty} 2^n f\left(\frac{x}{2^n}, \frac{y}{2^n}\right) = h(x+y) + g(y) - h(y), \tag{3.14}$$

for all  $x, y \in X$ . Put  $H_n = \|2^n f(x/2^n, y/2^n) - h(x) - g(y)\|$ . Since

$$\begin{aligned}
H_n &= \left\| 2^n d_n(x, 0, y) + 2^{n+1} f\left(\frac{x}{2^{n+1}}, 0\right) + 2^{n+1} f\left(0, \frac{y}{2^{n+1}}\right) - h(x) - g(y) \right\| \\
&\leq \left\| 2^n d_n(x, 0, y) \right\| + \left\| 2^{n+1} f\left(\frac{x}{2^{n+1}}, 0\right) - h(x) \right\| + \left\| 2^{n+1} f\left(0, \frac{y}{2^{n+1}}\right) - g(y) \right\| \\
&\leq 2^n \phi\left(\frac{x}{2^n}, 0, \frac{y}{2^n}\right) + \left\| 2^{n+1} f\left(\frac{x}{2^{n+1}}, 0\right) - h(x) \right\| + \left\| 2^{n+1} f\left(0, \frac{y}{2^{n+1}}\right) - g(y) \right\| \\
&\rightarrow 0,
\end{aligned} \tag{3.15}$$

as  $n \rightarrow \infty$ ,

$$\lim_{n \rightarrow \infty} 2^n f\left(\frac{x}{2^n}, \frac{y}{2^n}\right) = h(x) + g(y), \tag{3.16}$$

for all  $x, y \in X$ . It follows from (3.14) and (3.16) that

$$h(x + y) + g(y) - h(y) = h(x) + g(y), \quad (3.17)$$

for all  $x, y \in X$ . So  $h(x + y) = h(x) + h(y)$  for all  $x, y \in X$ .

Similarly, we can show that

$$g(x + y) = g(x) + g(y), \quad (3.18)$$

for all  $x, y \in X$ . Let  $F(x, y) = h(x) + g(y)$ . Then we conclude that

$$\begin{aligned} F(x + y, y + z) &= h(x) + h(y) + g(y) + g(z) \\ &= h\left(\frac{x}{2}\right) + 2h\left(\frac{y}{2}\right) + 2g\left(\frac{y}{2}\right) + 2g\left(\frac{z}{2}\right) \\ &= 2F\left(\frac{x}{2}, \frac{y}{2}\right) + 2F\left(\frac{y}{2}, \frac{z}{2}\right), \end{aligned} \quad (3.19)$$

for all  $x, y, z \in X$ .

Now, letting  $L_n = \|f(x, y) - 2^n f(x/2^n, 0) - 2^n f(0, y/2^n)\|$ , we obtain

$$\|f(x, y) - F(x, y)\| \leq \|f(x, y) - 2^n f\left(\frac{x}{2^n}, 0\right) - 2^n f\left(0, \frac{y}{2^n}\right)\| + L_n. \quad (3.20)$$

Since

$$\left\|2^n f\left(\frac{x}{2^n}, 0\right) + 2^n f\left(0, \frac{y}{2^n}\right) - F(x, y)\right\| \rightarrow 0, \quad (3.21)$$

as  $n \rightarrow \infty$ , it is sufficient to show that

$$L_n \leq \sum_{i=1}^{n-1} 2^i \left( \phi\left(\frac{x}{2^i}, 0, 0\right) + \phi\left(0, 0, \frac{y}{2^i}\right) \right) + \phi(x, 0, y). \quad (3.22)$$

We have

$$\begin{aligned} L_n &\leq \left\|f(x, y) - 2f\left(\frac{x}{2}, 0\right) - 2f\left(0, \frac{y}{2}\right)\right\| + \left\|2f\left(\frac{x}{2}, 0\right) - 2^n f\left(\frac{x}{2^n}, 0\right)\right\| \\ &\quad + \left\|2f\left(0, \frac{y}{2}\right) - 2^n f\left(0, \frac{y}{2^n}\right)\right\| \\ &\leq \sum_{i=1}^{n-1} 2^i \left( \phi\left(\frac{x}{2^i}, 0, 0\right) + \phi\left(0, 0, \frac{y}{2^i}\right) \right) + \phi(x, 0, y). \end{aligned} \quad (3.23)$$



As  $n \rightarrow \infty$ , we have

$$\|f(x, y) - F(x, y)\| \leq \sum_{i=1}^{\infty} 2^i \left( \phi\left(\frac{x}{2^i}, 0, 0\right) + \phi\left(0, 0, \frac{y}{2^i}\right) \right) + \phi(x, 0, y). \quad (3.24)$$

This completes the proof.  $\square$

**Corollary 3.2.** *Suppose that  $f : X \times X \rightarrow Y$  is a mapping such that*

$$\left\| f(x + y, y + z) - 2f\left(\frac{x}{2}, \frac{y}{2}\right) - 2f\left(\frac{y}{2}, \frac{z}{2}\right) \right\| \leq \epsilon(\|x\|^p + \|y\|^p + \|z\|^p), \quad (3.25)$$

where  $p > 1$  and  $\epsilon > 0$ . Then there exists a mapping  $F : X \times X \rightarrow Y$  satisfying (1.1),

$$\begin{aligned} \|f(x, y) - F(x, y)\| &\leq \frac{2^p}{2^p - 2} \epsilon(\|x\|^p + \|y\|^p), \\ \lim_{n \rightarrow \infty} 2^n f\left(\frac{x}{2^n}, \frac{y}{2^n}\right) &= F(x, y), \end{aligned} \quad (3.26)$$

for all  $x, y \in X$ .

**Corollary 3.3.** *Suppose that  $f : X \times X \rightarrow Y$  is a mapping such that*

$$\left\| f(x + y, y + z) - 2f\left(\frac{x}{2}, \frac{y}{2}\right) - 2f\left(\frac{y}{2}, \frac{z}{2}\right) \right\| \leq \epsilon(\|x\|^p \|y\|^p + \|y\|^p \|z\|^p + \|z\|^p \|x\|^p), \quad (3.27)$$

where  $p > 1/2$  and  $\epsilon > 0$ . Then there exists a mapping  $F : X \times X \rightarrow Y$  satisfying (1.1),

$$\begin{aligned} \|f(x, y) - F(x, y)\| &\leq \epsilon \|x\|^p \|y\|^p, \\ \lim_{n \rightarrow \infty} 2^n f\left(\frac{x}{2^n}, \frac{y}{2^n}\right) &= F(x, y), \end{aligned} \quad (3.28)$$

for all  $x, y \in X$ .

*Example 3.4.* Let  $X = Y = C[a, b]$  with the norm  $\|x\| = \sup_{t \in [a, b]} |x(t)|$ , where  $b > a > 0$ . Define  $f : C[a, b] \times C[a, b] \rightarrow C[a, b]$  by

$$f(x, y)(t) = x(t) + ty(t)^2, \quad (3.29)$$

for all  $t \in [a, b]$ , and  $\phi : X \times X \times X \rightarrow \mathbb{R}$  by

$$\phi(x, y, z) = b(\|x\| + \|y\|)^2. \quad (3.30)$$

It is not difficult to show that  $\phi$  satisfies Theorem 3.1 since

$$\begin{aligned}
& \left\| f(x+y, y+z) - 2f\left(\frac{x}{2}, \frac{y}{2}\right) - 2f\left(\frac{y}{2}, \frac{z}{2}\right) \right\| \\
&= \sup_{t \in [a, b]} \left| f(x+y, y+z)(t) - 2f\left(\frac{x}{2}, \frac{y}{2}\right)(t) - 2f\left(\frac{y}{2}, \frac{z}{2}\right)(t) \right| \\
&= \sup_{t \in [a, b]} \left| \frac{t}{2}x(t)^2 + \frac{t}{2}y(t)^2 + 2tx(t)y(t) \right| \\
&\leq b\|x\|^2 + b\|y\|^2 + 2b\|x\|\|y\| = b(\|x\| + \|y\|)^2 \\
&= \phi(x, y, z),
\end{aligned} \tag{3.31}$$

for all  $x, y, z \in X$ . Theorem 3.1 implies that there exists a mapping  $F : X \times X \rightarrow Y$  satisfying (1.1) and

$$\|f(x, y) - F(x, y)\| \leq 2b\|x\|^2, \tag{3.32}$$

for all  $x, y \in X$ .

*Example 3.5.* Let  $X$  be a normed space. Define the function  $\phi_0 : X \times X \rightarrow [0, \infty)$  by

$$\phi_0(x, y) = \left( \sup_{f, g \in B_{X'}} \left| \frac{f(x)}{g(x)} - \frac{f(y)}{g(y)} \right| \right)^p, \tag{3.33}$$

where  $B_{X'}$  is the closed unit ball of the dual space  $X'$  and  $p > 1$ . Then one can show that the function  $\phi : X \times X \times X \rightarrow [0, \infty)$ , defined by

$$\phi(x, y, z) = \phi_0(x, y) + \phi_0(y, z), \tag{3.34}$$

satisfies Theorem 3.1.

*Example 3.6.* Define  $\phi(x, y, z) = \ell^p \times \ell^p \times \ell^p \rightarrow [0, \infty)$  by

$$\phi(x, y, z) = \sum_{i=1}^{\infty} |\xi_i|^p + \sum_{i=1}^{\infty} |\eta_i|^p + \sum_{i=1}^{\infty} |\zeta_i|^p, \tag{3.35}$$

where  $x = (\xi_1, \xi_2, \dots)$ ,  $y = (\eta_1, \eta_2, \dots)$ , and  $z = (\zeta_1, \zeta_2, \dots)$ . It is obvious that  $\phi$  satisfies Theorem 3.1.

**Theorem 3.7.** Suppose that  $f : X \times X \rightarrow Y$  is a mapping and  $\phi : X \times X \times X \rightarrow [0, \infty)$  is a function such that

$$\left\| f(x+y, y+z) - 2f\left(\frac{x}{2}, \frac{y}{2}\right) - 2f\left(\frac{y}{2}, \frac{z}{2}\right) \right\| \leq \phi(x, y, z), \tag{3.36}$$

$\phi(0,0,0) = 0$  and  $\sum_{i=1}^{\infty} (1/2^i)\phi(2^i x, 2^i y, 2^i z) < \infty$  for all  $x, y, z \in X$ . Then there exists a mapping  $F : X \times X \rightarrow Y$  satisfying (1.1),

$$\begin{aligned} \|f(x, y) - F(x, y)\| &\leq \sum_{i=2}^{\infty} \frac{1}{2^i} \left( \phi(2^i x, 0, 0) + \phi(0, 0, 2^i y) \right) + \phi(x, 0, y), \\ \lim_{n \rightarrow \infty} \frac{1}{2^n} f(2^n x, 2^n y) &= F(x, y), \end{aligned} \quad (3.37)$$

for all  $x, y \in X$ .

*Proof.* It is clear that  $f(0,0) = 0$ . Now, letting  $y = z = 0$ , we obtain

$$\left\| \frac{1}{2} f(2x, 0) - f(x, 0) \right\| \leq \frac{1}{2} \phi(2x, 0, 0). \quad (3.38)$$

By induction, we have

$$\left\| \frac{1}{2^n} f(2^n x, 0) - f(x, 0) \right\| \leq \sum_{i=1}^n \frac{1}{2^i} \phi(2^i x, 0, 0). \quad (3.39)$$

It follows that, for any  $\epsilon > 0$ , there exists  $N \in \mathbb{N}$  such that, for all  $n, m \in \mathbb{N}$  with  $m > n > N$ ,

$$\left\| \frac{1}{2^m} f(2^m x, 0) - \frac{1}{2^n} f(2^n x, 0) \right\| \leq \sum_{i=n+1}^m \frac{1}{2^i} \phi(2^i x, 0, 0) < \epsilon. \quad (3.40)$$

Thus  $\{(1/2^n)f(2^n x, 0)\}$  is a Cauchy sequence in  $Y$ . The completeness of  $Y$  implies that there exists  $h(x)$  such that

$$\lim_{n \rightarrow \infty} \left\| \frac{1}{2^n} f(2^n x, 0) - h(x) \right\| = 0. \quad (3.41)$$

Since

$$\left\| h(x) - \frac{1}{2} h(2x) \right\| \leq \left\| h(x) - \frac{1}{2^n} f(2^n x, 0) \right\| + \left\| \frac{1}{2^n} f(2^n x, 0) - \frac{1}{2} h(2x) \right\| \rightarrow 0, \quad (3.42)$$

as  $n \rightarrow \infty$ ,  $h(x) = (1/2)h(2x)$ .

Similarly, we conclude that there exists a mapping  $g : X \rightarrow Y$  such that

$$\lim_{n \rightarrow \infty} \left\| \frac{1}{2^n} f(0, 2^n z) - g(z) \right\| = 0, \quad (3.43)$$

and  $g(z) = (1/2)g(2z)$ . Let

$$d_n(x, y, z) = \frac{1}{2} f(2^{n+1}(x+y), 2^{n+1}(y+z)) - f(2^n x, 2^n y) - f(2^n y, 2^n z), \quad (3.44)$$

and  $F_n = \|(1/2^{n+1})f(2^{n+1}(x+y), 2^{n+1}y) - h(x+y) - g(y)\|$ . Then we have

$$\begin{aligned}
F_n &= \left\| \frac{1}{2^n} d_n(x+y, 0, y) + \frac{1}{2^n} f(2^n(x+y), 0) + \frac{1}{2^n} f(0, 2^n y) - h(x+y) - g(y) \right\| \\
&\leq \left\| \frac{1}{2^n} d_n(x+y, 0, y) \right\| + \left\| \frac{1}{2^n} f(2^n(x+y), 0) - h(x+y) \right\| + \left\| \frac{1}{2^n} f(0, 2^n y) - g(y) \right\| \\
&\leq \frac{1}{2^n} \|d_n(x+y, 0, y)\| + \left\| \frac{1}{2^n} f(2^n(x+y), 0) - h(x+y) \right\| + \left\| \frac{1}{2^n} f(0, 2^n y) - g(y) \right\| \\
&\leq \frac{1}{2^n} \phi(2^n(x+y), 0, 2^n y) + \left\| \frac{1}{2^n} f(2^n(x+y), 0) - h(x+y) \right\| + \left\| \frac{1}{2^n} f(0, 2^{n+1} y) - g(y) \right\|,
\end{aligned} \tag{3.45}$$

for all  $x, y \in X$ . By hypothesis, (3.41) and (3.43), we get

$$\lim_{n \rightarrow \infty} \frac{1}{2^{n+1}} f(2^{n+1}(x+y), 2^{n+1}y) = h(x+y) + g(y), \tag{3.46}$$

for all  $x, y \in X$ .

Now, let  $L_n = \|(1/2^{n-1})f(2^{n-1}x, 2^{n-1}y) - h(x+y) - g(y) + h(y)\|$ . Then we obtain

$$\begin{aligned}
L_n &= \left\| -\frac{1}{2^{n-1}} d_{n-1}(x, y, 0) + \frac{1}{2^n} f(2^n(x+y), 2^n y) - \frac{1}{2^n} f(2^n y, 0) - h(x+y) - g(y) + h(y) \right\| \\
&\leq \left\| \frac{1}{2^{n-1}} d_{n-1}(x, y, 0) \right\| + \left\| \frac{1}{2^n} f(2^n(x+y), 2^n y) - h(x+y) - g(y) \right\| \\
&\quad + \left\| \frac{1}{2^{n-1}} f(2^{n-1}y, 0) - h(y) \right\| \\
&\leq \frac{1}{2^n} \phi(2^n x, 2^n y, 0) + \left\| \frac{1}{2^n} f(2^n(x+y), 2^{n-1}y) - h(x+y) - g(y) \right\| \\
&\quad + \left\| \frac{1}{2^{n-1}} f(2^{n-1}y, 0) - h(y) \right\|,
\end{aligned} \tag{3.47}$$

where the last inequality follows from (3.46). By (3.41) and (3.43), we conclude

$$\lim_{n \rightarrow \infty} \frac{1}{2^{n-1}} f(2^{n-1}x, 2^{n-1}y) = h(x+y) + g(y) - h(y), \tag{3.48}$$

for all  $x, y \in X$ . Put  $M_n = \|(1/2^{n+1})f(2^{n+1}x, 2^{n+1}y) - h(x) - g(y)\|$ . Since

$$\begin{aligned}
M_n &= \left\| \frac{1}{2^n} d_n(x, 0, y) + \frac{1}{2^n} f(2^n x, 0) + \frac{1}{2^n} f(0, 2^n y) - h(x) - g(y) \right\| \\
&\leq \left\| \frac{1}{2^n} d_n(x, 0, y) \right\| + \left\| \frac{1}{2^n} f(2^n x, 0) - h(x) \right\| + \left\| \frac{1}{2^n} f(0, 2^n y) - g(y) \right\|
\end{aligned}$$

$$\begin{aligned}
&\leq \frac{1}{2^{n+1}}\phi\left(2^{n+1}x, 0, 2^{n+1}y\right) + \left\| \frac{1}{2^n}f(2^n x, 0) - h(x) \right\| \\
&\quad + \left\| \frac{1}{2^n}f(0, 2^n y) - g(y) \right\| \\
&\rightarrow 0,
\end{aligned} \tag{3.49}$$

as  $n \rightarrow \infty$ , where the last inequality follows from (3.41) and (3.43). So we have

$$\lim_{n \rightarrow \infty} \frac{1}{2^{n+1}}f\left(2^{n+1}x, 2^{n+1}y\right) = h(x) + g(y), \tag{3.50}$$

for all  $x, y \in X$ . By (3.48) and (3.50), we have

$$h(x + y) + g(y) - h(y) = h(x) + g(y), \tag{3.51}$$

for all  $x, y \in X$ . It follows that  $h(x + y) = h(x) + h(y)$ .

Similarly, we obtain  $g(x + y) = g(x) + g(y)$ .

Now, define the mapping  $F : X \times X \rightarrow Y$  by  $F(x, y) = h(x) + g(y)$ . Then we have

$$\begin{aligned}
F(x + y, y + z) &= h(x) + h(y) + g(y) + g(z) \\
&= 2h\left(\frac{x}{2}\right) + 2h\left(\frac{y}{2}\right) + 2g\left(\frac{y}{2}\right) + 2g\left(\frac{z}{2}\right) \\
&= 2F\left(\frac{x}{2}, \frac{y}{2}\right) + 2F\left(\frac{y}{2}, \frac{z}{2}\right).
\end{aligned} \tag{3.52}$$

Now, we have

$$\begin{aligned}
\|f(x, y) - F(x, y)\| &\leq \left\| f(x, y) - \frac{1}{2^n}f(2^n x, 0) - \frac{1}{2^n}f(0, 2^n y) \right\| \\
&\quad + \left\| \frac{1}{2^n}f(2^n x, 0) + \frac{1}{2^n}f(0, 2^n y) - F(x, y) \right\|.
\end{aligned} \tag{3.53}$$

Let  $H_n = \|f(x, y) - (1/2^n)f(2^n x, 0) - (1/2^n)f(0, 2^n y)\|$ . Since we know that

$$\lim_{n \rightarrow \infty} \left\| \frac{1}{2^n}f(2^n x, 0) + \frac{1}{2^n}f(0, 2^n y) - F(x, y) \right\| = 0, \tag{3.54}$$

to prove (3.37), it is sufficient to show that

$$H_n \leq \sum_{i=2}^n \frac{1}{2^i} \left( \phi\left(2^i x, 0, 0\right) + \phi\left(0, 0, 2^i y\right) \right) + \phi(x, 0, y), \tag{3.55}$$

for all  $n \in \mathbb{N}$ . We have

$$\begin{aligned} H_n &\leq \left\| f(x, y) - \frac{1}{2}f(2x, 0) - \frac{1}{2}f(0, 2y) \right\| + \left\| \frac{1}{2}f(2x, 0) - \frac{1}{2^n}f(2^n x, 0) \right\| \\ &\quad + \left\| \frac{1}{2}f(0, 2y) - \frac{1}{2^n}f(0, 2^n y) \right\| \\ &\leq \sum_{i=2}^n \frac{1}{2^i} \left( \phi(2^i x, 0, 0) + \phi(0, 0, 2^i y) \right) + \phi(x, 0, y), \end{aligned} \quad (3.56)$$

for all  $x, y \in X$ . As  $n \rightarrow \infty$ , we conclude

$$\|f(x, y) - F(x, y)\| \leq \sum_{i=2}^{\infty} \frac{1}{2^i} \left( \phi(2^i x, 0, 0) + \phi(0, 0, 2^i y) \right) + \phi(x, 0, y), \quad (3.57)$$

for all  $x, y \in X$ . This completes the proof.  $\square$

**Corollary 3.8.** *Suppose that  $\epsilon > 0$  and  $f : X \times X \rightarrow Y$  is a mapping satisfying*

$$\left\| f(x + y, y + z) - 2f\left(\frac{x}{2}, \frac{y}{2}\right) - 2f\left(\frac{y}{2}, \frac{z}{2}\right) \right\| \leq \epsilon(\|x\|^p + \|y\|^p + \|z\|^p), \quad (3.58)$$

for some  $p < 1$ . Then there exists a mapping  $F : X \times X \rightarrow Y$  satisfying (1.1),

$$\begin{aligned} \|f(x, y) - F(x, y)\| &\leq \frac{2 - 2^p + 2^{2p-1}}{2 - 2^p} \epsilon(\|x\|^p + \|y\|^p), \\ \lim_{n \rightarrow \infty} \frac{1}{2^n} f(2^n x, 2^n y) &= F(x, y), \end{aligned} \quad (3.59)$$

for all  $x, y \in X$ .

*Example 3.9.* (1) Let  $(X, \langle \cdot, \cdot \rangle)$  be an inner product space. Then one can show that the function  $\phi : X \times X \times X \rightarrow \mathbb{R}$ , defined by

$$\phi(x, y, z) = |\langle x, y \rangle|^p + |\langle y, z \rangle|^p \quad (p < 1), \quad (3.60)$$

satisfies Theorem 3.7.

(2) Suppose that  $(X, \|\cdot\|)$  is a normed space. Then the function  $\phi : X \times X \times X \rightarrow \mathbb{R}$ , defined by  $\phi(x, y, z) = \|x\|\|y\|\|z\|$ , satisfies Theorem 3.7.

#### 4. A Fixed Point Approach to the Stability

In this section, we apply the fixed point method to prove the stability of the functional equation (1.1).

Let  $X$  be a set. A function  $d : X \times X \rightarrow [0, \infty]$  is called a *generalized metric* on  $X$  if  $d$  satisfies the following conditions:

- (1)  $d(x, y) = 0$  for all  $x, y \in X$  if and only if  $x = y$ ;
- (2)  $d(x, y) = d(y, x)$  for all  $x, y \in X$ ;
- (3)  $d(x, z) \leq d(x, y) + d(y, z)$  for all  $x, y, z \in X$ .

We now introduce one of the fundamental results of fixed point theory.

**Theorem 4.1** (see [13]). *Let  $(X, d)$  be a complete generalized metric space and let  $J : X \rightarrow X$  be a contractive mapping with constant  $L$ . Then for each  $x \in X$ , either*

$$d(J^n x, J^{n+1} x) = \infty, \quad (4.1)$$

for all  $n \in \mathbb{N}$  or there exists an  $n_0 \in \mathbb{N}$  such that

- (a)  $d(J^n x, J^{n+1} x) < \infty$  for all  $n \geq n_0$ ;
- (b) the sequence  $\{J^n x\}$  converges to fixed point  $x_0$  of  $J$ ;
- (c)  $x_0$  is the unique fixed point of  $J$  in the set  $Y = \{y \in X : d(J^{n_0} x, y) < \infty\}$ ;
- (d)  $d(y, y_0) \leq (1/(1-L))d(y, Jy)$  for all  $y \in Y$ .

In 1996, Isac and Rassias [14] were the first to provide applications of stability theory of functional equations for the proof of new fixed point theorems with applications. By using fixed point methods, the stability problems of several functional equations have been extensively investigated by a number of authors (see [15–22]).

**Theorem 4.2.** *Suppose that  $X$  is a vector space and  $Y$  is a Banach space. Let  $f : X \times X \rightarrow Y$  be a mapping and let  $\phi : X \times X \times X \rightarrow [0, \infty)$  be a function satisfying the following conditions:*

- (a)  $\|f(x + y, y + z) - 2f(x/2, y/2) - 2f(y/2, z/2)\| \leq \phi(x, y, z)$ ;
- (b)  $\phi(0, 0, 0) = 0$ ;
- (c) for all  $x, y \in X$ ,

$$\lim_{n \rightarrow \infty} \frac{1}{2^n} \phi(2^n x, 2^n y, 2^n z) = 0; \quad (4.2)$$

- (d) there exists a number  $L \in [0, 1)$  such that

$$\phi(x, y, \sigma(x, y)) \leq 2L\phi\left(\frac{x}{2}, \frac{y}{2}, \sigma\left(\frac{x}{2}, \frac{y}{2}\right)\right), \quad (4.3)$$

where  $\sigma : X \times X \rightarrow X$  is a mapping such that  $\sigma(x, 0) = 0$  for all  $x \in X$ .

Then there exists a unique mapping  $F : X \times X \rightarrow Y$  satisfying (1.1) and

$$\|f(x, y) - F(x, y)\| \leq \frac{1}{2-2L} \phi(x, y, \sigma(x, y)), \quad (4.4)$$

for all  $x, y \in X$ .

*Proof.* Consider the set

$$\mathcal{S} = \{g : X \times X \rightarrow Y\}, \quad (4.5)$$

and define the generalized metric on  $\mathcal{S}$  by

$$d(g, h) = \inf\{C : \|g(x, y) - h(x, y)\| \leq C\phi(x, y, \sigma(x, y)), \quad \forall(x, y) \in X \times X\}. \quad (4.6)$$

It is easy to show that  $(\mathcal{S}, d)$  is complete.

Now, we consider the linear mapping  $J : \mathcal{S} \rightarrow \mathcal{S}$  such that

$$Jg(x, y) = \frac{1}{2}g(2x, 2y), \quad (4.7)$$

for all  $x, y \in X$ . The definition of  $J$  implies that

$$d(Jg, Jh) \leq Ld(g, h), \quad (4.8)$$

for all  $g, h \in \mathcal{S}$ . Replacing  $x$  by  $2x$  and putting  $y = z = 0$  in the first statement, we get

$$\|f(2x, 0) - 2f(x, 0)\| \leq \phi(x, 0, 0) = \phi(2x, 0, \sigma(2x, 0)), \quad (4.9)$$

for all  $x \in X$ . So

$$\left\|f(x, 0) - \frac{1}{2}f(2x, 0)\right\| \leq \frac{1}{2}\phi(2x, 0, \sigma(2x, 0)), \quad (4.10)$$

for all  $x \in X$ . Hence  $d(f, Jf) \leq 1/2$ . By Theorem 4.1, there exists a mapping  $F : X \times X \rightarrow Y$  such that  $F$  is a fixed point of  $J$ , that is,

$$F(2x, 2y) = 2F(x, y), \quad (4.11)$$

for all  $x, y \in X$ . The mapping  $F$  is a unique fixed point of  $J$  in the set

$$\mathcal{S}' = \{g \in \mathcal{S} : d(f, g) < \infty\}. \quad (4.12)$$

This implies that there exists a  $C < \infty$  such that

$$\|F(x, y) - f(x, y)\| \leq C\phi(x, y, \sigma(x, y)), \quad (4.13)$$

for all  $x, y \in X$ . Also, we have

$$d(J^n f, F) \rightarrow 0, \quad (4.14)$$



as  $n \rightarrow \infty$ . It follows that

$$\lim_{n \rightarrow \infty} \frac{1}{2^n} f(2^n x, 2^n y) = F(x, y), \quad (4.15)$$

for all  $x, y \in X$ . By the third statement of Theorem 4.1, we have  $d(f, F) \leq (1/(1-L))d(f, Jf)$ . This implies that

$$d(f, F) \leq \frac{1}{2-2L}. \quad (4.16)$$

Thus, by the definition of  $d(\cdot, \cdot)$ , we conclude

$$\|f(x, y) - F(x, y)\| \leq \frac{1}{2-2L} \phi(x, y, \sigma(x, y)), \quad (4.17)$$

for all  $x, y \in X$ . Let

$$D_F = F(x + y, y + z) - 2F\left(\frac{x}{2}, \frac{y}{2}\right) - 2F\left(\frac{y}{2}, \frac{z}{2}\right). \quad (4.18)$$

It follows from (4.15) that

$$\begin{aligned} \|D_F\| &= \lim_{n \rightarrow \infty} \frac{1}{2^n} \left\| f(2^n(x+y), 2^n(y+z)) - 2f\left(2^n \frac{x}{2}, 2^n \frac{y}{2}\right) - 2f\left(2^n \frac{y}{2}, 2^n \frac{z}{2}\right) \right\| \\ &\leq \lim_{n \rightarrow \infty} \frac{1}{2^n} \phi(2^n x, 2^n y, 2^n z) = 0. \end{aligned} \quad (4.19)$$

So we have

$$F(x + y, y + z) = 2F\left(\frac{x}{2}, \frac{y}{2}\right) + 2F\left(\frac{y}{2}, \frac{z}{2}\right), \quad (4.20)$$

for all  $x, y, z \in X$ . So  $F$  satisfies (1.1). This completes the proof.  $\square$

## Acknowledgments

Y. J. Cho was supported by the Korea Research Foundation Grant funded by the Korean Government KRF-2008-313-C00050.

## References

- [1] S. M. Ulam, *Problems in Modern Mathematics*, Science Editions John Wiley & Sons, New York, NY, USA, 1964.
- [2] T. Aoki, "On the stability of the linear transformation in Banach spaces," *Journal of the Mathematical Society of Japan*, vol. 2, pp. 64–66, 1950.

- [3] J. Bae and W. Park, "On the solution of a bi-Jensen functional equation and its stability," *Bulletin of the Korean Mathematical Society*, vol. 43, no. 3, pp. 499–507, 2006.
- [4] Z. Gajda, "On stability of additive mappings," *International Journal of Mathematics and Mathematical Sciences*, vol. 14, no. 3, pp. 431–434, 1991.
- [5] P. Găvruta, "A generalization of the Hyers-Ulam-Rassias stability of approximately additive mappings," *Journal of Mathematical Analysis and Applications*, vol. 184, no. 3, pp. 431–436, 1994.
- [6] D. H. Hyers, "On the stability of the linear functional equation," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 27, pp. 222–224, 1941.
- [7] C. Park, Y. S. Cho, and M. Han, "Functional inequalities associated with Jordan-von Neumann-type additive functional equations," *Journal of Inequalities and Applications*, vol. 2007, Article ID 41820, 13 pages, 2007.
- [8] C. Park and J. Cui, "Generalized stability of  $C^*$ -ternary quadratic mappings," *Abstract and Applied Analysis*, vol. 2007, Article ID 23282, 6 pages, 2007.
- [9] C. Park and A. Najati, "Homomorphisms and derivations in  $C^*$ -algebras," *Abstract and Applied Analysis*, vol. 2007, Article ID 80630, 12 pages, 2007.
- [10] C. Park, W.-G. Park, and A. Najati, "Functional equations related to inner product spaces," *Abstract and Applied Analysis*, vol. 2009, Article ID 907121, p. 11, 2009.
- [11] T. M. Rassias, "On the stability of the linear mapping in Banach spaces," *Proceedings of the American Mathematical Society*, vol. 72, no. 2, pp. 297–300, 1978.
- [12] T. M. Rassias and P. Semrl, "On the behavior of mappings which do not satisfy Hyers-Ulam stability," *Proceedings of the American Mathematical Society*, vol. 114, no. 4, pp. 989–993, 1992.
- [13] J. B. Diaz and B. Margolis, "A fixed point theorem of the alternative, for contractions on a generalized complete metric space," *Bulletin of the American Mathematical Society*, vol. 74, pp. 305–309, 1968.
- [14] G. Isac and T. M. Rassias, "Stability of  $\varphi$ -additive mappings: applications to nonlinear analysis," *International Journal of Mathematics and Mathematical Sciences*, vol. 19, no. 2, pp. 219–228, 1996.
- [15] L. Cădariu and V. Radu, "Fixed points and the stability of Jensen's functional equation," *Journal of Inequalities in Pure and Applied Mathematics*, vol. 4, no. 1, Article ID 4, 2003.
- [16] L. Cădariu and V. Radu, "On the stability of the Cauchy functional equation: a fixed point approach," in *Grazer Mathematische Berichte*, vol. 346, pp. 43–52, 2004.
- [17] L. Cădariu and V. Radu, "Fixed point methods for the generalized stability of functional equations in a single variable," *Fixed Point Theory and Applications*, vol. 2008, Article ID 749392, 15 pages, 2008.
- [18] D. Mihecontractible, "The fixed point method for fuzzy stability of the Jensen functional equation," *Fuzzy Sets and Systems*, vol. 160, no. 11, pp. 1663–1667, 2009.
- [19] M. Mirzavaziri and M. S. Moslehian, "A fixed point approach to stability of a quadratic equation," *Bulletin of the Brazilian Mathematical Society*, vol. 37, no. 3, pp. 361–376, 2006.
- [20] C. Park, "Fixed points and Hyers-Ulam-Rassias stability of Cauchy-Jensen functional equations in Banach algebras," *Fixed Point Theory and Applications*, vol. 2007, Article ID 50175, 15 pages, 2007.
- [21] C. Park, "Generalized Hyers-Ulam stability of quadratic functional equations: a fixed point approach," *Fixed Point Theory and Applications*, vol. 2008, Article ID 493751, 9 pages, 2008.
- [22] V. Radu, "The fixed point alternative and the stability of functional equations," *Fixed Point Theory*, vol. 4, no. 1, pp. 91–96, 2003.



# Hindawi

Submit your manuscripts at  
<http://www.hindawi.com>

