



RESEARCH

Open Access

Mechanisms involved in PGE₂-induced transactivation of the epidermal growth factor receptor in MH₁C₁ hepatocarcinoma cells

Ingun Heiene Tveteraas^{1*}, Kristin Meisdalen Müller¹, Monica Aasrum¹, John Ødegård¹, Olav Dajani², Tormod Guren², Dagny Sandnes¹ and Thoralf Christoffersen¹

Abstract

Background: It is important to understand the mechanisms by which the cells integrate signals from different receptors. Several lines of evidence implicate epidermal growth factor (EGF) receptor (EGFR) in the pathophysiology of hepatocarcinomas. Data also suggest a role of prostaglandins in some of these tumours, through their receptors of the G protein-coupled receptor (GPCR) family. In this study we have investigated mechanisms of interaction between signalling from prostaglandin receptors and EGFR in hepatocarcinoma cells.

Methods: The rat hepatocarcinoma cell line MH₁C₁ and normal rat hepatocytes in primary culture were stimulated with EGF or prostaglandin E₂ (PGE₂) and in some experiments also PGF_{2α}. DNA synthesis was determined by incorporation of radiolabelled thymidine into DNA, phosphorylation of proteins in signalling pathways was assessed by Western blotting, mRNA expression of prostaglandin receptors was determined using qRT-PCR, accumulation of inositol phosphates was measured by incorporation of radiolabelled inositol, and cAMP was determined by radioimmunoassay.

Results: In the MH₁C₁ hepatocarcinoma cells, stimulation with PGE₂ or PGF_{2α} caused phosphorylation of the EGFR, Akt, and ERK, which could be blocked by the EGFR tyrosine kinase inhibitor gefitinib. This did not occur in primary hepatocytes. qRT-PCR revealed expression of EP1, EP4, and FP receptor mRNA in MH₁C₁ cells. PGE₂ stimulated accumulation of inositol phosphates but not cAMP in these cells, suggesting signalling via PLCβ. While pretreatment with EP1 and EP4 receptor antagonists did not inhibit the effect of PGE₂, pretreatment with an FP receptor antagonist blocked the phosphorylation of EGFR, Akt and ERK. Further studies suggested that the PGE₂-induced signal was mediated via Ca²⁺ release and not PKC activation, and that it proceeded through Src and shedding of membrane-bound EGFR ligand precursors by proteinases of the ADAM family.

Conclusion: The results indicate that in MH₁C₁ cells, unlike normal hepatocytes, PGE₂ activates the MEK/ERK and PI3K/Akt pathways by transactivation of the EGFR, thus diversifying the GPCR-mediated signal. The data also suggest that the underlying mechanisms in these cells involve FP receptors, PLCβ, Ca²⁺, Src, and proteinase-mediated release of membrane-associated EGFR ligand(s).

Keywords: EGF receptor, Prostaglandin receptors, Transactivation, Hepatocarcinoma cells, Hepatocytes

* Correspondence: ingunhei@medisin.uio.no

¹Department of Pharmacology, Institute of Clinical Medicine, Faculty of Medicine, University of Oslo, P.O.Box 1057 Blindern, N-0316 Oslo, Norway
Full list of author information is available at the end of the article

Background

Malignant cells are exposed to a variety of active agents, including hormones, peptide growth factors, cytokines, and many other locally acting substances such as prostaglandins, which together control or modulate the cellular functions. It is of interest to understand the mechanisms by which the cells integrate signals from different bioactive molecules via their receptors. A notable example is the interaction between pathways from G protein-coupled receptors (GPCRs) and receptor tyrosine kinases (RTKs). Studies in many cells have shown that signals from GPCRs may involve interaction with the epidermal growth factor receptor (EGFR), an ErbB family RTK [1-5]. EGFR, which serves important functions in normal cells [6,7], is involved in several malignancies [8,9], and is a target of novel antitumour therapies [10,11]. In studies including tumour cells from colon and pancreatic cancer, we have found that different mechanisms may be involved in the interaction of pathways from GPCRs and EGFR [12].

EGFR conveys strong mitogenic stimulation in normal hepatocytes [13-16], and several lines of evidence suggest a role of EGFR in hepatocarcinogenesis [17-20]. For example, overexpression of the EGFR agonist transforming growth factor alpha (TGF α) in mice causes hepatic hyperplasia and tumour formation [21,22], and EGFR is upregulated in a majority of human hepatocarcinomas [23]. Inhibition of the EGFR by antibodies or tyrosine kinase blockers can attenuate the growth of hepatocarcinoma cells in vitro [24,25], and are currently being tested in clinical trials in hepatocarcinomas [26].

Prostaglandins, acting through different receptors of the GPCR family, regulate many cellular functions [27]. In epithelial cells, prostaglandins often enhance proliferation and survival, and several lines of evidence implicate them in oncogenesis [28]. In many tumours, cyclooxygenases (COX-1 and COX-2), which catalyze the rate-limiting step in prostaglandin synthesis, are overexpressed, and the levels of prostaglandins, notably prostaglandin E₂ (PGE₂), are elevated [28-31]. In hepatocytes, PGE₂ and other prostaglandins enhance DNA synthesis [15,32-34], and COX-2 is overexpressed in many hepatocarcinomas [35,36].

In the study presented here we examined the Morris hepatocarcinoma cell line MH₁C₁, which was chosen due to its responsiveness to both EGF and the prostaglandins PGE₂ and PGF_{2 α} , and investigated the interaction between the pathways mediated by prostaglandin receptors and EGFR. We previously observed that while there was no evidence of transactivation of EGFR induced by prostaglandins or other GPCR agonists in hepatocytes, PGE₂ induced phosphorylation of the EGFR in the MH₁C₁ cells [37,38]. We have now investigated further the signalling mechanisms involved in this effect.

Methods

Chemicals

Dulbecco's Modified Eagle's Medium, Dulbecco's phosphate-buffered saline, William's Medium E, glutamine, and Pen-Strep (10.000 U/ml) were from Lonza (Verviers, Belgium). HEPES was from Gibco (Grand Island, NY). Dexamethasone, insulin, bovine serum albumin, collagen (type I, rat tail), prostaglandin F_{2 α} (Tris salt) and epidermal growth factor (EGF) were obtained from Sigma-Aldrich (St. Louis, MO). GF109203X ([2-[1-(3-dimethylaminopropyl)-1*H*-indol-3-yl]-male-imide]) and GM6001/Galardin (N-[(2R)-2 (hydroxamidocarbonylmethyl)-4-methylpentanoyl]-L-tryptophan methylamide) were from Calbiochem (San Diego, CA). Gefitinib was a gift from AstraZeneca (Cheshire, UK). [6-³H]-thymidine (20–30 Ci/mmol) and *myo*-[2-³H]inositol (15.0 Ci/mmol) were from PerkinElmer (Boston, MA). AL8810 (9 α ,15R-dihydroxy-11 β -fluoro-15-(2,3-dihydro-1*H*-inden-2-yl)-16,17,18,19,20-pentanor-prosta-5Z,13E-dien-1-oic acid), L161982 (N-[[4'-[[3-butyl-1,5-dihydro-5-oxo-1-[2-(trifluoromethyl)phenyl]-4*H*-1,2,4-triazol-4-yl]methyl][1,1'-biphenyl]-2-yl]sulfonyl]-3-methyl-2-thiophenecarboxamide), (+)fluprostenol, and prostaglandin E₂ (PGE₂) were from Cayman Chemical (Ann Arbor, MI). SC51322 (8-chloro-2-[3-[(2-furanylmethyl)thio]-1-oxopropyl]hydrazide, dibenz[b,f][1,4]oxazepine-10(11*H*)-carboxylic acid) was obtained from BIOMOL Research Laboratories (Plymouth Meeting, PA). The Src inhibitor CGP77675 was a gift from Novartis Pharma AG (Basel, Switzerland). All other chemicals were of analytical quality.

Antibodies against phosphorylated Akt^{Ser473}, total Akt, dually phosphorylated ERK^{Thr202/Tyr204}, GAPDH and phospho-Shc^{Tyr239/240} were obtained from Cell Signaling Technology (Boston, MA). Antibody against phospho-EGF receptor^{Tyr1173} was obtained from Invitrogen. Anti-ERK antibody was from Upstate/Millipore (Billerica, MA). Secondary antibodies were purchased from Bio-Rad Laboratories (Hercules, CA) and Licor Biosciences (Lincoln, NE).

Cells and culturing

The rat hepatocarcinoma cell line MH₁C₁, derived from a Morris hepatoma [39], was obtained from ATCC (Manassas, VA). The cells were seeded onto Costar plastic flasks and cultured in Dulbecco's Modified Eagle's medium. The Medium was supplemented with horse serum (10%), glutamine (2 mM), and 100 U/ml Pen-Strep. The cultures were kept in a humidified 5% CO₂ incubator at 37°C. Cells were seeded onto culture wells at a density of 40 000–50 000 cells per cm². After 24 hours, the medium was changed and the cells were cultured under serum-free conditions 24 h prior to stimulation.

Hepatocytes were isolated from male Wistar rats as previously described [40]. The hepatocytes were seeded onto Costar plastic culture wells at a density of 15 000–20 000 per cm². The culture medium was a serum-free 1:1 combination of William's Medium E and Dulbecco's Modified Eagle's Medium. The medium was supplemented with 100 U/ml Pen-Strep, collagen (3 µg/ml), insulin (100 nM) and dexamethasone (25 nM).

Immunoblotting

Aliquots containing ~30000 MH₁C₁ cells or hepatocytes (total cell lysate prepared in Laemmli or RIPA buffer) were electrophoresed on 6–12% (w/v) polyacrylamide gels (acrylamide: N,N'-bis-methylene acrylamide 30:1). This was followed by protein electrotransfer to nitrocellulose membranes and immunoblotting with antibodies against proteins as described in the figures. Usually the same membrane was stripped and reincubated with different antibodies, and then one single loading control was used as the final incubation. Immunoreactive bands were visualized with enhanced chemiluminescence using Lumi-GLO (KPL Protein research Products, Gaithersburg, MD) or by infrared imaging using Odyssey Infrared Imaging System, supplied by Licor Biosciences (Lincoln, NE).

RNA isolation and cDNA synthesis

RNA from MH₁C₁ cells was isolated with Qiagen RNeasy kit according to the manufacturer's instructions, and was treated with DNase. The integrity of RNA was evaluated by ethidium bromide agarose gel electrophoresis, and the quantity and purity was measured spectrophotometrically (OD 260/280). cDNA was synthesized from 1.0 µg RNA with Superscript[®] III reverse transcriptase (Invitrogen) according to manufacturer's protocol. Reactions without reverse transcriptase were run in parallel to control for contamination with chromosomal DNA. Standard curves with RNA ranging from 0.25 to 2.0 µg of total RNA were made to control for the reverse transcription and PCR quantification.

Quantitative real-time PCR (qRT-PCR)

The cDNA was analyzed in triplicate by real time quantitative PCR on an ABI Prism 7900 HT Sequence detector (Applied Biosystems) with the following cycling parameters: 50°C for 2 min, 95°C for 10 min and 40 cycles of 15 s at 95°C and 60 s at 60°C, followed by melting point analysis when using SYBR green. Raw data were collected and analyzed in the Sequence Detector Software (SDS ver. 2.2, Applied Biosystems), and cycle of threshold value (Ct) was calculated from each amplification plot. Standard curves (Ct value versus log initial RNA concentration) were used to calculate the relative input amount of RNA for each sample based on the Ct

value [41]. Satisfactory and comparable amplification efficiency was verified by the slopes of standard curves. Primers were designed using Primer Express[®] software v2.1 (ABI Prism, Applied Biosystems), and were validated by the production of single products of expected size on agarose gels, as well as uniformity of melting temperature, which was routinely performed. Prostaglandin receptor cDNA was detected with SYBR Green methodology and the following primers: EP1: forward 5'-CCT GCT GGT ATT GGT GGT GTT-3' and reverse 5'-GGG GTA GGA GGC GAA GAA GTT-3'; EP2: forward 5'-GCT CCC TGC CTT TCA CAA TCT-3' and reverse 5'-GGA CTG GTG GTC TAA GGA TGA CA-3'; EP3: forward 5'-GGT CGC CGC TAT TGA TAA TGA T-3' and reverse 5'-CAG GCG AAC GGC GAT TAG-3'; EP4: forward 5'-CTC GTG GTG CGA GTG TTC AT-3' and reverse 5'-TGT AGA TCC AAG GGT CCA GGA T-3'; FP: forward 5'-GTC ATT CAG CTC CTG GCC ATA-3' and reverse 5'-AGC GTC GTC TCA CAG GTC ACT-3'. GAPDH cDNA was quantified using the dual hybridization probe Double Dye oligonucleotide 5' labelled with the fluorescent dye Yakima yellow and quenched with Dark Quencher, 5'-CTC ATG ACC ACA GTC CAT GCC ATC ACT-3' and the following primers: forward 5'-CCA AGG TCA TCC ATG ACA ACT T-3' and reverse 5'-AGG GGC CAT CCA CAG TCT T-3'. Results were normalized to GADPH.

Accumulation of inositol phosphates and cAMP

[³H]inositol, 5 µCi/well was added simultaneously with the serum-free medium. 30 minutes before agonist stimulation for 30 minutes in serum-starved cells, medium was removed and replaced with Krebs-Ringer-Hepes buffer pH 7.4, containing 10 mM glucose and 15 mM LiCl. MH₁C₁ cells were stimulated with PGE₂, fluprostenol or isoproterenol as indicated, and the reaction was stopped by removing buffer and adding 1 ml ice-cold 0.4 M perchloric acid. Samples were harvested and neutralized with 1.5 M KOH, 60 mM EDTA and 60 mM Hepes, in the presence of Universal indicator. The neutralized supernatants were applied on columns containing 1 ml Dowex AG 1-X8 resin. The columns were washed with 20 ml distilled water and 10 ml 5 mM sodium tetraborate/60 mM ammonium formate, and inositol phosphates were eluted with 10 ml 1 M ammonium formate/0.1 M formic acid. cAMP was determined by radioimmunoassay as previously described [42].

Measurement of DNA synthesis

MH₁C₁ cells were seeded onto culture wells, and after 24 hours, the medium was changed and the cells were cultured under serum-free conditions. 24 h after change to serum-free medium, cells were treated with various concentrations of gefitinib and harvested at 48 hours,

after three hours of pulsing with [³H]thymidine. DNA synthesis was measured as the amount of radioactivity incorporated into DNA as previously described [34].

Results


In preliminary experiments we investigated the effect of PGE₂ in the rat hepatocarcinoma cell lines MH₁C₁, McA7777, and M4IIE, and the human hepatocarcinoma cell line HepG2. Although some of these cell lines had strong responses to EGF (data not shown), the MH₁C₁ were the only cells showing consistent responses to both EGF and prostaglandins, and we therefore used these cells in further experiments.

Transactivation of EGFR induced by PGE₂ and PGF_{2α} in MH₁C₁ cells

We previously observed that in the MH₁C₁ cells, unlike normal hepatocytes, PGE₂ induced phosphorylation of the EGFR and activated ERK by a mechanism that was sensitive to EGFR inhibition [37]. Further investigation (Figure 1), showed that in addition to inducing phosphorylation of

EGFR and ERK, PGE₂ treatment also led to phosphorylation of Akt. All these effects were inhibited by gefitinib (1 μM) (Figure 1A), providing further support for a transactivation of EGFR in the MH₁C₁ cells. In contrast, the effects of PGE₂ on ERK and Akt in hepatocytes were not dependent on the EGFR, since they were not inhibited by gefitinib (Figure 1B). We also observed that in the MH₁C₁ cells, the phosphorylation of the EGFR was somewhat slower after stimulation with PGE₂ than with EGF (data not shown), suggesting an indirect mechanism consistent with PGE₂-induced transactivation. As shown in Figure 1C, PGF_{2α} also induced a gefitinib-sensitive phosphorylation of EGFR, Akt and ERK in these cells.

Figure 1D shows that the EGFR tyrosine kinase blocker gefitinib dose-dependently inhibited DNA synthesis in MH₁C₁, indicating that EGFR is involved in the growth in these cells. Most likely there is an autocrine release of EGFR agonist(s) in these long-term experiments (48 h culturing). This has not been explored further in the present study, as the experiments below focus on early receptor-mediated mechanisms.



Prostaglandin receptors and involvement of PLC β

We next investigated which prostaglandin receptors are expressed in the MH₁C₁ cells. qRT-PCR analysis revealed mRNA expression of EP1, EP4, and FP subtypes of prostaglandin receptors, whereas only traces of EP3 receptor mRNA were present and no EP2 expression was detected (Figure 2A). The hepatocytes expressed EP2, EP3, EP4, and FP (Figure 2B).

The available evidence indicates that the EP4 receptors are coupled to G_s proteins and adenylyl cyclase activity and thereby cAMP elevation, and that FP receptors couple to G_q proteins which mediate activation of phospholipase C- β (PLC β) leading to formation of inositol trisphosphate (InsP₃) and diacylglycerol (DAG) [27,43]. The G proteins and signalling mechanisms stimulated by the EP1 receptors are not fully clarified [43,44]. PGE₂ has high affinity for EP1 and EP4 receptors, and while the FP receptor has the highest affinity for PGF_{2 α} , PGE₂ also binds to this receptor [27]. In the MH₁C₁ cells no cAMP response to PGE₂ could be detected, although the cells had a functional adenylyl cyclase, as shown by their marked cAMP elevation in response to the β -adrenergic agonist isoproterenol (Figure 2C left). In contrast, PGE₂ stimulated accumulation

of inositol phosphates (Figure 2C right). Thus, it is likely that PGE₂ induces signalling through PLC β activation in these cells.

To investigate which receptors are involved in the EGFR transactivation by PGE₂, we studied the effect of pretreating the cells with selective inhibitors of different prostaglandin receptors. The results suggested that EP4 did not mediate this transactivation since the EP4 receptor antagonist L161982 did not inhibit the effect of PGE₂ on the phosphorylation of EGFR, Akt, or ERK (Figure 3A), consistent with the lack of PGE₂-induced cAMP response in these cells (Figure 2C). We then examined the roles of EP1 and FP receptors. Pretreatment of the cells with 10 μ M of the EP1 receptor antagonist SC51322 did not affect PGE₂-induced phosphorylation of EGFR, Akt, or ERK (Figure 3B). In contrast, the FP receptor antagonist AL8810 at 10 μ M significantly inhibited the effect of PGE₂ on the phosphorylation of ERK, while 100 μ M inhibited phosphorylation of EGFR and Akt and blocked the effects on ERK almost completely (Figure 3C). These concentrations of AL8810 were not toxic to the cells. Although AL8810 is a less potent antagonist than L161982 or SC51322 [27,45,46], it was the only antagonist that had effect at 10 μ M. It was previously shown that at 10 μ M,





Figure 2 Prostaglandin receptors and cAMP and PLC β responses. **A)** and **B)** Expression of prostaglandin receptor mRNA in MH₁C₁ cells (data from three experiments, measured in triplicate) and hepatocytes (data from one experiment measured in triplicate). Quantitative RT-PCR of EP1, EP2, EP3, EP4 and FP normalized to GAPDH. RNA was isolated as described in Materials and Methods. * not detected # low levels-not quantifiable. **C)** Left: Accumulation of cAMP in MH₁C₁ cells after stimulation with either PGE₂ (100 μ M) or isoproterenol (10 μ M) in the presence of 0.5 mM IBMX. cAMP was measured after 3 minutes. Right: Accumulation of inositol phosphates in MH₁C₁ cells after stimulation with PGE₂ (100 μ M) for 30 minutes in the presence of 15 mM LiCl. The data shown are mean \pm S.E.M of three independent experiments.




AL8810 did not inhibit functional responses through other prostaglandin receptors, suggesting that it is a selective antagonist at the FP receptor [45]. Further support for a functional role of FP receptors in these cells was obtained in the results given in Figure 3D, demonstrating that AL8810 inhibited the inositol phosphate accumulation induced by the FP receptor agonist fluprostenol. Taken together, these results suggest that the PGE₂-induced transactivation of EGFR in MH₁C₁ hepatoma cells is mediated primarily by FP receptors and signalling via Gq and PLCβ.

Evidence of a role for Ca²⁺, but not PKC, in the PGE₂-induced transactivation of EGFR

We next tried to determine which pathways downstream of PLCβ are mediating the PGE₂-induced transactivation of EGFR. InsP₃ and DAG stimulate cytosolic Ca²⁺ release and protein kinase C (PKC) activity, respectively.

Pretreatment of the cells with the PKC inhibitor GF109203X did not prevent the effects of PGE₂ on the phosphorylation of the EGFR, ERK, or Akt in the MH₁C₁ cells (Figure 4A). Furthermore, the data in Figure 4B, comparing PGE₂ and the direct PKC activator tetradecanoylphorbol acetate (TPA), showed that TPA did not mimic the effect of PGE₂ on Akt, and its stimulation of ERK, unlike the effect of PGE₂, was blocked by GF109203X. Interestingly, pretreatment of the cells with GF109203X consistently increased basal and PGE₂-induced Akt phosphorylation in the cells. This might result from a reduced feedback inhibition by PKC [47]. In contrast to TPA, thapsigargin, which increases the intracellular Ca²⁺ level by inhibiting the 'sarco/endoplasmic reticulum Ca²⁺-ATPase' (SERCA) pump [48], induced gefitinib-sensitive phosphorylation of EGFR, ERK, and Akt (Figure 4C). Taken together, these data suggest that Ca²⁺ rather than PKC mediates the PGE₂-induced transactivation of the EGFR in these cells.



Role of Src and metalloproteinases in the transactivation of the EGFR


To further elucidate mechanisms involved in transactivation of the EGFR, we investigated the effects of Src inhibitors. As shown in Figure 5A, pretreatment of the cells with the Src inhibitor CGP77675 almost completely abolished the PGE₂-induced phosphorylation of EGFR and the activation of ERK and Akt, but, in contrast, had little or no effect on the phosphorylation of these proteins elicited by EGF. The Src inhibitor PP2 similarly prevented the phosphorylation of ERK in response to PGE₂, while the response to EGF was not significantly affected (Figure 5B). These results suggest an involvement of a Src family kinase in the PGE₂-induced transactivation of EGFR in MH₁C₁ cells.

Previous evidence has implicated proteinases of the 'a-disintegrin-and-metalloproteinase' (ADAM) family in EGFR transactivation by GPCRs in various cells [2,49,50]. To test the role of ADAMs in the PGE₂-induced EGFR transactivation in MH₁C₁, we pretreated the cells with GM6001, which is a broad-spectrum metalloproteinase inhibitor [50]. This pretreatment resulted in complete inhibition of PGE₂-induced

phosphorylation of EGFR, ERK, and Akt, while the EGF-induced phosphorylation of these proteins was not affected (Fig 5C and D), indicating that the transactivation is dependent on mechanisms involving ADAM-mediated release of EGFR ligand(s). We also examined the effect of this inhibitor in the primary cultures of rat hepatocytes, and found neither inhibition of PGE₂-induced phosphorylation of ERK and Akt in these cells nor any effect on EGF-induced phosphorylation of EGFR, ERK and Akt (Figure 5E).

Discussion


We have shown that in the MH₁C₁ hepatocarcinoma cells stimulation with PGE₂ or PGF_{2α} causes phosphorylation of the EGFR and an EGFR-dependent phosphorylation of ERK and Akt, indicating that these prostaglandins induced transactivation of EGFR. Further study of the PGE₂ effect suggested that the transactivation was mediated by the Gq-coupled FP receptor and activation of PLCβ with downstream signalling by Ca²⁺ release, Src, and ADAM-mediated shedding of membrane-bound EGFR ligand precursors. In contrast, in primary hepatocytes, PGE₂ did not phosphorylate the EGFR, and gefitinib did not prevent



phosphorylation of Akt or ERK after PGE₂-stimulation, which lends further support to our previous data suggesting that GPCR agonists do not transactivate the EGFR in normal rat hepatocytes, but rather signal via mechanisms that synergistically enhance the effects of EGF [34,37,38,51,52] (Figure 6).

Different receptors and pathways may be involved in mitogenic and tumour-promoting effects of prostaglandins [28]. qRT-PCR analysis showed that the prostaglandin receptors expressed in these cells are EP1, EP4, and FP. No significant increase in cAMP accumulation was detected, in accordance with previous results [53], suggesting either that the EP4 protein levels are low, or that

these receptors are functionally uncoupled from adenylyl cyclase. In contrast, PGE₂ stimulated accumulation of inositol phosphates. Pretreatment with the EP4 antagonist L161982 or the EP1 antagonist SC51322, had no effect on the PGE₂-induced phosphorylation of EGFR, ERK, or Akt, while the phosphorylation of these proteins were markedly inhibited by the FP antagonist AL8810. PGF_{2α}, which binds to FP receptors with high affinity, mimicked the effects of PGE₂. Together, these results suggest that in contrast to the normal rat hepatocytes, where the effect of PGE₂ seems to be mediated primarily through the EP3 receptor [37,52,54], the MH₁C₁ cells, which do not express EP3 receptors, respond to PGE₂



through FP receptors, Gq, and PLCβ. It is of interest that expression of EP3 receptors has been found to be suppressed or absent in colon cancer in vivo and in vitro, as compared to normal mucosa [55].

PLCβ can regulate cellular functions via two distinct pathways, involving DAG-mediated activation of PKC and InsP₃-induced release and elevation of cytosolic Ca²⁺, respectively. Our findings suggest that in the MH₁C₁ cells, the effect of PGE₂ was mediated through Ca²⁺, since it was not mimicked by TPA and not inhibited by a PKC blocker, while thapsigargin, which elevates intracellular Ca²⁺, mimicked the PGE₂ effect, inducing a gefitinib-sensitive phosphorylation of EGFR.

In other cells, both ligand-dependent and ligand-independent mechanisms have been found to mediate

EGFR transactivation [5]. Ligand-dependent mechanisms involve the release of EGFR agonists by cleavage and shedding of membrane-associated precursors by proteases of the ADAM family [2,49]. Ligand-independent mechanisms have been suggested to involve intracellular molecules including Src family kinases and Pyk2 [1,3,56,57]. Han et al. reported that in Hep3B cells, PGE₂ induced phosphorylation of the EGFR through EP1 receptors and an intracellular mechanism involving Src [57]. Itabashi et al. demonstrated that in some hepatocarcinoma cell lines EGFR transactivation triggered by angiotensin II stimulation was mediated through release of EGFR ligand by members of the ADAM family [58]. In the MH₁C₁ cells, we observed that Src inhibitors abolished PGE₂-stimulated phosphorylation of the EGFR, ERK, and Akt, but in contrast, only slightly affected the response to EGF, suggesting a role of Src in the transactivation in these cells. We also found evidence for the involvement of ligand shedding in the transactivation of EGFR after PGE₂ stimulation, since pretreatment of the cells with the metalloproteinase inhibitor GM6001 almost completely prevented PGE₂-induced, but not EGF-induced, phosphorylation of EGFR, Akt and ERK. GM6001 did not affect the effects of PGE₂ in the normal hepatocytes. The lack of transactivation in response to PGE₂ in these cells could be due to the low expression of metalloproteinases in hepatocytes as compared to hepatocarcinoma cells [59]. These results indicate that in the MH₁C₁ cells Src is involved in activating ADAMs rather than directly stimulating the EGFR, thus combining the two mechanisms. The involvement of both Src and ADAMs has been reported in normal gastrointestinal epithelial and colon cancer cell lines [60].

Several signalling pathways seem to be important in hepatocarcinomas [19], and there is evidence that both EGFR-mediated mechanisms and the COX/prostaglandin system may be involved in the pathobiology of these tumours [17,18,20,35,36]. The results of the present study suggest a functional interaction between the EGFR and the prostaglandins. It has been proposed that transactivation can explain the mitogenic effect of GPCR ligands in some cell systems [61] and that it represents a means of diversifying signalling in the cells, by linking the input from a large number of ligands stimulating GPCRs to the pleiotypic and potentially tumorigenic effects of the EGFR [62]. However, there seems to be great variation between cell types with respect to the different pathways involved in the signalling. We have recently shown that while neurotensin, a GPCR agonist, activates ERK and Akt in an EGFR-independent way in pancreatic cancer Panc-1 cells, as also found by others [63], and activates ERK and Akt via EGFR transactivation in the colon cancer cell line HT 29, neurotensin

uses both EGFR-dependent and -independent pathways in the colon cancer cell line HCT 116 [12].

In the present study we have shown that PGE₂ has different ways of stimulating the cells, acting by FP-mediated EGFR transactivation in the hepatocarcinoma cells, whereas the effect is mediated mainly via EP3 receptors without any involvement of the EGFR in the hepatocytes [37,52]. This is further evidence of the diversity of intracellular cross-talk and underscores the importance of investigating such mechanisms in order to better understand the signalling in cancer cells.

Conclusion

The results indicate that in MH₁C₁ cells, unlike normal hepatocytes, PGE₂ activates the MEK/ERK and PI3K/Akt pathways by transactivation of the EGFR, thus diversifying the GPCR-mediated signal. The data also suggest that the underlying mechanisms in these cells involve FP receptors, PLCβ, Ca²⁺, Src, and proteinase-mediated release of membrane-associated EGFR ligand(s).

Abbreviations

ADAM: A disintegrin and metalloproteinase; DAG: Diacylglycerol; EGF: Epidermal growth factor; EGFR: Epidermal growth factor receptor; ERK: Extracellular regulated kinase; EP: E prostaglandin; FP: F prostaglandin; GPCR: G protein-coupled receptor; InsP₃: Inositol trisphosphate; PG: Prostaglandin; PGE₂: Prostaglandin E₂; PGF_{2α}: Prostaglandin F_{2α}; PLCβ: Phospholipase C-β; PKC: Protein kinase C; RTK: Receptor tyrosine kinase; TPA: Tetradecanoylphorbol acetate; SERCA: Sarco/endoplasmic reticulum Ca²⁺-ATPase.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

IHT participated in the design of the study, carried out immunoblotting experiments and drafted the manuscript. KMM carried out immunoblotting experiments, inositol phosphate experiments and helped revise the manuscript. MA helped revise the manuscript. JØ carried out qRT-PCR experiment and helped revise the manuscript. OD conceived of the study, carried out DNA synthesis and helped revise the manuscript. TG conceived of the study and helped revise the manuscript. DS conceived of the study, participated in the design of the study, carried out cAMP and inositol phosphate experiments and helped revise the manuscript. TC conceived of the study, participated in the design of the study and helped revise the manuscript. All authors read and approved of the final manuscript.

Acknowledgements

The work was supported by the Norwegian Cancer Society. We thank Eva Østby and Ellen Johanne Johansen for excellent technical assistance.

Author details

¹Department of Pharmacology, Institute of Clinical Medicine, Faculty of Medicine, University of Oslo, P.O.Box 1057 Blindern, N-0316 Oslo, Norway.

²Department of Oncology, Oslo University Hospital, Oslo, Norway.

Received: 27 April 2012 Accepted: 1 September 2012

Published: 11 September 2012

References

1. Daub H, Weiss FU, Wallasch C, Ullrich A: **Role of transactivation of the EGF receptor in signalling by G-protein-coupled receptors.** *Nature* 1996, **379**(6565):557–560.
2. Prenzel N, Zwick E, Daub H, Leserer M, Abraham R, Wallasch C, Ullrich A: **EGF receptor transactivation by G-protein-coupled receptors requires**

- metalloproteinase cleavage of proHB-EGF. *Nature* 1999, **402**(6764):884–888.
3. Buchanan FG, Wang D, Bargiacchi F, DuBois RN: **Prostaglandin E2 regulates cell migration via the intracellular activation of the epidermal growth factor receptor.** *J Biol Chem* 2003, **278**(37):35451–35457.
4. Schafer B, Marg B, Gschwind A, Ullrich A: **Distinct ADAM metalloproteinases regulate G protein-coupled receptor-induced cell proliferation and survival.** *J Biol Chem* 2004, **279**(46):47929–47938.
5. Bholra NE, Grandis JR: **Crosstalk between G-protein-coupled receptors and epidermal growth factor receptor in cancer.** *Front Biosci* 2008, **13**:1857–1865.
6. Carpenter G, Cohen S: **Epidermal growth factor.** *J Biol Chem* 1990, **265**(14):7709–7712.
7. Jorissen RN, Walker F, Pouliot N, Garrett TP, Ward CW, Burgess AW: **Epidermal growth factor receptor: mechanisms of activation and signalling.** *Exp Cell Res* 2003, **284**(1):31–53.
8. Holbro T, Civenni G, Hynes NE: **The ErbB receptors and their role in cancer progression.** *Exp Cell Res* 2003, **284**(1):99–110.
9. Normanno N, De Luca A, Bianco C, Strizzi L, Mancino M, Maiello MR, Carotenuto A, De Feo G, Caponigro F, Salomon DS: **Epidermal growth factor receptor (EGFR) signaling in cancer.** *Gene* 2006, **366**(1):2–16.
10. Christoffersen T, Guren TK, Spindler KL, Dahl O, Lonning PE, Gjertsen BT: **Cancer therapy targeted at cellular signal transduction mechanisms: strategies, clinical results, and unresolved issues.** *Eur J Pharmacol* 2009, **625**(1–3):6–22.
11. Ciardiello F, Tortora G: **EGFR antagonists in cancer treatment.** *N Engl J Med* 2008, **358**(11):1160–1174.
12. Müller KM, Tveteraas IH, Aasrum M, Odegard J, Dawood M, Dajani O, Christoffersen T, Sandnes DL: **Role of protein kinase C and epidermal growth factor receptor signalling in growth stimulation by neurotensin in colon carcinoma cells.** *BMC Cancer* 2011, **11**:421.
13. Sand TE, Christoffersen T: **Temporal requirement for epidermal growth factor and insulin in the stimulation of hepatocyte DNA synthesis.** *J Cell Physiol* 1987, **131**(2):141–148.
14. Mead JE, Fausto N: **Transforming growth factor alpha may be a physiological regulator of liver regeneration by means of an autocrine mechanism.** *Proc Natl Acad Sci U S A* 1989, **86**(5):1558–1562.
15. Christoffersen T, Thoresen GH, Dajani OF, Melien Ø, Puren T, Refsnes M, Sandnes D: **Mechanisms of hepatocyte growth regulation by hormones and growth factor.** In *The hepatocyte review*. edn. Edited by AM BMAE. Dordrecht/Boston/London: Kluwer Academic Publishers; 2000:209–246.
16. Scheving LA, Stevenson MC, Taylormoore JM, Traxler P, Russell WE: **Integral role of the EGF receptor in HGF-mediated hepatocyte proliferation.** *Biochem Biophys Res Commun* 2002, **290**(1):197–203.
17. Farazi PA, DePinho RA: **Hepatocellular carcinoma pathogenesis: from genes to environment.** *Nat Rev Cancer* 2006, **6**(9):674–687.
18. Villanueva A, Newell P, Chiang DY, Friedman SL, Llovet JM: **Genomics and signaling pathways in hepatocellular carcinoma.** *Semin Liver Dis* 2007, **27**(1):55–76.
19. Whittaker S, Marais R, Zhu AX: **The role of signaling pathways in the development and treatment of hepatocellular carcinoma.** *Oncogene* 2010, **29**(36):4989–5005.
20. Berasain C, Castillo J, Prieto J, Avila MA: **New molecular targets for hepatocellular carcinoma: the ErbB1 signaling system.** *Liver Int* 2007, **27**(2):174–185.
21. Jhappan C, Stahle C, Harkins RN, Fausto N, Smith GH, Merlino GT: **TGF alpha overexpression in transgenic mice induces liver neoplasia and abnormal development of the mammary gland and pancreas.** *Cell* 1990, **61**(6):1137–1146.
22. Sandgren EP, Luetette NC, Qiu TH, Palmiter RD, Brinster RL, Lee DC: **Transforming growth factor alpha dramatically enhances oncogene-induced carcinogenesis in transgenic mouse pancreas and liver.** *Mol Cell Biol* 1993, **13**(1):320–330.
23. Ito Y, Takeda T, Sakon M, Tsujimoto M, Higashiyama S, Noda K, Miyoshi E, Monden M, Matsuura N: **Expression and clinical significance of erb-B receptor family in hepatocellular carcinoma.** *Br J Cancer* 2001, **84**(10):1377–1383.
24. Hopfner M, Sutter AP, Huether A, Schuppan D, Zeitz M, Scherubl H: **Targeting the epidermal growth factor receptor by gefitinib for treatment of hepatocellular carcinoma.** *J Hepatol* 2004, **41**(6):1008–1016.

25. Huether A, Hopfner M, Baradari V, Schuppan D, Scherubl H: **EGFR blockade by cetuximab alone or as combination therapy for growth control of hepatocellular cancer.** *Biochem Pharmacol* 2005, **70**(11):1568–1578.
26. Villanueva A, Llovet JM: **Targeted therapies for hepatocellular carcinoma.** *Gastroenterology* 2011, **140**(5):1410–1426.
27. Breyer RM, Bagdassarian CK, Myers SA, Breyer MD: **Prostanoid receptors: subtypes and signaling.** *Annu Rev Pharmacol Toxicol* 2001, **41**:661–690.
28. Wang D, Dubois RN: **Eicosanoids and cancer.** *Nat Rev Cancer* 2010, **10**(3):181–193.
29. Dubois RN, Abramson SB, Crofford L, Gupta RA, Simon LS, Van De Putte LB, Lipsky PE: **Cyclooxygenase in biology and disease.** *FASEB J* 1998, **12**(12):1063–1073.
30. Gupta RA, Dubois RN: **Colorectal cancer prevention and treatment by inhibition of cyclooxygenase-2.** *Nat Rev Cancer* 2001, **1**(1):11–21.
31. Zha S, Yegnasubramanian V, Nelson WG, Isaacs WB, De Marzo AM: **Cyclooxygenases in cancer: progress and perspective.** *Cancer Lett* 2004, **215**(1):1–20.
32. Andreis PG, Whitfield JF, Armato U: **Stimulation of DNA synthesis and mitosis of hepatocytes in primary cultures of neonatal rat liver by arachidonic acid and prostaglandins.** *Exp Cell Res* 1981, **134**(2):265–272.
33. Refsnes M, Dajani OF, Sandnes D, Thoresen GH, Rottingen JA, Iversen JG, Christoffersen T: **On the mechanisms of the growth-promoting effect of prostaglandins in hepatocytes: the relationship between stimulation of DNA synthesis and signaling mediated by adenyl cyclase and phosphoinositide-specific phospholipase C.** *J Cell Physiol* 1995, **164**(3):465–473.
34. Refsnes M, Thoresen GH, Dajani OF, Christoffersen T: **Stimulation of hepatocyte DNA synthesis by prostaglandin E2 and prostaglandin F2 alpha: additivity with the effect of norepinephrine, and synergism with epidermal growth factor.** *J Cell Physiol* 1994, **159**(1):35–40.
35. Koga H, Sakisaka S, Ohishi M, Kawaguchi T, Taniguchi E, Sasatomi K, Harada M, Kusaba T, Tanaka M, Kimura R, et al: **Expression of cyclooxygenase-2 in human hepatocellular carcinoma: relevance to tumor dedifferentiation.** *Hepatology* 1999, **29**(3):688–696.
36. Tang TC, Poon RT, Lau CP, Xie D, Fan ST: **Tumor cyclooxygenase-2 levels correlate with tumor invasiveness in human hepatocellular carcinoma.** *World J Gastroenterol* 2005, **11**(13):1896–1902.
37. Dajani OF, Meisdalen K, Guren TK, Aasrum M, Tveteraas IH, Lilleby P, Thoresen GH, Sandnes D, Christoffersen T: **Prostaglandin E2 upregulates EGF-stimulated signaling in mitogenic pathways involving Akt and ERK in hepatocytes.** *J Cell Physiol* 2008, **214**(2):371–380.
38. Nilssen LS, Odegard J, Thoresen GH, Molven A, Sandnes D, Christoffersen T: **G protein-coupled receptor agonist-stimulated expression of ATF3/LRF-1 and c-myc and comitogenic effects in hepatocytes do not require EGF receptor transactivation.** *J Cell Physiol* 2004, **201**(3):349–358.
39. Richardson UI, Tashjian AH Jr, Levine L: **Establishment of a clonal strain of hepatoma cells which secrete albumin.** *J Cell Biol* 1969, **40**(1):236–247.
40. Christoffersen T, Refsnes M, Bronstad GO, Ostby E, Huse J, Haffner F, Sand TE, Hunt NH, Sonne O: **Changes in hormone responsiveness and cyclic AMP metabolism in rat hepatocytes during primary culture and effects of supplementing the medium with insulin and dexamethasone.** *Eur J Biochem* 1984, **138**(2):217–226.
41. Bustin SA: **Absolute quantification of mRNA using real-time reverse transcription polymerase chain reaction assays.** *J Mol Endocrinol* 2000, **25**(2):169–193.
42. Skomedal T, Grynne B, Osnes JB, Sjetnan AE, Oye I: **A radioimmunoassay for cyclic AMP (cAMP) obtained by acetylation of both unlabeled and labeled (3 H-cAMP) ligand, or of unlabeled ligand only.** *Acta Pharmacol Toxicol (Copenh)* 1980, **46**(3):200–204.
43. Sugimoto Y, Narumiya S: **Prostaglandin E receptors.** *J Biol Chem* 2007, **282**(16):11613–11617.
44. Ji R, Chou CL, Xu W, Chen XB, Woodward DF, Regan JW: **EP1 prostanoid receptor coupling to G i/o up-regulates the expression of hypoxia-inducible factor-1 alpha through activation of a phosphoinositide-3 kinase signaling pathway.** *Mol Pharmacol* 2010, **77**(6):1025–1036.
45. Griffin BW, Klimko P, Crider JY, Sharif NA: **AL-8810: a novel prostaglandin F2 alpha analog with selective antagonist effects at the prostaglandin F2 alpha (FP) receptor.** *J Pharmacol Exp Ther* 1999, **290**(3):1278–1284.
46. Machwate M, Harada S, Leu CT, Seedor G, Labelle M, Gallant M, Hutchins S, Lachance N, Sawyer N, Slipetz D, et al: **Prostaglandin receptor EP(4) mediates the bone anabolic effects of PGE(2).** *Mol Pharmacol* 2001, **60**(1):36–41.
47. Santiskulvong C, Rozengurt E: **Protein kinase Calpha mediates feedback inhibition of EGF receptor transactivation induced by Gq-coupled receptor agonists.** *Cell Signal* 2007, **19**(6):1348–1357.
48. Thastrup O, Cullen PJ, Drobak BK, Hanley MR, Dawson AP: **Thapsigargin, a tumor promoter, discharges intracellular Ca2+ stores by specific inhibition of the endoplasmic reticulum Ca2(+)-ATPase.** *Proc Natl Acad Sci U S A* 1990, **87**(7):2466–2470.
49. Ohtsu H, Dempsey PJ, Eguchi S: **ADAMs as mediators of EGF receptor transactivation by G protein-coupled receptors.** *Am J Physiol Cell Physiol* 2006, **291**(1):C1–C10.
50. Santiskulvong C, Rozengurt E: **Galardin (GM 6001), a broad-spectrum matrix metalloproteinase inhibitor, blocks bombesin- and LPA-induced EGF receptor transactivation and DNA synthesis in rat-1 cells.** *Exp Cell Res* 2003, **290**(2):437–446.
51. Odegard J, Aasrum M, Tveteraas IH, Bharath SP, Sandnes D, Christoffersen T: **Role of ErbB2 in the prostaglandin E(2)-induced enhancement of the mitogenic response to epidermal growth factor in cultured hepatocytes.** *Biochem Biophys Res Commun* 2012, **421**(2):255–260.
52. Meisdalen K, Dajani OF, Christoffersen T, Sandnes D: **Prostaglandins enhance epidermal growth factor-induced DNA synthesis in hepatocytes by stimulation of E prostanoid 3 and F prostanoid receptors.** *J Pharmacol Exp Ther* 2007, **322**(3):1044–1050.
53. Bronstad GO, Gladhaug IP, Haffner F, Rugstad HE, Christoffersen T: **The regulation of cyclic AMP levels in cultured MH1C1 rat hepatoma cells and in solid tumours derived from MH1C1 cell inoculates.** *Anticancer Res* 1987, **7**(2):155–160.
54. Hashimoto N, Watanabe T, Ikeda Y, Yamada H, Taniguchi S, Mitsui H, Kurokawa K: **Prostaglandins induce proliferation of rat hepatocytes through a prostaglandin E2 receptor EP3 subtype.** *Am J Physiol* 1997, **272**(3 Pt 1):G597–G604.
55. Shoji Y, Takahashi M, Kitamura T, Watanabe K, Kawamori T, Maruyama T, Sugimoto Y, Negishi M, Narumiya S, Sugimura T, et al: **Downregulation of prostaglandin E receptor subtype EP3 during colon cancer development.** *Gut* 2004, **53**(8):1151–1158.
56. Andreev J, Galisteo ML, Kranenburg O, Logan SK, Chiu ES, Okigaki M, Cary LA, Moolenaar WH, Schlessinger J: **Src and Pyk2 mediate G-protein-coupled receptor activation of epidermal growth factor receptor (EGFR) but are not required for coupling to the mitogen-activated protein (MAP) kinase signaling cascade.** *J Biol Chem* 2001, **276**(23):20130–20135.
57. Han C, Michalopoulos GK, Wu T: **Prostaglandin E2 receptor EP1 transactivates EGFR/MET receptor tyrosine kinases and enhances invasiveness in human hepatocellular carcinoma cells.** *J Cell Physiol* 2006, **207**(1):261–270.
58. Itabashi H, Maesawa C, Oikawa H, Kotani K, Sakurai E, Kato K, Komatsu H, Nitta H, Kawamura H, Wakabayashi G, et al: **Angiotensin II and epidermal growth factor receptor cross-talk mediated by a disintegrin and metalloprotease accelerates tumor cell proliferation of hepatocellular carcinoma cell lines.** *Hepatology* 2008, **38**(6):601–613.
59. Caja L, Ortiz C, Bertran E, Murillo MM, Miro-Obradors MJ, Palacios E, Fabregat I: **Differential intracellular signalling induced by TGF-beta in rat adult hepatocytes and hepatoma cells: implications in liver carcinogenesis.** *Cell Signal* 2007, **19**(4):683–694.
60. Pai R, Soreghan B, Szabo IL, Pavelka M, Baatar D, Tarnawski AS: **Prostaglandin E2 transactivates EGF receptor: a novel mechanism for promoting colon cancer growth and gastrointestinal hypertrophy.** *Nat Med* 2002, **8**(3):289–293.
61. Daub H, Wallasch C, Lankenau A, Herrlich A, Ullrich A: **Signal characteristics of G protein-transactivated EGF receptor.** *EMBO J* 1997, **16**(23):7032–7044.
62. Fischer OM, Hart S, Gschwind A, Ullrich A: **EGFR signal transactivation in cancer cells.** *Biochem Soc Trans* 2003, **31**(Pt 6):1203–1208.
63. Kisfalvi K, Guha S, Rozengurt E: **Neurotensin and EGF induce synergistic stimulation of DNA synthesis by increasing the duration of ERK signaling in ductal pancreatic cancer cells.** *J Cell Physiol* 2005, **202**(3):880–890.

doi:10.1186/1756-9966-31-72

Cite this article as: Tveteraas et al.: Mechanisms involved in PGE₂-induced transactivation of the epidermal growth factor receptor in MH₁C₁ hepatocarcinoma cells. *Journal of Experimental & Clinical Cancer Research* 2012 **31**:72.