


Molecular Physics: An International Journal at the Interface Between Chemistry and Physics

[Select Language](#) ▼

[Volume 109, Issue 2, 2011](#)



ESR and optical study of Cu^{2+} -doped bis-(5,5'-diethylbarbiturato)bis picoline Zn(II)

DOI:

10.1080/00268976.2010.519350

[Ram Kripal](#)^{a*}, [Sanjay Misra](#)^a & [Indrajeet Mishra](#)^{ab}

pages 239-249

[Publishing models and article dates explained](#)

Received: 9 Feb 2010

Accepted: 14 Aug 2010

Published online: 02 Feb 2011

Article Views: 48

Abstract

ESR studies were conducted on Cu^{2+} -doped bis-(5,5'-diethylbarbiturato)bis picoline Zn(II). Two Cu^{2+} lattice sites, Cu^{2+} (I) and Cu^{2+} (II), were identified. These sites exhibit two sets of four hyperfine lines in all directions. The g factor and hyperfine splitting were calculated from ESR absorption spectra: $g_x = 2.0201 \pm 0.002$, $g_y = 2.0900 \pm 0.002$, $g_z = 2.1634 \pm 0.002$, $A_x = (30 \pm 2) \times 10^{-4} \text{ cm}^{-1}$, $A_y = (40 \pm 2) \times 10^{-4} \text{ cm}^{-1}$ and $A_z = (154 \pm 2) \times 10^{-4} \text{ cm}^{-1}$. It was found that Cu^{2+} enters the lattice substitutionally. The ground-state wavefunction of the Cu^{2+} ion in this lattice was determined from the spin

Hamiltonian constants obtained from the ESR studies. With the help of an optical absorption study, the nature of the bonding in the complex is also discussed.

- [View full text](#)
- [Download full text](#)

Keywords

- [ESR](#),
- [spin Hamiltonian](#),
- [absorption](#),
- [angular variation](#),
- [rhombic symmetry](#)

Related articles

*Local tax will be added as applicable

DOI:

10.1080/00268976.2010.519350

[Ram Kripal](#)^{a*}, [Sanjay Misra](#)^a & [Indrajeet Mishra](#)^{ab}

pages 239-249