



UNIVERSITI PUTRA MALAYSIA

**LARVAL FISH COMPOSITION, DISTRIBUTION AND DIETS IN THE
SEAGRASS-MANGROVE ECOSYSTEM OF GELANG PATAH, JOHOR,
MALAYSIA**

ROUSHON ARA

FP 2011 14

**LARVAL FISH COMPOSITION, DISTRIBUTION
AND DIETS IN THE SEAGRASS-MANGROVE
ECOSYSTEM OF GELANG PATAH, JOHOR,
MALAYSIA**



ROUSHON ARA


**DOCTOR OF PHILOSOPHY
UNIVERSITI PUTRA MALAYSIA**

2011

**LARVAL FISH COMPOSITION, DISTRIBUTION AND DIETS IN THE
SEAGRASS-MANGROVE ECOSYSTEM OF GELANG PATAH, JOHOR,
MALAYSIA**

By

ROUSHON ARA



**Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia,
in Fulfilment of the Requirements for the Degree of Doctor of Philosophy**

July 2011

DEDICATION

To my parents who always kept praying for me to achieve my goal

To my husband 'Dr. S. M. Nurul Amin' and son 'Md. Jahin Zawad', 'AL-Moid Amin'; daughter 'Tazri Amin' who have sacrificed so much for me during this study period



Abstract of thesis presented to the Senate of Universiti Putra Malaysia in fulfilment of the requirement for the degree of Doctor of Philosophy

LARVAL FISH COMPOSITION, DISTRIBUTION AND DIETS IN THE SEAGRASS-MANGROVE ECOSYSTEM OF GELANG PATAH, JOHOR, MALAYSIA

By

ROUSHON ARA

July 2011

Chairperson: Associate Professor Aziz Arshad, PhD

Faculty: Agriculture

Fish larval composition, spatio-temporal distribution, density, family richness, Shannon-Wiener index and feeding habits were determined by analyzing samples collected from the seagrass-mangrove ecosystem of Gelang Patah, Johor Strait, Peninsular Malaysia between October 2007 and September 2008. Five stations were selected namely Upper estuary (S1), middle estuary (S2) and lower estuary (S3) of Pendas River; seagrass beds (S4) and open seas (S5). Each sampling station was approximately 1 km apart from each other. Fish larvae were collected by subsurface towing of bongo net equipped with a flow meter. In total, 24 families of fish larvae belong to six orders were identified from seagrass-mangrove ecosystem of Gelang Patah, Johor Strait. Similarity matrix and cluster analysis revealed that there are six different families under the order Perciformes.

In total, 24 families were identified from the coastal waters of Gelang Patah, Johor, Peninsular Malaysia. Among them, 14 occurred in upper estuary, 17 in middle

estuary, 16 in lower estuary, 20 in seagrass beds and 16 in open sea. Overall five (Clupeidae, Blenniidae, Terapontidae, Gobiidae and Sillaginidae) were the most dominant in study areas. Shannon-Wiener index varied significantly within monsoon and intermonsoon seasons peaking in the months October-January and May-August. The highest density of larval fishes was recorded at seagrass station (S4) and the spatial variations in larval density were significant ($p < 0.05$) between seagrass and other four sampling sides. None of the diversity indices showed significant among-stations except only family richness was significantly ($p < 0.05$) higher in seagrass beds than upper estuary. Among 24 families, 18 families were found to have correlation either positively or negatively with the water parameters. The highest and significant regression coefficient was observed in Sillaginidae which indicated 72% abundance of Sillaginidae was influenced by the major water parameters and remaining 28% by other factors.

In total, 267 Blenniidae, 401 Clupeidae, 126 Gobiidae and 117 Terapontidae stomachs were examined during the study period. Analyses of prey in the stomachs identified 24 important items (%Rs > 0.05) belonging to six major groups: phytoplankton, zooplankton, algae, plant like matter, debris and unidentified materials. According to the Simple Resultant Index (%Rs), the predominant food item in the stomach of all four families (Blenniidae, Clupeidae, Gobiidae and Terapontidae) was phytoplankton ($> 60\%$). This was followed by zooplankton in Blenniidae (18.24%) and Clupeidae (8.60%). On the other hand, the second diet composition was plant like matter in Gobiidae (14.73%) and Terapontidae (8.02%). Among phytoplankton, *Dacytyloccopsis fascicularis* (26.31%) was large quantity in the stomachs of Blenniidae and this was followed by *Nitzschia baccata* (23.38%). Conversely, *Nitzschia* sp. was

the highest quantity (26.33%) in the stomachs of Clupeidae larvae and subsequently *Dacytyloccopsis fascicularis* (13.56%), *Biddulphia sinensis* (10.51%), *Rhizosolenia araturensis* (8.63%), *Lauderia borealis* (8.06%) and *Fragilaria intermedia* (3.47%). In Gobiidae larvae, Chromophyta (28.30%) was observed as the highest quantity and second highest component was *Nitzschia* sp. (15.97%) and then *Dacytyloccopsis fascicularis* (8.27%). Similarly, Chromophyta (29.12%) was the highest quantity in the stomachs of Terapontidae (Table 6.17). Second and third component was *Nitzschia* sp. (15.95%) and *Dacytyloccopsis fascicularis* (13.80%). Overall, two most dominant phytoplankton (*Dacytyloccopsis fascicularis* and *Nitzschia* sp.) was observed among the four larval families. It is revealed that various food items were found in the stomachs of Bleniidae, Clupeidae, Gobiidae and Terapontidae larvae and remarkable that phytoplankton were more than 60% in the diet composition. Therefore, all studied fish larval families in the study areas are exclusively herbivore.

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk ijazah Doktor Falsafah

**KOMPOSISI, TABURAN DAN DIET LARVA IKAN PADA EKOSISTEM
BAKAU-RUMPUT LAUT DI GELANG PATAH, JOHOR, MALAYSIA**

Oleh

ROUSHON ARA

Julai 2011

Chairperson: Profesor Madya Aziz Arshad, PhD

Fakulti: Pertanian

Komposisi larva ikan, taburan kawasan dan masa, kepadatan, kepelbagaian famili, Shannon-Wiener indeks dan habitat pemakanan ditentukan melalui analisis sampel yang diambil dari ekosistem bakau-rumput laut di Gelang Patah, Selat Johor, Semenanjung Malaysia dari Oktober 2007 hingga September 2008. Sebanyak lima stesen yang terpilih dinamakan sebagai 'bahagian atas muara' (S1), 'pertengahan muara' (S2), dan 'bahagian rendah muara' (S3) Sungai Pendas, hamparan rumput laut (S4) dan kawasan laut terbuka (S5). Jarak di antara setiap stesen adalah lebih kurang 1km. Larva ikan disampel di permukaan air dengan menggunakan jaring bongo yang dilengkapi dengan jangka aliran. Secara keseluruhan, 24 famili larva ikan dari enam order telah dikenalpasti dari ekosistem bakau-rumput laut di Gelang Patah, Selat Johor. Persamaan matriks dan analisis kluster menunjukkan terdapat enam famili berbeza di bawah order Perciformes.

Secara keseluruhan, 24 famili telah dikenalpasti dari kawasan perairan Gelang Patah, Selat Johor, Semenanjung Malaysia. Di antara 24 famili tersebut, 14 famili terdapat di kawasan 'bahagian atas muara', 17 di kawasan 'pertengahan muara', 16 di kawasan 'bahagian rendah muara', 20 di kawasan hamparan rumput laut dan 16 di kawasan laut terbuka. Clupeidae, Blenniidae, Terapontidae, Gobiidae and Sillaginidae merupakan famili yang paling dominan di kawasan kajian. Indeks Shannon-Wiener menunjukkan perbezaan yang signifikan di antara kemuncak musim monsun dan antara monsun yang berlaku pada bulan Oktober-Januari dan Mei-Ogos. Kepadatan larva ikan yang tertinggi direkodkan di kawasan hamparan rumput laut (S4) dan variasi taburan kawasan untuk kepadatan larva ikan adalah signifikan ($p < 0.05$) di antara kawasan rumput laut dengan 4 kawasan kajian yang lain. Indeks diversiti tidak menunjukkan perbezaan yang signifikan di antara stesen, kecuali kepelbagaian famili yang lebih tinggi secara signifikan ($p < 0.05$) di kawasan hamparan rumput laut berbanding bahagian atas muara. Di antara 24 famili, 18 famili telah didapati mempunyai korelasi sama ada positif atau negatif dengan parameter air. Regresi tertinggi dan pekali yang signifikan telah diperhatikan dalam Sillaginidae yang menunjukkan sebanyak 72% Sillaginidae dipengaruhi oleh parameter air utama dan baki 28% dipengaruhi oleh faktor lain.

Secara keseluruhan, struktur perut larva untuk 267 Blenniidae, 401 Clupeidae, 126 Gobiidae and 117 Terapontidae telah dianalisa sepanjang tempoh kajian. Analisa kandungan perut menunjukkan terdapat 24 item penting ($\%Rs > 0.05$) yang tergolong kepada enam kumpulan: fitoplankton, zooplankton, alga, bahan berbentuk tumbuhan, debris dan bahan yang tidak dikenalpasti. Berdasarkan Simple Resultant Index ($\%Rs$), makanan predomian di dalam perut kesemua 4 famili (Blenniidae, Clupeidae,

Gobiidae and Terapontidae) adalah fitoplankton (> 60%), diikuti oleh zooplankton pada Blenniidae (18.24%) dan Clupeidae (8.60%). Komposisi diet yang kedua di dalam Gobiidae dan Terapontidae merupakan bahan berbentuk tumbuhan iaitu sebanyak (14.73%) dan (8.02%). *Dacytyloccopsis fascicularis* merupakan fitoplankton yang mempunyai kuantiti terbesar di dalam perut Blenniidae iaitu sebanyak (26.31%) diikuti oleh *Nitzschia baccata* (23.38%). Manakala *Nitzschia* sp. mencatatkan kuantiti tertinggi (26.33%) di dalam perut larva Clupeidae dan ini diikuti oleh *Dacytyloccopsis fascicularis* (13.56%), *Biddulphia sinensis* (10.51%), *Rhizosolenia araturensis* (8.63%), *Lauderia borealis* (8.06%) dan *Fragilaria intermedia* (3.47%). Chromophyta mencatatkan kuantiti tertinggi (28.30%) pada larva Gobiidae manakala komponen kedua tertinggi pula adalah *Nitzschia* sp. (15.97%) dan *Dacytyloccopsis fascicularis* (8.27%). Chromophyta mencatatkan kuantiti tertinggi (29.12%) di dalam Terapontidae (Jadual 6.17). Komponen kedua dan ketiga tertinggi adalah merupakan *Nitzschia* sp. (15.95%) dan *Dacytyloccopsis fascicularis* (13.80%). Secara keseluruhan, dua fitoplankton yang paling dominan (*Dacytyloccopsis fascicularis* dan *Nitzschia* sp.) telah dikenalpasti di kalangan empat famili larva. Kajian menunjukkan terdapat pelbagai variasi makanan ditemui di dalam perut larva Blenniidae, Clupeidae, Gobiidae dan Terapontidae termasuklah fitoplankton yang telah mencatatkan lebih dari 60% daripada komposisi diet keseluruhan. Oleh yang demikian, dapat dirumuskan yang kesemua larva ikan yang dikaji di kawasan kajian merupakan herbivor.

ACKNOWLEDGEMENTS

All the admires and appreciation is for Allah, the Almighty, Beneficial and the most Merciful, who has enabled me to submit this thesis.

It is my pleasure to express my profound sense of gratitude and indebtedness to my respected research supervisor, Associate Professor Dr. Aziz Arshad, the chairman of my supervisory committee for his guidance, valuable collaboration and inspiration during the research period. Without his friendly and quality supervision, this work would not have come to complete. I am profound indebted to my co-supervisors Associate Professor Dr. Siti Khalijah Daud and Professor Dr. Mazlan Abdul Ghaffar for their advice, critical thought, thoroughness to this thesis and for the continue constructive discussions and suggestions.

I would like to thank to Ministry of Science, Technology and Innovation (MOSTI), Malaysia (Grant No. 05-01-04-SF0613) for providing financial support to carry out this research work. Special thanks to TWOWS (Third World Organization for Women in Science) for providing financial support through a fellowship during the study period. In addition, thanks go to Mr. Perumal Kuppan, Science Officer, Laboratory of Marine Science and Aquaculture, Institute of Bioscience, Universiti Putra Malaysia for the assistance during field sampling.

My special appreciation goes to my parents, husband, sons, elder brother and sisters for their unfailing support and encouragement for my higher study. My respect also goes to Dr. Jeff Leis, Australian Museum of Sydney for his valuable advice, suggestions and encouragement of the research on fish larvae.

I would like to express my gratitude and thanks to the officers, technicians, undergraduate and graduate students who helped me throughout this study. I would especially like to thank my friends and lab mates Latifa, Izyan, Hafiza, Laila, Amani, Mahvas, Oh and Leena who have contributed in ways too numerous to list. Last, but not least, I am thankful to all of my well-wishers whom have helped me in any form.

I certify that a Thesis Examination Committee has met on 7th July to conduct the final examination of Roushon Ara on her thesis entitled “Fish Larval Composition, Distribution and Diets in the Seagrass-Mangrove Ecosystem of Gelang Patah, Johor Strait, Peninsular Malaysia” in accordance with the Universities and University Colleges Act 1971 and the Constitution of the Universiti Putra Malaysia [P.U.(A) 106] 15 March 1998. The Committee recommends that the student be awarded the Doctor of Philosophy.

Members of the Examination Committee were as follow:

Siti Shapor binti Hj Siraj, PhD

Professor
Faculty of Agriculture
Universiti Putra Malaysia
(Chairman)

Annie Christianus, PhD

Associate Professor
Faculty of Agriculture
Universiti Putra Malaysia
(Internal Examiner)

Hishamuddin b Omar, PhD

Lecturer
Faculty of Agriculture
Universiti Putra Malaysia
(Internal Examiner)

Miguel Dino Fortes, PhD

Professor
Marine Science Institute
University of The Philippines
(External Examiner)

BUJANG KIM HUAT, PhD

Professor and Deputy Dean
School of Graduate Studies
Universiti Putra Malaysia

Date:

This thesis was submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfillment of the requirement for the degree of Doctor of Philosophy. The members of the Supervisory Committee were as follows:

Aziz Arshad, PhD

Associate Professor
Faculty of Agriculture
Universiti Putra Malaysia
(Chairman)

Siti Khalijah Daud, PhD

Associate Professor
Faculty of Science
Universiti Putra Malaysia
(Member)

Mazlan Abdul Ghaffar, PhD

Professor
Faculty of Science and Technology
Universiti Kebangsaan Malaysia
(Member)


HASANAH MOHD GHAZALI, PhD

Professor and Dean
School of Graduate Studies
Universiti Putra Malaysia

Date:

DECLARATION

I declare that the thesis is my original work except for quotations and citations which have been duly acknowledged. I also declare that it has not been previously, and is not concurrently, submitted for any other degree at Universiti Putra Malaysia or at any other institution.



ROUSHON ARA

Date: 7 July 2011

TABLE OF CONTENTS

	Page
DEDICATION	ii
ABSTRACT	iii
ABSTRAK	vi
ACKNOWLEDGEMENTS	ix
APPROVAL	xi
DECLARATION	xiii
LIST OF TABLES	xvii
LIST OF FIGURES	xx
LIST OF ABBREVIATIONS	xxiv
CHAPTER	
1 GENERAL INTRODUCTION	1
1.1 Background of the study	1
1.2 Statement of the problems	4
1.3 Objectives of the study	6
1.4 Organization of the study	6
2 LITERATURE REVIEW	8
2.1 The fish larvae	8
2.2 The egg stage	9
2.3 The yolk-sac stage	10
2.4 Larval fish taxonomy	11
2.5 The post-larvae	12
2.6 Pigmentation	13
2.7 Structural characteristics	17
2.8 Identification of post larvae	18
2.9 Habitat of fish larvae	21
2.10 Food and feeding habits of fish larvae	26
2.11 Fish larval abundance and distribution	29
3 GENERAL METHODOLOGY	33
3.1 Introduction	33
3.2 Methods	33
4 TAXONOMY OF LARVAL FISHES IN THE SEAGRASS-MANGROVE ECOSYSTEM OF GELANG PATAH, JOHOR, PENINSULAR MALAYSIA	38
4.1 Introduction	38
4.2 Materials and Methods	39
4.3 Results	40
4.3.1 Morphometric description	42

	4.3.2 Morphometric variation	85
	4.4 Discussion	86
	4.5 Conclusions	94
5	SPATIO-TEMPORAL DISTRIBUTION OF FISH LARVAE IN THE SEAGRASS-MANGROVE ECOSYSTEM OF GELANG PATAH, JOHOR, PENINSULAR MALAYSIA	95
	5.1 Introduction	95
	5.2 Materials and Methods	97
	5.3 Results	98
	5.3.1 Fish larval family composition in five sampling sites	98
	5.3.2 Monthly abundance of fish larvae in five sampling sites	105
	5.3.3 Diversity index of fish larvae in five sampling sites	110
	5.3.4 Size structure of fish larval family in five sampling sites	115
	5.4 Discussion	121
	5.5 Conclusions	127
6	INFLUENCE OF ENVIRONMENTAL VARIABLES ON FISH LARVAL ABUNDANCE	128
	6.1 Introduction	128
	6.2 Materials and Methods	129
	6.3 Results	130
	6.3.1 Environmental variables	130
	6.3.2 Relationship between fish larval abundance and environmental variables	137
	6.4 Discussion	140
	6.5 Conclusions	141
7	FOOD AND FEEDING HABITS OF FOUR MOST DOMINANT FAMILIES OF LARVAL FISHES IN THE SEAGRASS-MANGROVE ECOSYSTEM OF GELANG PATAH, JOHOR, PENINSULAR MALAYSIA	142
	7.1 Introduction	142
	7.2 Materials and Methods	145
	7.3 Results	148
	7.3.1 Food item compositions in four families of fish larvae	148
	7.3.2 Monthly variation of diet composition in four families of fish larvae	153
	7.4 Discussion	162
	7.5 Conclusions	167
8	GENERAL DISCUSSION, CONCLUSIONS AND RECOMMENDATIONS	168
	8.1 Discussion	168
	8.2 Conclusions	174
	8.3 Recommendations	175

REFERENCES	176
BIODATA OF STUDENT	188
LIST OF PUBLICATIONS	190



© COPYRIGHT UPM