



**UNIVERSITI PUTRA MALAYSIA**

**EXTRACTION, PARTIAL PURIFICATION AND CHARACTERISATION  
OF A PROTEASE EXTRACTED FROM KESINAI (Streblusasper Lour.)  
LEAVES, AND ITS POTENTIAL APPLICATION IN  
THE PRODUCTION OF CHEDDAR CHEESE**

**MYLVAGANAM PAGTHINATHAN**

**FSTM 2011 1**



**EXTRACTION, PARTIAL PURIFICATION AND CHARACTERISATION  
OF A PROTEASE EXTRACTED FROM KESINAI (*Streblus asper* Lour.)  
LEAVES, AND ITS POTENTIAL APPLICATION IN  
THE PRODUCTION OF CHEDDAR CHEESE**

**By**

**MYLVAGANAM PAGTHINATHAN**

**Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia,  
in Fulfilment of the Requirements for the Degree of Doctor of Philosophy.**

**January 2011**



## **DEDICATION**

This thesis is dedicated to

My father Kandia Mylvaganam

My mother Mylvaganam Paramesvari

My wife Pagthinathan Kumuthini



Abstract of thesis presented to the Senate of Universiti Putra Malaysia in fulfilment of the requirement for the degree of Doctor of Philosophy.

**EXTRACTION, PARTIAL PURIFICATION AND CHARACTERISATION  
OF A PROTEASE EXTRACTED FROM KESINAI (*Streblus asper* Lour.)  
LEAVES, AND ITS POTENTIAL APPLICATION IN  
THE PRODUCTION OF CHEDDAR CHEESE**

By

**MYLVAGANAM PAGTHINATHAN**

**January 2011**

**Chair : Hasanah Mohd Ghazali, PhD**

**Faculty : Food Science and Technology**

Calf-rennet is a conventional milk-clotting enzyme, which is most widely used coagulant in cheese-making all over the world. The world wide reduced supply of calf rennet and increase of cheese production have led to a systematic investigation of calf-rennet substitutes. Extracts of 'kesinai' (*Streblus asper*) leaves have been investigated as a source of enzymes to be used in cheese-making as an alternative source of calf rennet. The overall aim of this study was to evaluate the properties and application of *Streblus asper* leaf protease as a rennet substitute in cheese-making.

A milk-clotting protease was extracted from kesinai leaves using 100 mM Tris-HCl buffer (pH 7.4) and several other buffers. Purification was carried out using acetone precipitation, and ion-exchange and size-exclusion chromatography. Results obtained showed that 100 mM Tris-HCl buffer (pH 7.4) was found to be the most effective



extraction buffer. A higher yield of the kesinai protease was obtained using acetone precipitation as compared to ammonium sulphate precipitation. The ratio of cold acetone to crude extract of 1.25:1 was found to be the most suitable ratio for the partial purification of the protease. After the final purification step, the enzyme was partially purified 3.3 fold with a 42.3% of recovery.

The partially purified protease was characterised and it showed an optimum activity at 60 °C and pH 7.4. The enzyme was stable up to 70 °C for one hour, and residual activities at pH 6.0 to 10.0 were 72.4% and 70.2% of the optimum activity, respectively. The enzyme was found to have a higher temperature stability at -10 °C and 4 °C. It retained more than 98% of its activity 7 days after storage at both -10 and 4°C. The enzyme was inhibited by PMSF and trypsin inhibitor by 98% and 95.87% relative to the initial activity, respectively, suggesting the presence of serine residue at the active site.  $\text{Ca}^{2+}$  had a slight stimulating effect while,  $\text{Hg}^{2+}$ ,  $\text{Zn}^{2+}$  and  $\text{Pb}^{2+}$  had strong inhibitory effects on the enzyme activity. The partially purified enzyme appeared as a single band on SDS PAGE with an apparent molecular weight of 75.8 kDa.

The maximum coagulation activity was observed at pH 6.0 and at temperature 70 °C. The presence of  $\text{CaCl}_2$  up to 10 mM increased the milk-clotting activity while addition of NaCl decreased the milk-clotting activity. The milk-coagulation was also strengthened by increase enzyme concentration

Cheddar cheese was prepared using the acetone-precipitated enzyme and commercial enzyme (chymosin), and the physico-chemical, biochemical and sensory properties



were determined throughout two months of ripening. The yield of cheese produced using the commercial enzyme was 6% higher ( $p < 0.05$ ) than the yield obtained with kesinai enzymes. Cheeses made from kesinai enzyme and commercial enzyme were proximate compositionally alike except for moisture, protein contents and pH. The moisture content ( $42.67 \pm 0.47\%$ ) of cheese made with commercial enzyme was lower than the moisture ( $43.88 \pm 0.99\%$ ) of cheese made with kesinai enzyme, while the protein content ( $21.02 \pm 0.41\%$ ) and pH ( $4.79 \pm 0.03$ ) were ( $p < 0.05$ ) higher in cheese made with commercial enzyme than the protein content ( $18.87 \pm 0.32\%$ ) and pH ( $4.72 \pm 0.01$ ) obtained from cheese made from kesinai enzyme. In both cheeses, the moisture contents declined ( $p < 0.05$ ) with ripening, while the pH increased with ripening time. Non protein nitrogen (NPN) ( $0.04 \pm 0.00\%$ ) and non-casein nitrogen (NCN) ( $0.09 \pm 0.00\%$ ), were higher ( $p < 0.05$ ) in the whey obtained with kesinai enzyme compared to those (NPN and NCN were  $0.03 \pm 0.00\%$  and  $0.06 \pm 0.00\%$ , respectively), obtained with the commercial enzyme. Protein nitrogen (PN) ( $2.75 \pm 0.06\%$ ) and casein nitrogen (CN) ( $2.56\% \pm 0.05$ ) contents of cheese made with kesinai enzyme were ( $p < 0.05$ ) lower than those (CN and TN were  $2.93 \pm 0.11\%$  and  $3.29 \pm 0.12\%$ , respectively), made with the commercial enzyme. In general, all NPN and NCN values were found to increase ( $p < 0.05$ ) throughout the ripening period in both cheeses. Cheese made with the commercial enzyme exhibited a slightly higher level of free amino acids ( $27.06 \pm 0.64$  mg per g of cheese) compared to that of cheese made with kesinai enzyme ( $26.57 \pm 2.05$  mg per g of cheese). The principal component analysis (PCA) using a zNose® (Electronic Sensor Technology Co., Newbury Park, CA, USA) indicated the presence of seven volatile (aroma) compounds that are common in both cheeses. PCA analysis was performed to discriminate the cheeses in terms of aroma with their ripening time. Textural



characteristics such as hardness ( $811.8 \pm 36.6$  g for cheese made from kesinai enzyme and  $984.5 \pm 24.2$  g for cheese made from commercial enzyme), gumminess ( $622.8 \pm 39.2$  g for cheese made from kesinai enzyme and  $820.9 \pm 20.9$  g for cheese made from commercial enzyme), and chewiness ( $534.7 \pm 17.1$  g mm for cheese made from kesinai enzyme and  $735.3 \pm 14.6$  g mm for cheese made from commercial enzyme), increased in both types of cheese during the early stage of ripening and decrease gradually at end of ripening whereas springiness and cohesiveness showed similar changes in both cheeses during the ripened period.

It is concluded that kesinai protease could be used for the production of Cheddar cheese or it can be used together with another commercial enzyme in cheese production.



Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk ijazah Doktor Falsafah

**PENGEKSTRAKAN, PENULENAN SEPARA DAN PENCIRIAN PROTEASE  
DARIPADA DAUN KESINAI (*Streblus asper* Lour.), DAN POTENSI  
PENGUNAANNYA DALAM PENGELUARAN KEJU CEDAR**

Oleh

**MYLVAGANAM PAGTHINATHAN**

**Januari 2011**

**Pengerusi : Hasanah Mohd Ghazali, PhD**

**Fakulti : Sains dan Teknologi Makanan**

'Calf-rennet' ialah enzim pengkoagulasi susu konvensional, yang digunakan secara meluas sebagai koagulan dalam pembuatan keju di seluruh dunia. Kekurangan bekalan 'calf-rennet' di seluruh dunia dan peningkatan penghasilan keju telah membawa kepada kajian sistematik pengganti 'calf-rennet'. Ekstrak daun 'kesinai' (*Streblus asper*) telah dikenalpasti sebagai sumber enzim yang digunakan dalam penghasilan keju sebagai sumber alternatif 'calf-rennet'. Objektif keseluruhan kajian ini ialah untuk menilai kesesuaian protease *Streblus asper* sebagai pengganti rennet dalam pembuatan keju.

Protease pengkoagulasi-susu telah diekstrak dari daun kesinai menggunakan 100 mM penimbal Tris-HCl (pH 7.4) dan beberapa larutan penimbal lain. Penulenan telah dijalankan menggunakan pemendakan aseton, kromatografi penukaran ion dan kromatografi 'size-exclusion'. Keputusan menunjukkan bahawa 100 mM penimbal Tris-HCl (pH 7.4) telah didapati menjadi penimbal pengekstrakan paling berkesan.





Hasil protease kesinai yang lebih tinggi telah diperoleh melalui pemendakan aseton berbanding pemendakan ammonium sulfat. Nisbah aseton sejuk dan ekstrak mentah iaitu 1.25:1 telah didapati menjadi nisbah yang paling sesuai untuk penulenan separa protease. Selepas langkah terakhir penulenan, enzim ditulenan dengan 3.3 lipatan penulenan dan menghasilkan 42.3 % perolehan.

Enzim separa tulen telah dicirikan dan ia mempamerkan suhu optimum 60°C dan pH 7.4. Enzim stabil sehingga 70°C selama satu jam, dan aktiviti residual pada pH 6.0 dan 10.0 dengan aktiviti optimum 72.4% dan 70.2%, masing-masing. Enzim telah didapati mempunyai kestabilan suhu yang tinggi pada -10°C dan 4°C. Ia mengekalkan lebih dari 98% aktivitinya selepas 7 hari penyimpanan pada -10°C dan 4°C. Enzim telah direncat oleh PMSF dan perencat tripsin dengan 98% dan 95.87% relatif terhadap aktiviti permulaan, masing-masing, mencadangkan kehadiran residu serin pada tapak aktif.  $\text{Ca}^{2+}$  mempunyai kesan rangsangan lemah manakala,  $\text{Hg}^{2+}$ ,  $\text{Zn}^{2+}$  dan  $\text{Pb}^{2+}$  mempunyai kesan rencatan kuat pada aktiviti enzim. Enzim separa tulen muncul sebagai satu jalur tunggal pada SDS PAGE dengan berat molekul ketara 75.8 kDa.

Ciri-ciri pengkoagulasi susu bagi kesinai protease mentah, protease separa tulen dan rennet komersial telah dikaji. Aktiviti koagulasi sangat bergantung kepada pH dan suhu susu, dan aktiviti koagulasi maksimum telah didapati pada pH 6 dan 70°C. Kehadiran  $\text{CaCl}_2$  sehingga 10 mM meningkatkan aktiviti pengkoagulasi-susu manakala penambahan  $\text{NaCl}$  menurunkan aktiviti pengkoagulasi-susu. Pengkoagulasian susu juga dipertingkatkan dengan penambahan kepekatan enzim.

Keju cedar telah disediakan dengan menggunakan enzim hasil pemendakan aseton dan enzim komersial (chymosin), seterusnya ciri-ciri fiziko-kimia, biokimia dan sensori telah ditentukan sepanjang dua bulan tempoh pematangan keju cedar. Hasil perolehan keju yang disediakan dengan menggunakan enzim komersial adalah 6% lebih tinggi ( $p < 0.05$ ) berbanding keju yang dihasil dengan menggunakan enzim kesinai. Komposisi bagi kedua-dua keju ini adalah sama kecuali kelembapan, kandungan protein dan pH. Kandungan kelembapan ( $42.67 \pm 0.47\%$ ) keju yang dibuat menggunakan enzim komersial adalah lebih rendah daripada kandungan kelembapan ( $43.88 \pm 0.99\%$ ) keju dibuat menggunakan enzim kesinai, manakala kandungan protein ( $21.02 \pm 0.41\%$ ) dan pH ( $4.79 \pm 0.03$ ) adalah lebih tinggi dalam keju yang dibuat menggunakan enzim komersial berbanding kandungan protein ( $18.87 \pm 0.32\%$ ) dan pH ( $4.72 \pm 0.01$ ) diperolehi dalam keju yang dibuat daripada enzim kesinai. Pada kedua-dua keju, kandungan kelembapan adalah berkurangan ( $p < 0.05$ ) dengan masa pematangan, manakala nilai pH pula bertambah dengan masa pematangan. Kandungan nitrogen bukan-protein (NPN) ( $0.04 \pm 0.001\%$ ) dan nitrogen bukan-kasein (NCN) ( $0.09 \pm 0.001\%$ ), adalah lebih tinggi ( $p < 0.05$ ) dalam 'whey' yang diperolehi dengan enzim kesinai berbanding dengan enzim komersial (NPN dan NCN ialah  $0.03 \pm 0.002\%$  dan  $0.06 \pm 0.002\%$ , masing-masing). Kandungan nitrogen protein (PN) ( $2.75 \pm 0.06\%$ ) dan nitrogen kasein (CN) ( $2.56\% \pm 0.05$ ) dalam keju yang dibuat menggunakan enzim kesinai adalah ( $p < 0.05$ ) lebih rendah daripada enzim komersial (CN dan TN ialah  $2.93 \pm 0.11\%$  dan  $3.29 \pm 0.12\%$ , masing-masing). Secara keseluruhannya, nilai NPN dan NCN meningkat ( $p < 0.05$ ) sepanjang tempoh pematangan dalam kedua-dua keju. Keju yang dibuat menggunakan enzim komersial menunjukkan tahap asid amino bebas ( $27.06 \pm 0.64$  mg per g keju) yang tinggi sedikit berbanding dengan keju dibuat menggunakan

enzim kesinai ( $26.57 \pm 2.05$  mg per g keju). Perubahan pada kandungan asid lemak bebas adalah sama bagi kedua-dua keju tersebut semasa tempoh pematangan. Analisis komponen utama (PCA) menggunakan zNose® (Electronic Sensor Technology Co, Newbury Park, CA, USA) menunjukkan kehadiran tujuh sebatian meruap (aroma) yang biasa dijumpai dalam kedua-dua keju tersebut. Analisis PCA dijalankan bertujuan untuk membezakan aroma bagi kedua-dua keju semasa pematangan berlaku. Ciri-ciri tekstur seperti kekerasan ( $811.8 \pm 36.6$  g untuk keju dibuat daripada enzim kesinai dan  $984.5 \pm 24.2$  g untuk keju dibuat daripada enzim komersial), kebergetahan ( $622.8 \pm 39.2$  g untuk keju dibuat daripada enzim kesinai dan  $820.9 \pm 20.9$  g untuk keju dibuat daripada enzim komersial), dan kekenyalan ( $534.7 \pm 17.1$  g mm untuk keju dibuat daripada enzim kesinai dan  $735.3 \pm 14.6$  g mm untuk keju dibuat daripada enzim komersial) bertambah dalam kedua-dua jenis keju semasa peringkat awal pematangan dan berkurang secara beransur-ansur pada peringkat akhir pematangan, manakala keanjalan dan kekohesifan menunjukkan perubahan yang sama dalam kedua-dua keju semasa tempoh matang.

## ACKNOWLEDGEMENTS

Praise and thanks be first to the unseen power that made me exist in the universe along with my teachers, relative and friends without whom this thesis would have not come to a reality.

I am indebted to my supervisor, Professor Dr. Hasanah Mohd Ghazali, Faculty of Food Science and Technology, for her invaluable guidance, supervision, advises encouragement and constructive criticisms throughout the study.

I am also very grateful to Professor Dr. Mohd Yazid Abd Manap, a member of my supervisory committee, for his valuable discussions and help. He devoted much of his time to guide me during my work with patience giving enormous encouragement.

I am also thankful to Associate Professor Dr. Foo Hooi Ling, another member of supervisory committee, for her guidance, supervision, encouragement, and support in all steps of this study.

I would also like to thank Improving Relevance and Quality of Undergraduate Education (IRQUE) project, Government of Sri Lanka for the financial support.

I extend my thanks to all staff and colleagues at the Faculty of Food Science and Technology, Universiti Putra Malaysia, Malaysia.

Finally, I am grateful to my wife, my parents, my sister and my brother for their endless support and love during this thesis as in all stages of my life.



I certify that a Thesis Examination committee has met on 13<sup>th</sup> of January, 2011 to conduct the final examination of Mylvaganam Pagthinathan on his thesis entitled “Extraction, partial purification and characterisation of a protease extracted from kesinai (*streblus asper*) leaves, and its potential application in the development of Cheddar cheese ” in accordance with Universities and University Colleges Act 1971 and the Constitution of the Universiti Putra Malaysia [P.U.(A) 106] 15 March 1998. The Committee recommends that the student be awarded the Doctor of Philosophy.

Members of the Thesis Examination Committee were as follows:

Md Zaidul Islam Sarker, PhD  
Associated Professor,  
Faculty of Food Science and Technology  
Universiti Putra Malaysia.  
(Chairman)

Abdul Karim Sabo Mohamed, PhD  
Associated Professor,  
Faculty of Food Science and Technology  
Universiti Putra Malaysia.  
(Internal Examiner)

Seyed Hamed Mirhosseini, PhD  
Lecturer,  
Faculty of Food Science and Technology  
Universiti Putra Malaysia.  
(Internal Examiner)

Athanasios A. Koutinas, PhD  
Professor,  
Department of Chemistry  
University of Patras  
Patras, Greece.  
(External Examiner)

---

BUJANG KIM HUAT, PhD  
Professor and Deputy Dean  
School of Graduate Studies  
Universiti Putra Malaysia

Date :



The thesis was submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfilment of the requirement for the degree of Doctor of Philosophy. The members of Supervisory Committee were as follows:

**Hasanah Mohd Ghazali, PhD**  
Professor,  
Faculty of Food Science and Technology  
Universiti Putra Malaysia.  
(Chairman)

**Mohd Yazid Abd Manap, PhD**  
Professor,  
Faculty of Food Science and Technology  
Universiti Putra Malaysia.  
(Member)

**Foo Hooi Ling, PhD**  
Associated Professor,  
Faculty of Biotechnology and Biomolecular Sciences  
Universiti Putra Malaysia.  
(Member)

---

**HASANAH MOHD GHAZALI, PhD**  
Professor and Dean  
School of Graduate Studies  
Universiti Putra Malaysia

Date :



## **DECLARATION**

I declare that the thesis is my original work except for quotations and citations which have been duly acknowledged. I also declare that it has not been previously, and is not concurrently, submitted for any other degree at Universiti Putra Malaysia or at any other institution.

---

**MYLVAGANAM PAGTHINATHAN**

Date: 13<sup>th</sup> January 2011

## TABLE OF CONTENTS

	<b>Page</b>
<b>ABSTRACT</b>	III
<b>ABSTRAK</b>	VII
<b>ACKNOWLEDGEMENT</b>	XI
<b>APPROVAL</b>	XII
<b>DECLARATION</b>	XIV
<b>LIST OF TABLES</b>	XIX
<b>LIST OF FIGURES</b>	XXI
<b>LIST OF ABBREVIATIONS</b>	XXIV
<b>CHAPTER</b>	
<b>1 INTRODUCTION</b>	<b>1</b>
<b>2 LITERATURE REVIEW</b>	<b>5</b>
2.1 Milk	5
2.1.1 Casein	5
2.1.2 Structure of casein micelles	6
2.2 Milk Coagulation	11
2.2.1 Enzymatic coagulation	12
2.2.2 Factors affecting coagulation	15
2.3 Cheese	17
2.3.1 Ripening of Cheese	18
2.3.2 Microbial ripening	30
2.3.3 Accelerating of cheese ripening	30
2.3.4 Texture of cheese	31
2.4 Milk Coagulation Enzymes.	31
2.4.1 Animal rennet	33
2.4.2 Microbial rennet	34
2.4.3 Recombinant rennet	35
2.4.4 Plant rennet	35
2.5 Proteases	41
2.5.1 Classification of proteases	42
2.6 Protein Purification	48
2.6.1 Purification techniques	48
2.6.2 Evidence of purity	52
<b>3 EXTRACTION AND PURIFICATION OF PROTEASE FROM KESINAI (<i>Streblus asper</i>) LEAVES</b>	<b>53</b>
3.1 Introduction	53
3.2 Materials and Methods	55
3.2.1 Sample collection	55
3.2.2 Materials	56
3.2.3 Preparation of crude kesinai extract	56
3.2.4 Analysis of protease activity in crude kesinai extract	57
	xv





3.2.5	Protease activity assay	59
3.2.6	Milk-clotting activity assay	60
3.2.7	Protein determination	60
3.2.8	Partial purification of kesinai protease	61
3.2.9	Statistical analysis	64
3.3	Results and Discussion	65
3.3.1	Preparation of crude enzyme extract	65
3.3.2	Proteolytic activity of crude extract	72
3.3.3	Partial purification of kesinai protease	77
3.4	Conclusions	88
<b>4</b>	<b>CHARACTERISATION OF PARTIALLY PURIFIED PROTEASE FROM KESINAI (<i>Streblus asper</i>) LEAVES</b>	<b>91</b>
4.1	Introduction	91
4.2	Materials and Methods	93
4.2.1	Materials	93
4.2.2	Enzyme preparation	94
4.2.3	Determination of protease activity	94
4.2.4	Milk-clotting activity assay	94
4.2.5	Protein determination	95
4.2.6	Enzyme concentration on kesinai protease activity	95
4.2.7	Determination of optimum pH and pH stability	95
4.2.8	Determination of optimum temperature and temperature stability	96
4.2.9	Determination of $K_m$ and $V_{max}$ values	97
4.2.10	Determination of substrate specificity	98
4.2.11	Effect of protease inhibitors on kesinai protease activity	99
4.2.12	Effect of various metal cations on kesinai protease activity	100
4.2.13	Determination of native molecular mass by gel- filtration chromatography	101
4.2.14	Determination of purity using native polyacrylamide gel electrophoresis (NATIVE-PAGE)	101
4.2.15	Determination of molecular mass sodium dodecyl sulphate polyacrylamide gel electrophoresis (SDS-PAGE)	102
4.2.16	Statistical analysis	103
4.3	Results and Discussion	103
4.3.1	Partially purified enzyme concentration on kesinai protease activity	104
4.3.2	Optimum pH of partially purified kesinai protease	105
4.3.3	pH Stability of partially purified kesinai protease	106
4.3.4	Optimum temperature of partially purified kesinai protease	108
4.3.5	Temperature stability of partially purified kesinai protease	109
4.3.6	$K_m$ and $V_{max}$ values of partially purified kesinai	114



	protease	
	4.3.7 Substrate specificity of partially purified kesinai protease	115
	4.3.8 Effect of protease inhibitors on kesinai protease activity	116
	4.3.9 Effect of metal cations on kesinai protease activity	118
	4.3.10 Molecular mass of partially purified kesinai protease	121
	4.3.11 Milk-clotting activity of partially purified kesinai protease	125
	4.4. Conclusions	126
<b>5</b>	<b>COMPARISON OF MILK-CLOTTING PROPERTIES OF CRUDE AND ACETONE-PRECIPIATED KESINAI PROTEASE WITH COMMERCIAL RENNET</b>	<b>127</b>
	5.1 Introduction	127
	5.2 Materials and Methods	128
	5.2.1 Materials	129
	5.2.2 Preparation of enzyme	129
	5.2.3 Milk-clotting activity assay	130
	5.2.4 Protease activity assay	130
	5.2.5 Protein determination	130
	5.2.6 Effect of enzyme concentration on milk-clotting activity	131
	5.2.7 Effect of temperature on milk-clotting activity	131
	5.2.8 Effect of milk pH on milk-clotting activity	132
	5.2.9 Effect of calcium chloride concentration on milk-clotting activity	132
	5.2.10 Effect of sodium chloride concentration on milk-clotting activity	132
	5.2.11 Syneresis test	133
	5.2.12 Determination of coagulum strength	134
	5.2.13 Statistical analysis	136
	5.3 Results and Discussion	137
	5.3.1 Effect of pH of extraction buffer on milk-clotting activity	137
	5.3.2 Enzyme concentration on milk-clotting activity	138
	5.3.3 Effect of milk temperature on milk-clotting activity	141
	5.3.4 Effect of milk pH on milk-clotting activity	143
	5.3.5 Effect of calcium chloride concentration on milk-clotting activity	146
	5.3.6 Effect of Sodium chloride concentration on milk-clotting activity	148
	5.3.7 Syneresis of milk-coagulum	149
	5.3.8 Assessment of milk-clotting time and coagulum strength of kesinai protease mixture with pepsin	151
	5.4 Conclusions	154



6	<b>APPLICATION OF KESINAI (<i>Stebulus asper</i>) LEAVE PROTEASE EXTRACT IN CHEDDAR CHEESE MAKING</b>	156
	6.1 Introduction	156
	6.2 Materials and Methods	158
	6.2.1 Materials	158
	6.2.2 Production of Cheddar cheese	159
	6.2.3 Sampling of cheese for analysis	163
	6.2.4 Proximate analysis	163
	6.2.5 Determination of nitrogen contents in cheese and whey	163
	6.2.6 pH determination	164
	6.2.7 Free Amino acid composition	164
	6.2.8 Fatty acid composition	166
	6.2.9 Aroma profile	167
	6.2.10 Texture profile analysis	168
	6.2.11 Statistical analysis	169
	6.3 Results and Discussion	170
	6.3.1 Chemical composition of cheese	170
	6.3.2 Nitrogen fractions of cheese and whey	174
	6.3.3 Free Amino acid content of cheese	180
	6.3.4 Fatty acid composition of cheese	182
	6.3.5 Aroma profile	184
	6.3.6 Texture profile	195
	6.4 Conclusions	200
7	<b>SUMMARY, GENERAL CONCLUSIONS AND RECOMMENDATIONS FOR FUTURE RESEARCH</b>	201
	7.1 Summary and General Conclusions	201
	7.2 Recommendations for Future Research	206
	<b>REFERENCES</b>	207
	<b>APPENDICES</b>	232
	<b>BIODATE OF STUDENT</b>	239
	<b>LIST OF PUBLICATIONS</b>	240

