

Malaysian Journal of Mathematical Science 7(1): 79-95 (2013)

Application of Homotopy Perturbation Sumudu Transform Method for Solving Heat and Wave-Like Equations

¹Jagdev Singh, ²Devendra Kumar and ³Adem Kilicman

¹*Department of Mathematics, JaganNath University,
Village- Rampura, Tehsil- Chaksu, Jaipur-303901, Rajasthan, India*

²*Department of Mathematics,
JaganNath Gupta Institute of Engineering and Technology,
Jaipur-302022, Rajasthan, India*

³*Department of Mathematics and Institute for Mathematical Research,
Universiti Putra Malaysia, 43400 UPM, Serdang, Selangor, Malaysia*

*E-mail: jagdevsinghrathore@gmail.com, devendra.maths@gmail.com
and akilicman@science.upm.edu.my*

ABSTRACT

In this paper, we use the homotopy perturbation sumudu transform method (HPSTM) to solve heat and wave-like equations. The proposed scheme finds the solution without any discretization or restrictive assumptions and avoids the round-off errors. Several examples are given to verify the reliability and efficiency of the method. The fact that the proposed technique solves nonlinear problems without using Adomian's polynomials can be considered as a clear advantage of this algorithm over the decomposition method.

Keywords: homotopy perturbation sumudu transform method, sumudu transform, heat and wave-like equations, He's Polynomials.

1. INTRODUCTION

The heat and wave-like models are the integral part of applied sciences and arise in various physical phenomena. Several techniques including spectral, characteristic, modified variational iteration, Adomian's decomposition method and He's polynomials have been used for solving these problems (see Noor and Mohyud-Din (2008), Wazwaz and Gorguis (2004), Wilcox (1970) and Mohyud-Din (2009)) and references therein. Most of these methods have their inbuilt deficiencies like the calculation of Adomian's polynomials, the Lagrange multiplier, divergent results and huge computational work. He (1999, 2003, 2004) and references therein developed the homotopy perturbation method (HPM) by merging the standard homotopy and perturbation for solving various physical problems.

It is worth mentioning that the HPM is applied without any discretization, restrictive assumption or transformation and is free from round off errors. The homotopy perturbation method is also combined with the well-known Laplace transformation method (Madani and Fathizadeh (2010) and Khan and Wu (2011)) to produce a highly effective technique for handling many nonlinear problems. Very recently Singh, Kumar and Sushila (2011) have introduced, a new technique called homotopy perturbation sumudu transform method (HPSTM) for solving nonlinear equations. It is worth mentioning that HPSTM is an elegant combination of the sumudu transform method, the homotopy perturbation method and He's polynomials and is mainly due to Ghorbani and Saberi-Nadjafi (2007) and Ghorbani (2009).

The use of He's polynomials in the nonlinear term was first introduced by Ghorbani and Saberi-Nadjafi (2007) and Ghorbani (2009). HPSTM provides the solution in a rapid convergent series which may lead to the solution in a closed form. The advantage of this method is its capability of combining two powerful methods for obtaining exact and approximate solutions for nonlinear equations. Inspired and motivated by the ongoing research in this area, we apply HPSTM for solving the heat and wave-like equations in the present article. Several examples are given to verify the reliability and efficiency of the technique.

2. SUMUDU TRANSFORM

In early 90's, Watugala (1998) introduced a new integral transform, named the sumudu transform and applied it to the solution of ordinary differential equation in control engineering problems. The sumudu transform is defined over the set of functions

$$A = \{f(t) \mid \exists M, \tau_1, \tau_2 > 0, |f(t)| < M e^{t/\tau_j}, \text{ if } t \in (-1)^j \times [0, \infty)\}$$

by the following formula

$$\bar{f}(u) = S[f(t)] = \int_0^\infty f(ut) e^{-t} dt, u \in (-\tau_1, \tau_2). \quad (1)$$

Some of the properties were established by Weerakoon in Kilicman *et al.* (2011) and Weerakoon (1994). In Asiru (2004), further fundamental properties of this transform were also established. Similarly, this transform was applied to the one-dimensional neutron transport equation in Kadem

(2005). In fact it was shown that there is strong relationship between Sumudu and other integral transform (see Kilicman and Eltayeb (2010)). In particular the relation between Sumudu transform and Laplace transforms was proved in Kilicman and Eltayeb (2010).

Further, in Eltayeb *et al.* (2010), the Sumudu transform was extended to the distributions and some of their properties were also studied in Kilicman *et al.* (2010). Recently, this transform is applied to solve the system of differential equations (see Kilicman *et al.* (2010)).

Note that a very interesting fact about Sumudu transform is that the original function and its Sumudu transform have the same Taylor coefficients except the factor n (see Zhang (2007)).

Thus if $f(t) = \sum_{n=0}^{\infty} a_n t^n$ then $F(u) = \sum_{n=0}^{\infty} n! a_n u^n$, see Kilicman and

Eltayeb (2010). Similarly, the Sumudu transform sends combinations, $C(m, n)$, into permutations, $P(m, n)$ and hence it will be useful in the discrete systems.

3. HOMOTOPY PERTURBATION SUMUDU TRANSFORM METHOD (HPSTM)

To illustrate the basic idea of this method, we consider a general nonlinear non-homogenous partial differential equation with the initial conditions of the form

$$DU(x,t) + RU(x,t) + NU(x,t) = g(x,t), \quad (2)$$

$$U(x,0) = h(x), \quad U_t(x,0) = f(x),$$

where D is the second order linear differential operator $D = \partial^2 / \partial t^2$, R is the linear differential operator of less order than D , N represents the general nonlinear differential operator and $g(x,t)$ is the source term.

Taking the sumudu transform on both sides of Equation (2), we get

$$S[DU(x,t)] + S[RU(x,t)] + S[NU(x,t)] = S[g(x,t)]. \quad (3)$$

Using the differentiation property of the sumudu transform and above initial conditions, we have

$$S[U(x,t)] = u^2 S[g(x,t)] + h(x) + uf(x) - u^2 S[RU(x,t)] + S[NU(x,t)]. \quad (4)$$

Now, applying the inverse sumudu transform on both sides of Equation (4), we get

$$U(x,t) = G(x,t) - S^{-1} \left[u^2 S[RU(x,t) + NU(x,t)] \right], \quad (5)$$

where $G(x,t)$ represents the term arising from the source term and the prescribed initial conditions. Now, we apply the homotopy perturbation method

$$U(x,t) = \sum_{n=0}^{\infty} p^n U_n(x,t) \quad (6)$$

and the nonlinear term can be decomposed as

$$NU(x,t) = \sum_{n=0}^{\infty} p^n H_n(U), \quad (7)$$

for some He's polynomials $H_n(U)$ (see Ghorbani (2009) and Mohyud-Din *et al.* (2009)) that are given by

$$H_n(U_0, U_1, \dots, U_n) = \frac{1}{n!} \frac{\partial^n}{\partial p^n} \left[N \left(\sum_{i=0}^{\infty} p^i U_i \right) \right]_{p=0}, \quad n = 0, 1, 2, 3, \dots \quad (8)$$

Substituting Equations (6) and (7) in Equation (5), we get

$$\begin{aligned} & \sum_{n=0}^{\infty} p^n U_n(x,t) \\ & = G(x,t) - p \left(S^{-1} \left[u^2 S \left[R \sum_{n=0}^{\infty} p^n U_n(x,t) + \sum_{n=0}^{\infty} p^n H_n(U) \right] \right] \right), \quad (9) \end{aligned}$$

which is the coupling of the sumudu transform and the homotopy perturbation method using He's polynomials.

By comparing the coefficient of like powers of p , the following approximations are obtained

$$\begin{aligned}
 p^0 : U_0(x,t) &= G(x,t), \\
 p^1 : U_1(x,t) &= -S^{-1} \left[u^2 S [RU_0(x,t) + H_0(U)] \right], \\
 p^2 : U_2(x,t) &= -S^{-1} \left[u^2 S [RU_1(x,t) + H_1(U)] \right], \\
 p^3 : U_3(x,t) &= -S^{-1} \left[u^2 S [RU_2(x,t) + H_2(U)] \right], \\
 &\vdots
 \end{aligned} \tag{10}$$

4. NUMERICAL APPLICATIONS

In this section, we apply the homotopy perturbation Sumudu transform method [HPSTM] for solving heat and wave-like equations.

Example 4.1. Consider the following one-dimensional initial boundary value problem which describes the heat-like models (Noor and Mohyud-Din (2008) and Wazwaz and Gorguis (2004).

$$U_t = \frac{1}{2}x^2U_{xx}, \quad 0 < x < 1, \quad t > 0, \tag{11}$$

with boundary conditions

$$U(0,t) = 0, \quad U(1,t) = e^t, \tag{12}$$

and the initial condition

$$U(x,0) = x^2. \tag{13}$$

Taking the sumudu transform on both sides of equation (11) subject to the initial condition, we have

$$S[U(x,t)] = x^2 + \frac{1}{2}x^2u S[U_{xx}]. \tag{14}$$

The inverse of sumudu transform implies that

$$U(x,t) = x^2 + \frac{1}{2}x^2S^{-1}[uS[U_{xx}]]. \tag{15}$$

Now, applying the homotopy perturbation method, we get

$$\sum_{n=0}^{\infty} p^n U_n(x,t) = x^2 + p \left(\frac{1}{2}x^2S^{-1} \left[uS \left[\left(\sum_{n=0}^{\infty} p^n U_n(x,t) \right)_{xx} \right] \right] \right). \tag{16}$$

Comparing the coefficients of like powers of p , we have

$$\begin{aligned} p^0 : U_0(x,t) &= x^2, \\ p^1 : U_1(x,t) &= \frac{1}{2}x^2S^{-1}[uS[(U_0)_{xx}]] = x^2 t, \\ p^2 : U_2(x,t) &= \frac{1}{2}x^2S^{-1}[uS[(U_1)_{xx}]] = x^2 \frac{t^2}{2!} \end{aligned} \tag{17}$$

Proceeding in a similar manner, we have

$$\begin{aligned} p^3 : U_3(x,t) &= x^2 \frac{t^3}{3!}, \\ p^4 : U_4(x,t) &= x^2 \frac{t^4}{4!}, \\ &\vdots \end{aligned} \tag{18}$$

Therefore the solution $U(x,t)$ is given by

$$U(x,t) = x^2 \left(1 + t + \frac{t^2}{2!} + \frac{t^3}{3!} + \frac{t^4}{4!} + \dots \right), \tag{19}$$

in a series form, and

$$U(x,t) = x^2 e^t, \tag{20}$$

in closed form.

Example 4.2. Consider the following two-dimensional initial boundary value problem which describes the heat-like models (Noor and Mohyud-Din (2008) and Wazwaz and Gorguis (2004)).

$$U_t = \frac{1}{2}(y^2 U_{xx} + x^2 U_{yy}), \quad 0 < x, y < 1, \quad t > 0, \quad (21)$$

with boundary conditions

$$\begin{aligned} U_x(0, y, t) = 0, & \quad U_x(1, y, t) = 2 \sinh t, \\ U_y(x, 0, t) = 0, & \quad U_y(x, 1, t) = 2 \cosh t, \end{aligned} \quad (22)$$

and initial condition

$$U(x, y, 0) = y^2. \quad (23)$$

In a similar way as above, we have

$$\begin{aligned} \sum_{n=0}^{\infty} p^n U_n(x, y, t) = y^2 + p \left(\frac{1}{2} y^2 S^{-1} \left[u S \left[\left(\sum_{n=0}^{\infty} p^n U_n(x, y, t) \right)_{xx} \right] \right] \right. \\ \left. + \frac{1}{2} x^2 S^{-1} \left[u S \left[\left(\sum_{n=0}^{\infty} p^n U_n(x, y, t) \right)_{yy} \right] \right] \right). \end{aligned} \quad (24)$$

Comparing the coefficients of like powers of p , we have

$$\begin{aligned} p^0 : U_0(x, y, t) &= y^2, \\ p^1 : U_1(x, y, t) &= x^2 t, \\ p^2 : U_2(x, y, t) &= y^2 \frac{t^2}{2!}, \\ p^3 : U_3(x, y, t) &= x^2 \frac{t^3}{3!}, \\ p^4 : U_4(x, y, t) &= y^2 \frac{t^4}{4!}, \\ &\vdots \end{aligned} \quad (25)$$

Therefore the solution $U(x, y, t)$ is given by

$$U(x, y, t) = x^2 \left(t + \frac{t^3}{3!} + \frac{t^5}{5!} + \dots \right) + y^2 \left(1 + \frac{t^2}{2!} + \frac{t^4}{4!} + \dots \right), \quad (26)$$

which is in series form, and

$$U(x, y, t) = x^2 \sinh t + y^2 \cosh t, \quad (27)$$

in closed form.

Example 4.3. Consider the following three-dimensional inhomogeneous initial boundary value problem which describes the heat-like models (Noor and Mohyud-Din (2008) and Wazwaz and Gorguis (2004)).

$$U_t = x^4 y^4 z^4 + \frac{1}{36} (x^2 U_{xx} + y^2 U_{yy} + z^2 U_{zz}), \quad 0 < x, y, z < 1, \quad t > 0, \quad (28)$$

subject to the following boundary conditions

$$\begin{aligned} U(0, y, z, t) &= 0, & U(1, y, z, t) &= y^4 z^4 (e^t - 1), \\ U(x, 0, z, t) &= 0, & U(x, 1, z, t) &= x^4 z^4 (e^t - 1), \\ U(x, y, 0, t) &= 0, & U(x, y, 1, t) &= x^4 y^4 (e^t - 1), \end{aligned} \quad (29)$$

and the initial condition

$$U(x, y, z, 0) = 0. \quad (30)$$

In a similar way as above, we have

$$\begin{aligned} \sum_{n=0}^{\infty} p^n U_n(x, y, z, t) &= x^4 y^4 z^4 t + p \left[\frac{1}{36} x^2 S^{-1} \left[u S \left[\left(\sum_{n=0}^{\infty} p^n U_n(x, y, z, t) \right)_{xx} \right] \right] \right. \\ &\quad + \frac{1}{36} y^2 S^{-1} \left[u S \left[\left(\sum_{n=0}^{\infty} p^n U_n(x, y, z, t) \right)_{yy} \right] \right] \\ &\quad \left. + \frac{1}{36} z^2 S^{-1} \left[u S \left[\left(\sum_{n=0}^{\infty} p^n U_n(x, y, z, t) \right)_{zz} \right] \right] \right]. \end{aligned} \quad (31)$$

Comparing the coefficients of like powers of p , we have

$$\begin{aligned} p^0 : U_0(x, y, z, t) &= x^4 y^4 z^4 t, \\ p^1 : U_1(x, y, z, t) &= x^4 y^4 z^4 \frac{t^2}{2!}, \end{aligned}$$

$$\begin{aligned}
 p^2 : U_2(x, y, z, t) &= x^4 y^4 z^4 \frac{t^3}{3!}, \\
 p^3 : U_3(x, y, z, t) &= x^4 y^4 z^4 \frac{t^4}{4!}, \\
 p^4 : U_4(x, y, z, t) &= x^4 y^4 z^4 \frac{t^5}{5!}, \\
 &\vdots
 \end{aligned}
 \tag{32}$$

Therefore the solution $U(x, y, z, t)$ is given by

$$U(x, y, z, t) = x^4 y^4 z^4 \left(t + \frac{t^2}{2!} + \frac{t^3}{3!} + \frac{t^4}{4!} + \frac{t^5}{5!} + \dots \right),
 \tag{33}$$

in a series form, and

$$U(x, y, z, t) = x^4 y^4 z^4 (e^t - 1),
 \tag{34}$$

in closed form.

Example 4.4. Consider the following one-dimensional initial boundary value problem which describes the wave-like models (Noor and Mohyud-Din (2008) and Wazwaz and Gorguis (2004))

$$U_{tt} = \frac{1}{2} x^2 U_{xx}, \quad 0 < x < 1, \quad t > 0,
 \tag{35}$$

subject to the boundary conditions

$$U(0, t) = 0, \quad U(1, t) = 1 + \sinh t,
 \tag{36}$$

and the initial conditions

$$U(x, 0) = x, \quad U_t(x, 0) = x^2.
 \tag{37}$$

In a similar way as above, we have

$$\sum_{n=0}^{\infty} p^n U_n(x, t) = x + x^2 t + p \left(\frac{1}{2} x^2 S^{-1} \left[u^2 S \left[\left(\sum_{n=0}^{\infty} p^n U_n(x, t) \right)_{xx} \right] \right] \right).
 \tag{38}$$

Comparing the coefficients of like powers of p , we have

$$\begin{aligned}
 p^0 : U_0(x,t) &= x + x^2t, \\
 p^1 : U_1(x,t) &= x^2 \frac{t^3}{3!}, \\
 p^2 : U_2(x,t) &= x^2 \frac{t^5}{5!}, \\
 p^3 : U_3(x,t) &= x^2 \frac{t^7}{7!}, \\
 &\vdots
 \end{aligned}
 \tag{39}$$

Therefore the solution $U(x,t)$ is given by

$$U(x,t) = x + x^2 \left(t + \frac{t^3}{3!} + \frac{t^5}{5!} + \frac{t^7}{7!} + \dots \right),
 \tag{40}$$

in series form, and

$$U(x,t) = x + x^2 \sinh t,
 \tag{41}$$

in closed form.

Example 4.5. Consider the following two-dimensional initial boundary value problem which describes the wave-like models (Noor and Mohyud-Din (2008) and Wazwaz and Gorguis (2004)

$$U_{tt} = \frac{1}{12} (x^2 U_{xx} + y^2 U_{yy}), \quad 0 < x, y < 1, \quad t > 0,
 \tag{42}$$

subject to the Neumann boundary conditions

$$\begin{aligned}
 U_x(0, y, t) &= 0, & U_x(1, y, t) &= 4 \cosh t, \\
 U_y(x, 0, t) &= 0, & U_y(x, 1, t) &= 4 \sinh t,
 \end{aligned}
 \tag{43}$$

and the initial conditions

$$U(x, y, 0) = x^4, \quad U_t(x, y, 0) = y^4.
 \tag{44}$$

In a similar way as above, we have

$$\sum_{n=0}^{\infty} p^n U_n(x, y, t) = x^4 + y^4 t + p \left(\frac{1}{12} x^2 S^{-1} \left[u^2 S \left[\left(\sum_{n=0}^{\infty} p^n U_n(x, y, t) \right)_{xx} \right] \right] + \frac{1}{12} y^2 S^{-1} \left[u^2 S \left[\left(\sum_{n=0}^{\infty} p^n U_n(x, y, t) \right)_{yy} \right] \right] \right). \quad (45)$$

Comparing the coefficients of like powers of p , we have

$$\begin{aligned} p^0 : U_0(x, y, t) &= x^4 + y^4 t, \\ p^1 : U_1(x, y, t) &= x^4 \frac{t^2}{2!} + y^4 \frac{t^3}{3!}, \\ p^2 : U_2(x, y, t) &= x^4 \frac{t^4}{4!} + y^4 \frac{t^5}{5!}, \\ p^3 : U_3(x, y, t) &= x^4 \frac{t^6}{6!} + y^4 \frac{t^7}{7!}, \\ &\vdots \end{aligned} \quad (46)$$

Therefore the solution $U(x, y, t)$ is given by

$$U(x, y, t) = x^4 \left(1 + \frac{t^2}{2!} + \frac{t^4}{4!} + \dots \right) + y^4 \left(t + \frac{t^3}{3!} + \frac{t^5}{5!} + \dots \right), \quad (47)$$

in series form, and

$$U(x, y, t) = x^4 \cosh t + y^4 \sinh t, \quad (48)$$

in closed form.

Example 4.6. Consider the following three-dimensional inhomogeneous initial boundary value problem which describes the wave-like models (Noor and Mohyud-Din (2008) and Wazwaz and Gorguis (2004)

$$U_{tt} = (x^2 + y^2 + z^2) + \frac{1}{2}(x^2 U_{xx} + y^2 U_{yy} + z^2 U_{zz}), \quad 0 < x, y, z < 1, \quad t > 0, \quad (49)$$

subject to the boundary conditions

$$\begin{aligned} U(0, y, z, t) &= y^2(e^t - 1) + z^2(e^{-t} - 1), & U(1, y, z, t) &= (1 + y^2)(e^t - 1) + z^2(e^{-t} - 1), \\ U(x, 0, z, t) &= x^2(e^t - 1) + z^2(e^{-t} - 1), & U(x, 1, z, t) &= (1 + x^2)(e^t - 1) + z^2(e^{-t} - 1), \\ U(x, y, 0, t) &= (x^2 + y^2)(e^t - 1), & U(x, y, 1, t) &= (x^2 + y^2)(e^t - 1) + (e^{-t} - 1), \end{aligned} \tag{50}$$

and having the initial conditions

$$U(x, y, z, 0) = 0, \quad U_t(x, y, z, 0) = x^2 + y^2 - z^2. \tag{51}$$

In a similar way as above, we have

$$\begin{aligned} \sum_{n=0}^{\infty} p^n U_n(x, y, z, t) &= (x^2 + y^2 + z^2) \frac{t^2}{2} \\ &+ (x^2 + y^2 - z^2)t + p \left[\frac{1}{2} x^2 S^{-1} \left[u^2 S \left[\left(\sum_{n=0}^{\infty} p^n U_n(x, y, z, t) \right)_{xx} \right] \right] \right] \\ &+ \frac{1}{2} y^2 S^{-1} \left[u^2 S \left[\left(\sum_{n=0}^{\infty} p^n U_n(x, y, z, t) \right)_{yy} \right] \right] \\ &+ \frac{1}{2} z^2 S^{-1} \left[u^2 S \left[\left(\sum_{n=0}^{\infty} p^n U_n(x, y, z, t) \right)_{zz} \right] \right]. \end{aligned} \tag{52}$$

Comparing the coefficients of like powers of p , we have

$$\begin{aligned} p^0 : U_0(x, y, z, t) &= (x^2 + y^2 + z^2) \frac{t^2}{2} + (x^2 + y^2 - z^2)t, \\ p^1 : U_1(x, y, z, t) &= (x^2 + y^2 + z^2) \frac{t^4}{4!} + (x^2 + y^2 - z^2) \frac{t^3}{3!}, \\ p^2 : U_2(x, y, z, t) &= (x^2 + y^2 + z^2) \frac{t^6}{6!} + (x^2 + y^2 - z^2) \frac{t^5}{5!}, \\ p^3 : U_3(x, y, z, t) &= (x^2 + y^2 + z^2) \frac{t^8}{8!} + (x^2 + y^2 - z^2) \frac{t^7}{7!}, \\ &\vdots \end{aligned} \tag{53}$$

Therefore the solution $U(x, y, z, t)$ is given by

$$U(x, y, z, t) = (x^2 + y^2) \left(t + \frac{t^2}{2!} + \frac{t^3}{3!} + \frac{t^4}{4!} + \dots \right) + z^2 \left(-t + \frac{t^2}{2!} - \frac{t^3}{3!} + \frac{t^4}{4!} + \dots \right), \quad (54)$$

in the series form, and

$$U(x, y, z, t) = (x^2 + y^2)e^t + z^2e^{-t} - (x^2 + y^2 + z^2), \quad (55)$$

in closed form.

5. CONCLUSION

In this paper, we have applied the homotopy perturbation sumudu transform method (HPSTM) for solving heat and wave-like equations. It is worth mentioning that the proposed technique is capable of reducing the volume of the computational work as compared to the classical methods while still maintaining the high accuracy of the numerical result; the size reduction amounts to an improvement of the performance of the approach. The method gives more realistic series solutions that converge very rapidly in physical problems. The fact that the HPSTM solves nonlinear problems without using Adomian's polynomials is a clear advantage of this technique over the decomposition method. In conclusion, the HPSTM may be considered as a nice refinement in existing numerical techniques and might find the wide applications.

REFERENCES

- Asiru, M. A. 2002. Further properties of the Sumudu transform and its applications. *Int. J. Math. Educ.Sci.Tech.* **33**(3): 441-449.
- Asiru, M. A. 2001. Sumudu transform and the solution of integral equation of convolution type. *International Journal of Mathematical Education in Science and Technology.* **32**: 906-910.

- Belgacem, F. B. M., Karaballi, A. A. and Kalla, S. L. 2003. Analytical investigations of the Sumudu transform and applications to integral production equations. *Mathematical Problems in Engineering*. **3**: 103-118.
- Belgacem, F. B. M. and Karaballi, A. A. 2005. Sumudu transform fundamental properties investigations and applications. *International J. Appl. Math. Stoch. Anal.* 1-23.
- Eltayeb, H., Kilicman, A. and Fisher, B. 2010. A new integral transform and associated distributions. *Int. Trans. Spec. Func.* **21**(5): 367-379.
- Ganji, D. D. 2006. The applications of He's homotopy perturbation method to nonlinear equation arising in heat transfer. *Physics Letters A*. **335**: 337-341.
- Ghorbani, A. and Saberi-Nadjafi, J. 2007. He's homotopy perturbation method for calculating Adomian polynomials. *International Journal of Nonlinear Sciences and Numerical Simulation*. **8**: 229-232.
- Ghorbani, A. 2009. Beyond Adomian's polynomials: He polynomials. *Chaos Solitons Fractals*. **39**: 1486-1492.
- He, J. H. 1999. Homotopy perturbation technique. *Computer Methods in Applied Mechanics and Engineering*. **178**: 257-262.
- He, J. H. 2003. Homotopy perturbation method: a new nonlinear analytical technique. *Applied Mathematics and Computation*. **135**: 73-79.
- He, J. H. 2004. Comparison of homotopy perturbation method and homotopy analysis method. *Applied Mathematics and Computation*. **156**: 527-539.
- He, J. H. 2012. Asymptotic methods for solitary solutions and compactons. *Abstract and Applied Analysis*. Volume 2012, Article ID 916793, 130 pages
- Hesameddini, E. and Latifizadeh, H. 2009. An optimal choice of initial solutions in the homotopy perturbation method. *International Journal of Nonlinear Sciences and Numerical Simulation*. **101**: 389-1398.

- Hesameddini, E. and Latifizadeh, H. 2009. A new vision of the He's homotopy perturbation method. *International Journal of Nonlinear Sciences and Numerical Simulation*. **10**: 1415–1424.
- Kadem, A. 2005. Solving the one-dimensional neutron transport equation using Chebyshev polynomials and the Sumudu transform, *Analele Universitatii din Oradea. Fascicola Matematica*. **12**:153-171.
- Khan, Y. and Wu, Q. 2011. Homotopy perturbation transform method for nonlinear equations using He's polynomials. *Computer and Mathematics with Applications*. **61**(8): 1963-1967.
- Kilicman, A., Eltayeb, H. and Agarwal, P. Ravi. 2010. On Sumudu Transform and System of Differential Equations. *Abstract and Applied Analysis*. Article ID 598702, doi: 10.1155/2010/598702.
- Kilicman, A. and Eltayeb, H. 2010. A note on integral transforms and partial differential equations. *Applied Mathematical Sciences*. **4**(3):109-118.
- Kilicman, A. and Eltayeb, H. 2010. On the applications of Laplace and Sumudu transforms. *Journal of the Franklin Institute*. **347**(5): 848–862.
- Kilicman, A., Eltayeb, H. and Kamel Ariffin Mohd. Atan. 2011. A Note on the Comparison Between Laplace and Sumudu Transforms. *Bulletin of the Iranian Mathematical Society*. **37**(1): 131-141.
- Madani, M. and Fathizadeh, M. 2010. Homotopy perturbation algorithm using Laplace transformation. *Nonlinear Science Letters A*. **1**: 263–267.
- Mohyud-Din, S. T. 2009. Solving heat and wave-like equations using He's polynomials. *Mathematical Problems in Engineering*. doi:10.1155/2009/427516.
- Mohyud-Din, S. T. and Yildirim, A. 2010. Homotopy perturbation method for advection problems. *Nonlinear Science Letters A*. **1**: 307–312.

- Mohyud-Din, S.T., Noor, M. A. and Noor, K. I. 2009. Traveling wave solutions of seventh-order generalized KdV equation using He's polynomials. *International Journal of Nonlinear Sciences and Numerical Simulation*. **10**: 227–233.
- Noor, M.A. and Mohyud-Din, S.T. 2008. Modified variation iteration method for heat and wave-like equations. *Acta Applicandae Mathematicae*. **104**(3) 257-269.
- Rafei, M. and Ganji, D. D. 2006. Explicit solutions of helmhotz equation and fifth-order KdV equation using homotopy perturbation method. *International Journal of Nonlinear Sciences and Numerical Simulation*. **7**: 321–328.
- Saberi-Nadjafi, J. and Ghorbani, A. 2009. He's homotopy perturbation method: an effective tool for solving nonlinear integral and integro-differential equations. *Computers and Mathematics with Applications*. **58**: 1345–1351.
- Singh, J., Kumar, D. and Sushila. 2011. Homotopy perturbation sumudu transform method for nonlinear equations. *Adv. Theor. Appl. Mech.* **4**: 165-175.
- Watugala, G. K. 1998. Sumudu transform-a new integral transform to solve differential equations and control engineering problems. *Math. Eng. Indust.* **6**(4): 319-329.
- Wazwaz, A. M. and Gorguis, A. 2004, Exact solutions for heat-like and wave-like equations with variable coefficients. *Applied Mathematics and Computation*. **149** (1) 15-29.
- Weerakoon, S. 1994. Applications of Sumudu Transform to Partial Differential Equations. *Int. J. Math. Educ. Sci. Technol.* **25**(2): 277-283.
- Weerakoon, S. 1998. Complex inversion formula for Sumudu transforms. *Int. J. Math. Educ. Sci. Technol.* **29**(4): 618-621.
- Wilcox, R. 1970. Closed-form solution of the differential equation $(\partial^2/\partial x \partial y + ax(\partial/\partial x) + by(\partial/\partial y) + cxy + \partial/\partial t)P(x,y,t) = 0$ by normal-ordering exponential operators. *Journal of Mathematical Physics*. **11**(4): 1235-1237.

- Xu, L. 2007. He's homotopy perturbation method for a boundary layer equation in unbounded domain. *Computers and Mathematics with Applications*. **54**:1067–1070.
- Zhang, J. 2007. A Sumudu based algorithm for solving differential equations. *Comput. Sci. J. Moldova*. **15**: 303-313.