

PDF hosted at the Radboud Repository of the Radboud University Nijmegen

The following full text is a publisher's version.

For additional information about this publication click this link.

<http://hdl.handle.net/2066/128543>

Please be advised that this information was generated on 2017-12-05 and may be subject to change.

Measuring a caring culture in hospitals: a systematic review of instruments

G Hesselink,¹ E Kuis,² M Pijnenburg,¹ H Wollersheim¹

To cite: Hesselink G, Kuis E, Pijnenburg M, *et al*. Measuring a caring culture in hospitals: a systematic review of instruments. *BMJ Open* 2013;**3**:e003416. doi:10.1136/bmjopen-2013-003416

► Prepublication history and additional material for this paper is available online. To view these files please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2013-003416>).

Received 14 June 2013
Revised 20 August 2013
Accepted 21 August 2013

ABSTRACT

Objective: To identify instruments or components of instruments that aim to measure aspects of a caring culture-shared beliefs, norms and values that direct professionals and managers to act caring in hospitals, and to evaluate their psychometric properties.

Design: Systematic review.

Data sources: PubMed, CINAHL, EMBASE, PsychInfo, Web of Science and the International bibliography of the Social Sciences.

Study selection: Peer-reviewed articles describing (components of) instruments measuring aspects of a caring culture in a hospital setting. Studies had to report psychometric data regarding the reliability or validity of the instrument. Potentially useful instruments that were identified after the title and abstract scan were assessed on relevance by an expert panel (n=12) using the RAND-modified Delphi procedure.

Results: Of the 6399 references identified, 75 were examined in detail. 7 studies each covering a unique instrument met our inclusion criteria. On average, 24% of the instrument's items were considered relevant for measuring aspects of the hospital's caring culture. Studies showed moderate-to-high validity and reliability scores. Validity was addressed for 6 of the 7 instruments. Face, content (90%) and construct (60%) validity were the most frequently reported psychometric properties described. One study (14%) reported discriminant validity of the instrument. Reliability data were available for all of the instruments. Internal consistency was the most frequently reported psychometric property for the instruments and demonstrated by: a Cronbach's α coefficient (80%), subscale intercorrelations (60%), and item-total correlations (40%).

Conclusions: The ultimate standard for measuring a caring culture in hospitals does not exist. Existing instruments provide partial coverage and lack information on discriminant validity, responsiveness and feasibility. Characteristics of the instruments included in this review could provide useful input for the design of a reliable and valid instrument for measuring a caring culture in hospitals.

INTRODUCTION

Biomedical research and the concept of evidence-based medicine direct the discourse

ARTICLE SUMMARY

Strengths and limitations of this study

- We have reviewed an extensive body of literature and consulted a group of experts in multiple rounds (RAND-modified Delphi procedure) to identify relevant instruments.
- The possible biased inclusion of studies due to diffused research concepts (ie, 'caring' and 'organisational culture').
- Most instruments were successfully tested on their reliability as well as face- and construct-validity, but lack data on discriminant validity, feasibility and responsiveness.

on quality in healthcare. In the pursuit for better, safer and more cost-effective care numerous initiatives to measure, evaluate and improve care have been developed.¹⁻³ These result in a variety of objective quality indicators and 'good clinical practices' and contribute to a more rational and standardised way of organising care processes and professional decision-making. However, the focus on a rational and process-oriented view on healthcare improvement leaves another quality aspect undervalued that is of major importance to patients. This paper indicates this aspect as 'caring': the sensitivity of healthcare providers and organisations for what patients have to endure and for how patients experience the care they receive, and the art of attuning the care to these experiences.

The concept of 'caring' is closely related to the concept of 'patient-centred care'. In 2001, the Institute of Medicine formulated patient-centeredness as one of the six qualities of care domains to answer the current deficits in the quality of care provision.¹ Patient-centeredness is associated with positive outcomes such as patient satisfaction,⁴ medication adherence⁵⁻⁷ and more efficient use of health services,⁸ and reduction of costs.⁹ Although consensus on the definition of patient-centeredness lacks,^{7 10 11} authors agree to the following aspects: (1) eliciting

¹Scientific Institute for Quality of Healthcare (IQ healthcare), Radboud University Nijmegen Medical Centre, Nijmegen, The Netherlands

²University of Humanistic Studies, Utrecht, The Netherlands

Correspondence to

Dr Gijs Hesselink;
G.Hesselink@iq.umcn.nl



and respecting patients' preferences and values; (2) informing, involving and engaging patients and family members in the care process and (3) providing patients' physical comfort and emotional support. The concept of caring overlaps with patient-centeredness as it relates to being compassionate,¹² and being empathic,¹³ as a healthcare provider and as an institute. However, caring represents a broader meaning; it encompasses a careful attention to every single, unique patient: with interest, curiosity, concern and openness for what moves or puzzles a patient, and, then, rightly responding to it.^{14 15} Baart and Vosman,¹⁶ describe this as the fit or match between the need or wish of the patient and the care provided.

Although many studies focus on measuring aspects of care on a microlevel,^{6 17 18} and various studies found associations between caring professionals and improved clinical outcomes,^{12 13} research is limited to the extent to which these aspects are supported on a higher, 'meso' level.¹⁹ More specifically, hospitals require a caring culture, that is, beliefs, norms and values shared by professionals and the management throughout the organisation,^{20 21} that motivate, facilitate and direct these professionals to structurally act caring to patients and family. Appropriate means can help to diagnose the hospital's caring culture and thereby provide important input to evaluate the quality of care provided in a single hospital and between hospitals, over a period of time. It may also provide insight for further in-depth research and opportunities for improvement. The existence of appropriate instruments to measure a caring culture in hospitals is unknown. Therefore, the aim of this study is to identify possible existing instruments or components of instruments that aim to measure the extent to which hospitals are caring, and to systematically review existing instruments on their psychometric properties, feasibility and responsiveness.

METHODS

We planned and reported this systematic review in accordance with the preferred reporting items for systematic reviews and meta-analyses (PRISMA).²²

Data sources

We searched for English language, peer-reviewed, studies published between 1990 and 1 May 2012 using the following full-text databases: PubMed, CINAHL, EMBASE, PsychInfo, Web of Science and the International bibliography of the Social Sciences (IBSS). A specific search strategy was developed for each database. Online supplementary appendix 1 provides a detailed listing of search terms. The references of the selected studies were manually checked (snowballing) to identify additional relevant studies that were missed.

Study selection

Two reviewers (GH and EK) independently assessed inclusion eligibility of the retrieved studies using the

search strategy. The initial selection for inclusion was based on the title and abstract of the study. When the title and/or abstract provided insufficient information to determine the relevance, full-paper copies of the articles were retrieved and reviewed. For the final selection, full-text copies of the studies were examined by GH and EK to determine whether they fulfilled the inclusion criteria. Disagreement about inclusion was solved by discussion. When no consensus could be achieved, a third and fourth reviewer (HW/MP) decided. Studies included in this review had to meet all of the following criteria:

1. Peer-reviewed studies, published full-text, during the period from January 1990 to 1 May 2012, and with an abstract in English to compare studied instruments in the same language and to avoid misinterpretation of the purpose and the content of instruments due to language barriers.
2. Describing instruments or components (ie, items or domains) measuring aspects of a caring culture in a hospital setting, that is, beliefs, norms and values shared by professionals and the management throughout the organisation,^{20 21} that motivate, facilitate and direct these professionals and the management to structurally act caring to patients and family. No distinction was made between a 'caring culture' and a 'caring climate'. These constructs are highly inter-related which makes it difficult to determine where culture leaves and climate begins.²³ Studies with instruments examined in primary care settings (including nursing homes and rehabilitation centres) or administered to medical or nursing students were excluded.
3. Reporting psychometric data (ie, reliability or validity) regarding instruments or components to be included.

Systematic reviews, intervention studies or studies measuring patient satisfaction were excluded.

Evaluation of instrument items by a RAND-modified Delphi study

Items of potentially useful instruments, that were included after the title and abstract scan, were evaluated on relevance in a RAND-modified Delphi procedure.²⁴ The RAND-modified Delphi method facilitated a systematic process of evaluating instrument items and reaching a consensus on item relevance by the input of expert opinions. A multidisciplinary panel (n=12) of experts in the field of medical ethics, social and organisational sciences, one patient representative and persons with expertise in patient-centred care, caring and organisational culture were consulted in three rounds.

In step 1, the members of the expert panel received a tabulated list of the instruments included after the title and abstract scan and an additional number of instruments from studies that were excluded from this review. One instrument was included by snowballing references after the Delphi study and therefore not evaluated by the group of experts. Experts were instructed to individually rate on a 5-point scale (1 for lowest, up to 5 for

highest) by asking: "Please rate to what extent the item is a good measure for assessing the caring culture in hospitals." To support their choice, panel members were provided with the source and available psychometric properties of the instrument. The results of step 1 were processed into a summary report to facilitate step 2 (panel consensus meeting). In this report, based on the rating of the experts, the items were ranked on their mean score and categorised into three according to their potential to measure aspects of a caring culture in hospitals: a category of items with high potential, low potential, or uncertain potential (for discussion). Items were considered to be of high potential if the mean score was 4.2 or higher. This cut-off point for high-potential items was chosen to ensure a limited number of selected items, face validity and good reproducibility. A low overall ranking score (low-potential recommendations) included a mean score rating <4.0. For the category of uncertain potential or with dubious results (ie, ratings that were highly conflicting between panel members), the level of agreement between panel members was assessed in the consensus meeting.

In step 2, panel members were invited to the consensus meeting to discuss results from step 1 and to criticise instruments and specific items face-to-face. A personalised summary report provided panel members the opportunity to compare their individual scores to the overall distribution of scores and to discuss reasons for disagreement or conflict situations. The goal of the meeting was not to force consensus, but to distinguish well-founded disagreement and disagreement based on misunderstanding or irrational motives.²⁵ The following options were explained to the panel members: acceptance, rejection or adjustment of an item, or the formulation of a new item.

In step 3, a set of items was identified that passed the first round of individual rating as well as the second-round discussion. This set of items was sent to the expert panel by email. In addition, all panel members, including those not present at the meeting, were asked to rate the adjusted or the newly formulated items once more, were provided with a last opportunity to make remarks and were asked to approve the final set. Comments were discussed by the authors and final revisions were made.

Quality assessment of studies

The seven-criterion appraisal framework of Yu and Kirk,²⁶ based on the work of Greenhalgh *et al*,²⁷ Russel *et al*²⁸ and Grange *et al*²⁹ was modified to six quality criteria and applied to each included study. The total score possible for each instrument ranged from 0 to 12 (see online supplementary appendix 2). Two reviewers (GH and EK) separately assessed each study based on validity (eg, face, content, construct and criterion), reliability (eg, internal consistency, stability and equivalence), responsiveness, user-centeredness, sample size and feasibility. Discrepancies were resolved through discussion. If no consensus was reached, a third reviewer (HW) was consulted.

Data extraction

Data were abstracted into a standard data abstraction form covering general information about the instrument such as the name and source, the study setting (country, type of hospital and population), purpose, the way the instrument is administered to participants, items and scoring of items and subscales. Psychometric properties regarding the validity and reliability of measurement, the response rate, the feasibility in terms of time and cost investment and ease of use of the instrument, and information regarding responsiveness of the instrument were extracted as well. Data extraction was performed independently by two reviewers (GH and EK). Any disagreement was resolved by discussion among the reviewers and a final decision made by the third reviewer (HW).

RESULTS

Search results

Our initial search identified 6399 records (figure 1), of which 1935 were in PubMed, 1127 were in CINAHL, 1900 were in EMBASE, 324 were in PsychInfo, 764 were in Web of Science and 349 were in IBSS. The title and abstract scan resulted in 72 papers that, at first sight, met the inclusion criteria or raised doubt. Sixty-seven papers were excluded after full-text scan and based on the outcome of the Delphi study. Two additional articles were identified by manual review of the reference lists of the original 72 articles and were included after the full-text scan and the Delphi study. Thus, the final set consisted of seven unique studies that underwent full-text abstraction.

General description of the instruments

In total, seven instruments were included in the review (table 1). Two instruments, the Person-centered Climate Questionnaire-staff version (PCQ-S),^{30 31} and the




Figure 1 Flow diagram of the search process.

Table 1 General information of instruments included in the review

Instrument and reference	Country, hospital type and population	Purpose	Administration	Items and scoring	Subscales
Professional Practice Environment Scale ³⁶	USA Teaching hospital (n=1) Staff (n=849)	To measure eight characteristics of the professional practice environment in an acute care setting	Self-rating	38; 4-point Likert Scale ('strongly agree' to 'strongly disagree')	Handling disagreement and conflict (8); internal work motivation (7); control over practice (7); leadership and autonomy in clinical practice (5); staff relationships with physicians (2); teamwork (4); cultural sensitivity (3); communication about patients (2)
Swedish language Person-centred Climate Questionnaire—patient version ³⁰	Sweden Local hospitals (n=3) Patients (n=544)	To measure the extent to which hospital environments are experienced by patients as person-centred	Self-rating	17; 7-point Likert Scale ('no, I disagree completely' to 'yes, I agree completely') and 4-point Likert Scale ('of very little importance' to 'of very high importance')	Safety (10); everydayness (4); hospitality (3)
Scale for care quality climate ³⁵	UK Acute hospital trusts (n=86) Staff (n=17949)	To measure leadership to take account of the healthcare context; to measure the care quality orientation as perceived by hospital staff; to measure job satisfaction	Self-rating	7; 5-point scale ('strongly disagree' to 'strongly agree')	NR
English Language Person-centered Climate Questionnaire—patient Version ³¹	Australia Hospital facility providing short-stay elective surgery, diagnostic procedures and other planned services (n=1) Patients (108)	To measure the extent to which the climate of healthcare settings are perceived as being person-centered	Self-rating	17; 7-point Likert Scale ('no, I disagree completely' to 'yes, I agree completely')	Safety (NR); Hospitality (NR)
Swedish language Person-centred Climate Questionnaire—staff version ³²	Sweden Hospitals (n=3) Staff (n=600)	To measure the extent to which hospital environments are perceived by staff as person-centred	Self-rating	14; 6-point-Likert Scale ('no, I disagree completely' to 'yes, I agree completely')	A climate of safety (5); A climate of everydayness (5); A climate of community (4)

Continued

Table 1 Continued

Instrument and reference	Country, hospital type and population	Purpose	Administration	Items and scoring	Subscales
English language Person-centred Climate Questionnaire—staff version ³³	Australia Hospital facility providing short-stay elective surgery, diagnostic procedures and other planned services (n=1) Healthcare and support staff (n=52)	To measure the extent to which hospital environments are perceived by staff as person-centred	Self-rating	14; 6-point-Likert Scale ('no, I disagree completely' to 'yes, I agree completely')	A climate of safety (3); A climate of everydayness (4); A climate of community (3); A climate of comprehensibility (4)
Staff questionnaire ³⁴	UK Large district hospital (n=1) Nursing staff (n=97)	To measure the perceptions and experiences of the hospital staff around the organisation and delivery of patient-centred acute nursing care	Self-rating	45; 4-point Likert Scale ('always' to 'never')	NR

NR, not reported.

Person-centered Climate Questionnaire-patient version (PCQP),^{32 33} were distinctively studied in the Swedish as well as in the English language on their psychometric properties. Instruments comprised between 7 and 76 items.

Of the seven instruments, two were studied in Australia,^{31 33} two in the UK,^{34 35} two in Sweden,^{30 32} and one in the USA.³⁶ Instruments were studied in one to three hospitals, varying in type: local, district, acute, teaching and tertiary. One instrument was tested in 86 hospital trusts.³⁵ Of the seven instruments, two were tested with only patients or relatives,^{30 31} and five with only hospital (nursing, medical or support) staff.³²⁻³⁶ The sample size for the hospital staff ranged from 52 to 17 949 and for patients or relatives from 108 to 544.

Relevance of items

On average, 24% of the instrument's items were considered relevant for measuring aspects of the hospital's culture of caring. The percentage of relevant items for an instrument ranged between 4% and 47% (see [table 2](#)).

Quality assessment of studies

Studies fulfilled 3–8 of 12 quality item scores (mean fulfilled criteria (\pm SD), 5.7 (2.3)); see online supplementary appendix 3).

Validity

Validity was addressed in some way (eg, by tests in the study or by referring to previous tests) for all instruments, except for one instrument ([table 3](#)).³⁵ For five instruments,^{30-33 36} more than one type of validity was reported. Face or content validity was described for six,^{30-34 36} of the seven instruments. Face or content validity was evaluated by a panel of experts, clinicians or patients. Construct validity was established by principal component analysis for five^{30-33 36} of the seven instruments. Factors accounting for the total variance of the instrument varied between 60% and 72%. For two studies, construct validity was evaluated by confirmatory factor analysis.^{30 32} One study reported the ability of the instrument to detect true differences between hospital units by examining the dispersion of mean scores.³²

Reliability

Reliability data was available for all of the instruments ([table 3](#)). Internal consistency was the most frequently reported psychometric property for the instruments. Internal consistency was demonstrated by:

- ▶ A Cronbach's α coefficient,^{30-33 35 36}
- ▶ Subscale intercorrelations;^{30-33 36}
- ▶ Interitem or item-total correlations.³¹⁻³³

Most instruments showed an α ranging between 0.87 and 0.93. The α for the subscales of the instruments varied between 0.64 and 0.96. For three instruments the correlations between items and the total scale ranged between 0.24 and 0.71, 0.37 and 0.80, and 0.56 and 0.64. Stability was addressed for four instruments through test-

**Table 2** Relevant items of instruments to measure aspects of a caring culture within hospitals

Instrument	Formulation of relevant items	Percentage relevant items
Professional Practice Environment Scale ^{36*}	Freedom to make important patient care and work decisions Adequate support services allow me to spend time with patients Enough time and opportunity to discuss patient care problems with other staff Not being placed in a position of having to do things against my professional judgment	11
Person-Centered Climate Questionnaire—patient version ^{30 31†}	A place where I feel welcome A place where it is easy to talk to the staff A place where the staff takes notice of what I say A place where the staff come quickly when I need help A place where the staff uses language I can understand A place which is neat and clean A place where the staff have time for the patients	47
Scale for care quality climate ³⁵	A place where I have choices, for example, what to wear There is an emphasis on patient-focussed care in this organisation As a patient, I would be happy to have care provided by this organisation	29
Person-Centred Climate Questionnaire—staff version ^{32 33†}	A place where the staff use a language that the patients can understand A place where it is easy for the patients to keep in contact with their loved ones A place where it is easy for the patients to talk to the staff A place where the patients have someone to talk to if they so wish	29
Staff questionnaire ³⁴	I feel accountable for the care I give to my patients Patient care is organised around the needs of the individual patient	4

*The instrument was included after the Delphi study. Items were evaluated by the research team.

†Items of the Swedish language 'Person-Centered Climate Questionnaire—patient version' and the Swedish language 'Person-Centered Climate Questionnaire—staff version' are, after forward and back translation, identical to the English language versions and therefore not presented in this table.

retest reliability with 1 week interval between testing.^{30–33} Correlation coefficients varied between 0.51 and 0.75.

User-centeredness

Healthcare providers were involved to test the face and content validity for four instruments,^{32–34 36} and patients, respectively in two instruments.^{30 31} User views were taken into account in initial item generation for five instruments.^{30–33 36} An initial pool of items was usually generated from literature reviews and empirical research, and guided by theoretical constructs.^{30–33 35}

Sample size

Six instruments were tested with a sample size that was suitable for factor analysis based on Kass and Tinsley's³⁷ guideline for a ratio of 5–10 participants per item up to about 300 participants. If the number of participants reaches up to 300, test parameters tend to be stable regardless of the subject to variable ratio.³⁷ The sample size of four instruments,^{30 32 35 36} was high (ie, above 300) and for one instrument sufficient (ie, 5–10 participants per item).³¹

Feasibility and responsiveness

All instruments were self- (or peer) administered. Information regarding the time needed for completion, costs, perceived difficulties and training needs or instructions (eg, how to complete the questionnaire) were not reported for any of the studied instruments. Non-response was reported for all instruments, except for one.³⁶ However, the reasons for not participating were not evaluated in any of the studies. An assessment of responsiveness was conducted for none of the instruments.

DISCUSSION

This is the first systematic review of instruments evaluating aspects of a caring culture in hospital settings. Various instruments (ie, questionnaires) were found measuring aspects of a caring culture in hospitals. Moderate-to-high reliability and validity was reported for most of the instruments. However, the usefulness of these instruments is limited. The instruments consist of a low percentage of relevant items covering one or a few

Table 3 Psychometric properties of the included instruments

Instrument	Item generation and face/content validity	Construct and criterion validity	Reliability	Response rate (%)	Feasibility†	Responsiveness
Professional Practice Environment Scale ³⁶	Literature; empirical study on 'professional practice environment' <i>Face/content validity</i> Consultation of a multidisciplinary group of 7 clinicians (Original Scale)	<i>Construct validity</i> PCA: 8 factors accounted for 61.0% of the total variance	<i>Cronbach's α</i> $\alpha=0.93$ (total scale) <i>Inter-item correlation</i> 0.26–0.81	NR	NR	NR
Swedish language Person-Centred Climate Questionnaire—patient version ³⁰	Literature; qualitative research on care environments perceived as caring; stepwise item reduction technique using both statistics and theory <i>Face/content validity</i> Delphi assessment by an expert group (4 senior nurse researchers experienced in scale development) and 5 patients evaluating the relevance, clarity, readability and the scaling of the items	<i>Construct validity</i> ‡ PCA: 3 factors accounted for 65.1% of the total variance	<i>Cronbach's α</i> $\alpha=0.93$ (total scale) Test-retest reliability of the total scale $r=0.73$ (95% CI 0.58 to 0.85)	33	NR	NR
Scale for care quality climate ³⁵	Based on literature on care quality climate	NR	<i>Cronbach's α</i> $\alpha=0.87$ (total scale)	41	NR	NR
English Language Person-Centred Climate Questionnaire—patient version ³¹	Based on existing instrument: the Swedish version PCQ-P (forward and back translation by two translators)	<i>Construct validity</i> PCA: two factors accounted for 64.99% of the total variance	<i>Cronbach's α</i> $\alpha=0.90$ (total scale) Item-total correlations $r=0.37$ –0.80 <i>Test-retest reliability of the total scale</i> $r=0.70$ (95% CI 0.63 to 0.77)	29	NR	NR
Swedish language Person-centred Climate Questionnaire—staff version ³²	Literature and empirical research on the conceptualisation of organisational environments providing a person-centred climate <i>Face/content validity</i> Expert group (4 senior nurse researchers) evaluating the relevance, clarity, readability and the scaling of the items	<i>Construct validity</i> ‡ PCA: three factors accounted for 60.0% of the total variance	<i>Cronbach's α</i> $\alpha=0.88$ (total scale) Item-total correlations $r=0.56$ –0.64 <i>Test-retest reliability of the total scale</i> $r=0.51$ (95% CI 0.47–0.75)	57	NR	NR

Continued

Table 3 Continued

Instrument	Item generation and face/content validity	Construct and criterion validity	Reliability	Response rate (%)	Feasibility†	Responsiveness
English language Person-centred Climate Questionnaire—staff version ³³	Based on existing instrument: the Swedish version PCQ-S (forward and back translation by 2 translators)	<i>Construct validity</i> PCA: four factors accounted for 71.8% of the total variance	<i>Cronbach's α</i> α=0.89 (total scale) Item-total correlations r=0.24–0.71 <i>Test-retest reliability of the total scale</i> r=0.75 (95% CI 0.58 to 0.86)	66	NR	NR
Staff questionnaire ³⁴	Literature on nursing activity and the organisation and delivery of patient-centred care <i>Face/content validity</i> Focus group with nurses (n=10) piloting the questionnaires on appropriateness	NR	NR	25	NR	NR

CDI, Caring Dimension Inventory; NDI, Nursing Dimensions Inventory; NR, Not Reported; NUM, Nurse Unit Manager questionnaire; PCA, Principal Component Analysis.

Service climate refers to 'employee perceptions of the practices, procedures, and behaviours that get rewarded, supported and expected with regard to customer service and service quality'.

†In terms of time investment and/or costs.

‡Findings were supported by confirmatory factor analysis (CFA).

aspects of a caring culture in hospitals, leaving other important aspects unnoticed. Although most instruments were successfully tested on their reliability as well as face- and construct-validity, studies lack data on discriminant validity.³⁸ An instrument should demonstrate significant differences across hospitals if it is to be useful in discriminating between hospitals in terms of their caring culture. Information on feasibility in terms of instructions or training on rating, time investment, costs and non-response evaluation lacked for all of the instruments. Various studies revealed that it was not possible to explore reasons for not participating (by completing the questionnaire), because of the anonymous return and implied consent. Furthermore, the ability of the instrument to detect clinically important changes over time, such as tests for differences between individuals, factors associated with good outcome and treatment effect from group differences, were generally not examined or reported as well. All identified instruments were questionnaires. Questionnaires are useful in providing a first general overview of a hospital's caring culture by the input from a large sample within a short period of time. However, culture and caring are constructs that are difficult to identify and assess by quantitative research alone.^{14–16 39 40} Although being more time-consuming, in-depth interviews and observations are needed to identify and assess the underlying social constructions, attitudes and patterns of communication between care providers, patients and family members.³⁹

Our study has several limitations. First of all, this review focused on instruments measuring complex and disputed constructs such as 'patient-centred culture' and 'caring culture'. In the literature, for each of these constructs a widely agreed definition lacks. This hindered us in formulating strict inclusion and exclusion criteria and may have caused subjective selection of studies (and instruments). We tried to reduce the subjectivity on selecting studies by using the RAND-modified Delphi procedure. Cut-off points for selecting relevant instrument items were arbitrarily chosen as standard cut-off points for evaluating items on a 5-point Likert scale. Second, articles with potentially relevant instruments may not have been covered by our search strategy, because they did not describe one of our search terms related to a caring culture. For example, we did not find and examine the American Hospital-level Consumer Assessment of Health Plans Survey in this review. Third, only instruments measuring aspects of a caring culture in hospital settings were included in this review. This narrowed focus possibly left out two potentially useful instruments that were tested in the community care,^{41 42} which we identified in the title and abstract scan. Fourth, the authors did not investigate if instruments were sensitive to measure subcultures (eg, at the department level or among physicians or nurses).³⁸ This may raise questions about the appropriateness of the instruments to measure a caring culture in hospitals.

In conclusion, an ultimate standard for measuring a caring culture in hospitals does not exist. An instrument specifically aimed at measuring the caring culture in hospitals, covering a wide range of caring aspects, does not exist in one single instrument for patients nor for care providers. The items of the studied instruments included in this review that were appraised as relevant for measuring aspects of a caring culture could assist in the design of a comprehensive instrument. In particular, the PCQ-P and PCQ-S are useful, based on their relatively high number of relevant items. Further information on the reliability, validity, feasibility and responsiveness of such an instrument is warranted. A rigorous multimethod approach in which quantitative findings are further explored qualitatively and in-depth is important for providing an adequate diagnosis of a hospital's caring culture or its change over time.

Acknowledgements The authors would like to thank the members of the expert panel who participated in the Delphi study: Rianne van den Brink (Dutch Institute for Healthcare Improvement, CBO), Marjan Faber (Scientific Institute for Quality of Healthcare, IQ healthcare), Michel van Slobbe (Utrecht University School of Governance, USG), Guy Widdershoven (VU University Medical Center, VUmc), Evert van Leeuwen (Scientific Institute for Quality of Healthcare, IQ Healthcare), Esther Kuis (University of Humanistic Studies), Hans van Dartel (Leiden University Medical Center, LUMC), Jan den Bakker (Stichting Presentie), Jorke de Witte (Radboud University Nijmegen Medical Centre, RUNMC), Dolf de Boer (Netherlands Institute for Health Services Research, NIVEL), Frans Kingma (The Client Advisory Board for University Hospitals, CRAZ), Harriët Messing (Compassion for Care).

Contributors GH, EK, MP and HW were involved in conception and design of the study. GH and EK were responsible for data acquisition. GH and EK analysed and interpreted the data. GH and EK drafted the manuscript, which was critically revised for important intellectual content by all authors.

Funding This review was funded by the Dutch healthcare insurance organisation CZ.

Competing interests None.

Provenance and peer review Not commissioned; externally peer reviewed.

Data sharing statement No additional data are available.

Open Access This is an Open Access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 3.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/3.0/>

REFERENCES

1. Institute of Medicine. *Crossing the quality chasm: a new health system for the twenty-first century*. Washington: National Academies Press, 2001.
2. Hughes RG. *Patient safety and quality: an evidence-based handbook for nurses*. AHRQ, 2008.
3. Gandjour A, Kleinschmit F, Littmann V, et al. An evidence-based evaluation of quality and efficiency indicators. *Qual Manag Health Care* 2002;10:41–52.
4. Wolf DM, Lehman L, Quinlin R, et al. Effect of patient-centered care on patient satisfaction and quality of care. *J Nurs Care Qual* 2008;23:316–21.
5. Robinson JH, Callister LC, Berry JA, et al. Patient-centered care and adherence: definitions and applications to improve outcomes. *J Am Acad Nurse Pract* 2008;20:600–7.
6. Stewart M, Brown JB, Donner A, et al. The impact of patient-centered care on outcomes. *J Fam Pract* 2000;49:796–804.
7. Stewart M, Brown JB, Weston WW, et al. *Patient-centred medicine: transforming the clinical method*. 2nd edn. Radcliffe Medical Press, 2003.



8. Agency for Healthcare Research and Quality. National Healthcare Quality Report, 2005.
9. Little P, Everitt H, Williamson I, *et al*. Observational study of effect of patient-centredness and positive approach on outcomes of general practice consultations. *BMJ* 2001;323:908–11.
10. Epstein RM, Fiscella K, Lesser CS, *et al*. Why the nation needs a policy push on patient-centered health care. *Health Aff (Millwood)* 2010;29:1489–95.
11. Mead N, Bower P. Patient-centredness: a conceptual framework and review of the empirical literature. *Soc Sci Med* 2000;51:1087–110.
12. Del Canale S, Louis DZ, Maio V, *et al*. The relationship between physician empathy and disease complications: an empirical study of primary care physicians and their diabetic patients in Parma, Italy. *Acad Med* 2012;87:1243–9.
13. Lown BA, Rosen J, Martila J. An agenda for improving compassionate care: a survey shows about half of patients say such care is missing. *Health Aff (Millwood)* 2011;30:1772–8.
14. Goossens A, Baart A. Kwaliteit van zorg 2.0: menslievende, presente en zorgzame zorg. *Kwaliteit in Zorg* 2011;6:4–7.
15. van Heijst A. *Professional loving care. An ethical view of the healthcare sector*. Leuven: Peeters, 2011.
16. Baart A, Vosman F. Relationship based care and recognition. Part one: sketching good care from the theory of presence and five entries. In: Leget C, Gastman C, Verkerk M, eds. *Care, compassion and recognition: an ethical discussion*. Leuven: Peters, 2011.
17. Mallinger JB, Griggs JJ, Shields CG. Patient-centered care and breast cancer survivors' satisfaction with information. *Patient Educ Couns* 2005;57:342–9.
18. Smith F, Orrell M. Does the patient-centred approach help identify the needs of older people attending primary care? *Age Ageing* 2007;36:628–31.
19. Hudon C, Fortin M, Haggerty JL, *et al*. Measuring patients' perceptions of patient-centered care: a systematic review of tools for family medicine. *Ann Fam Med* 2011;9:155–64.
20. Schein EH. *Organisational culture and leadership*. 2nd edn. San Francisco: Jossey-Bass, 1992.
21. Davies H, Nutley SM, Mannion R. Organisational culture and quality of health care. *Qual Health Care*. 2000;9:111–19.
22. Moher D, Liberati A, Tetzlaff J, *et al*. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *Ann Intern Med* 2009;151:264–9.
23. Gershon RR, Stone PW, Bakken S, *et al*. Measurement of organizational culture and climate in healthcare. *J Nurs Adm* 2004;34:33–40.
24. Fitch K, Bernstein SJ, Aguilar MS, *et al*. *The RAND/UCLA appropriateness method user's manual*. Santa Monica, CA: RAND, 2001.
25. Uphoff EP, Wennekes L, Punt CJ, *et al*. Development of generic quality indicators for patient-centered cancer care by using a RAND modified Delphi method. *Cancer Nurs* 2012;35:29–37.
26. Yu J, Kirk M. Evaluation of empathy measurement tools in nursing: systematic review. *J Adv Nurs* 2009;65:1790–806.
27. Greenhalgh J, Long AF, Brettell AJ, *et al*. Reviewing and selecting outcome measures for use in routine practice. *J Eval Clin Pract* 1998;4:339–50.
28. Russell IT, Blasi ZD, Lambert MF, *et al*. Systematic reviews and meta-analyses: opportunities and threats. In: Templeton A, O'Brien P, eds. *Evidence-based fertility treatment*. London: RCOG Press, 1998.
29. Grange A, Bekker H, Noyes J, *et al*. Adequacy of health-related quality of life measures in children under 5-years-old: systematic review. *J Adv Nurs* 2007;59:197–220.
30. Edvardsson D, Sandman PO, Rasmussen B. Swedish language person-centered climate questionnaire—patient version: construction and psychometric evaluation. *J Adv Nurs* 2008;63:302–9.
31. Edvardsson D, Koch S, Nay R. Psychometric evaluation of the English language person-centered climate questionnaire—patient version. *West J Nurs Res* 2009;31:235–44.
32. Edvardsson D, Sandman PO, Rasmussen B. Construction and psychometric evaluation of the Swedish language person-centered climate questionnaire—staff version. *J Nurs Manag* 2009;17:790–5.
33. Edvardsson D, Koch S, Nay R. Psychometric evaluation of the English language person-centered climate questionnaire—staff version. *J Nurs Manag* 2010;18:54–60.
34. Haigh C, Ormandy P. Evaluation of the organisation and delivery of patient-centered acute nursing care. *Contemp Nurse* 2011;37:253–64.
35. Shipton H, Armstrong C, West M, *et al*. The impact of leadership and quality climate on hospital performance. *Int J Qual Health Care* 2008;20:439–45.
36. Erickson JI, Duffy ME, Gibbons MP, *et al*. Development and psychometric evaluation of the Professional Practice Environment (PPE) scale. *J Nurs Scholarsh* 2004;36:279–85.
37. Kass RA, Tinsley HEA. Factor analysis. *J Leisure Res* 1979;11:120–38.
38. Anderson N, West M. Measuring climate for work group innovation: development and validation of the team climate inventory. *J Organ Behav* 1998;19:235–58.
39. Scott JT, Mannion R, Davies HTO, *et al*. The quantitative measurement of organisational culture in health care: what instruments are available? *Health Serv Res* 2003;38:923–45.
40. Schein EH. Sense and nonsense about culture and climate. In: Ashkanasy NM, Wilderom CPM, Peterson MF, eds. *Handbook of organizational culture and climate*. Thousand Oaks: Sage Publications, 2000:xxiii–xxx.
41. Edvardsson D, Fetherstonhaugh D, Nay R, *et al*. Development and initial testing of the person-centered care assessment tool (P-CAT). *Int Psychogeriatr* 2010;22:101–8.
42. Lee M, Reuben DB, Ferrel BA. Multidimensional attitudes of medical residents and geriatrics follows toward older people. *J Am Geriatr Soc* 2005;53:489–94.

Measuring a caring culture in hospitals: a systematic review of instruments

G Hesselink, E Kuis, M Pijnenburg and H Wollersheim

BMJ Open 2013 3:
doi: 10.1136/bmjopen-2013-003416

Updated information and services can be found at:
<http://bmjopen.bmj.com/content/3/9/e003416>

Supplementary Material

Supplementary material can be found at:
<http://bmjopen.bmj.com/content/suppl/2013/09/23/bmjopen-2013-003416.DC1.html>

These include:

References

This article cites 32 articles, 8 of which you can access for free at:
<http://bmjopen.bmj.com/content/3/9/e003416#BIBL>

Open Access

This is an Open Access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 3.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/3.0/>

Email alerting service

Receive free email alerts when new articles cite this article. Sign up in the box at the top right corner of the online article.

Topic Collections

Articles on similar topics can be found in the following collections

[Ethics](#) (33)
[Patient-centred medicine](#) (154)

Notes

To request permissions go to:
<http://group.bmj.com/group/rights-licensing/permissions>

To order reprints go to:
<http://journals.bmj.com/cgi/reprintform>

To subscribe to BMJ go to:
<http://group.bmj.com/subscribe/>