



Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

1,5-Bis(2-methoxybenzylidene)thiocarbonohydrazide methanol monosolvate

Jianfeng Yu,^a Shiming Tang,^a Jingbin Zeng^a and Zifeng Yan^{b*}

^aDepartment of Chemistry, College of Science, China University of Petroleum, Qingdao 266555, People's Republic of China, and ^bSate Key Laboratory of Heavy Oil Processing, China University of Petroleum, Qingdao 266555, People's Republic of China

Correspondence e-mail: zfyancat@163.com


Received 20 May 2013; accepted 19 June 2013

Key indicators: single-crystal X-ray study; $T = 296\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.005\text{ \AA}$; R factor = 0.049; wR factor = 0.171; data-to-parameter ratio = 13.9.

The title compound, $\text{C}_{17}\text{H}_{18}\text{N}_4\text{O}_2\text{S}\cdot\text{CH}_3\text{OH}$, was synthesized by the condensation reaction of *o*-methoxybenzaldehyde with thiocarbonohydrazide in methanol. The two benzene rings are inclined each to other at $31.7(1)^\circ$. Intermolecular $\text{N}-\text{H}\cdots\text{O}$ and bifurcated $\text{O}-\text{H}\cdots\text{N}(\text{S})$ hydrogen bonds link two thiocarbonohydrazide and two solvent molecules into a centrosymmetric unit. These units, related by translation along the b axis, are further aggregated into columns through $\text{N}-\text{H}\cdots\text{S}$ hydrogen bonds.

Related literature

For biological activities of thiocarbonohydrazides, see: Liang (2003); Bacchi *et al.* (2005). For the crystal structures of related compounds, see: Fang *et al.* (2006); Feng *et al.* (2011); Zhao (2011).



Experimental

Crystal data

$\text{C}_{17}\text{H}_{18}\text{N}_4\text{O}_2\text{S}\cdot\text{CH}_3\text{OH}$	$\gamma = 79.550(3)^\circ$
$M_r = 374.46$	$V = 969.3(3)\text{ \AA}^3$
Triclinic, $P\bar{1}$	$Z = 2$
$a = 7.7223(15)\text{ \AA}$	Mo $K\alpha$ radiation
$b = 10.232(2)\text{ \AA}$	$\mu = 0.19\text{ mm}^{-1}$
$c = 12.648(3)\text{ \AA}$	$T = 296\text{ K}$
$\alpha = 85.938(3)^\circ$	$0.25 \times 0.21 \times 0.18\text{ mm}$
$\beta = 80.796(3)^\circ$	

Data collection

Bruker SMART APEX CCD area-detector diffractometer	4769 measured reflections
Absorption correction: multi-scan (<i>SADABS</i> ; Bruker, 2007)	3324 independent reflections
$T_{\min} = 0.954$, $T_{\max} = 0.966$	2766 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.024$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.049$	240 parameters
$wR(F^2) = 0.171$	H-atom parameters constrained
$S = 1.01$	$\Delta\rho_{\max} = 0.20\text{ e \AA}^{-3}$
3324 reflections	$\Delta\rho_{\min} = -0.21\text{ e \AA}^{-3}$

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
O3—H3—S1 ⁱ	0.82	2.80	3.534 (2)	150
O3—H3—N4 ⁱ	0.82	2.36	3.028 (3)	139
N3—H3A—O3	0.86	2.38	3.126 (3)	145
N1—H1—S1 ⁱⁱ	0.86	2.57	3.4184 (19)	169

Symmetry codes: (i) $-x + 1, -y + 1, -z$; (ii) $-x + 1, -y + 2, -z$.

Data collection: *SMART* (Bruker, 2007); cell refinement: *SAINT* (Bruker, 2007); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

The authors gratefully acknowledge the financial support of the Fundamental Research Funds for the Central Universities (grant No. 12CX04091A).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: CV5415).

References

- Bacchi, A., Carcelli, M., Pelagatti, P., Pelizzi, G., Rodriguez-Arguelles, M. C., Rogolino, D., Solinas, C. & Zani, F. (2005). *J. Inorg. Biochem.* **99**, 397–408.
- Bruker (2007). *SMART*, *SAINT* and *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Fang, X.-N., Xu, Y.-P., Guo, X.-F. & Zeng, X.-R. (2006). *Acta Cryst. E* **62**, o1052–o1054.
- Feng, L., Ji, H., Wang, R., Ge, H. & Li, L. (2011). *Acta Cryst. E* **67**, o1514.
- Liang, F.-Z. (2003). *J. Shandong Normal Univ. (Nat. Sci.)*, **18**, 50–51.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Zhao, X. (2011). *Acta Cryst. E* **67**, o2133.

supporting information

Acta Cryst. (2013). E69, o1147 [https://doi.org/10.1107/S1600536813016954]

1,5-Bis(2-methoxybenzylidene)thiocarbonohydrazide methanol monosolvate

Jianfeng Yu, Shiming Tang, Jingbin Zeng and Zifeng Yan

S1. Comment

Schiff bases of thiocarbonohydrazide are important organic intermediates owing to their biological activities (Liang, 2003; Bacchi *et al.*, 2005). In a continuation of structural study of Schiff bases of thiocarbonohydrazide (Fang *et al.*, 2006; Feng *et al.*, 2011; Zhao, 2011), we present here the title compound (I).

In (I) (Fig. 1), the bond lengths and angles are normal and correspond to those observed in 1,5-bis[(1E)-(2-methoxyphenyl)methylene]-thiocarbonohydrazide ethanol solvate (Fang, *et al.*, 2006), *N*^{''},*N*^{'''}-bis(1-phenylethylidene)thiocarbonohydrazide (Feng *et al.*, 2011) and *N*^{''},*N*^{'''}-bis(4-methoxybenzylidene)thiocarbonohydrazide methanol solvate (Zhao, 2011).


In the crystal, the benzene ring C3—C8 and the benzene ring C11—C16 are inclined each to other at 31.7 (1) $^{\circ}$. In the crystal, intermolecular N—H···O and bifurcated O—H···N(S) hydrogen bonds (Table 1) link two *M* and two solvent molecules into centrosymmetric unit. These units related by translation along the *b* axis are further aggregated into columns through the N—H···S hydrogen bonds (Table 1).

S2. Experimental

A 50 ml flask was charged with a magnetic stir bar, *o*-methoxybenzaldehyde (1 mmol), thiocarbonohydrazide (0.5 mmol) in 20 ml methanol. After stirring 3 h at 373 K, the resulting mixture was cooled to room temperature, and recrystallized from methanol, and afforded the title compound as a crystalline solid.

S3. Refinement

All H atoms were placed in geometrically idealized positions (C—H 0.93–0.96 Å, N—H 0.86 Å, O—H 0.82 Å) and treated as riding on their parent atoms, with $U_{\text{iso}}(\text{H}) = 1.2\text{--}1.5U_{\text{eq}}(\text{C}, \text{O}, \text{N})$.

**Figure 1**

The molecular structure of (I) showing the atomic numbering and 30% probability displacement ellipsoids.

1,5-Bis(2-methoxybenzylidene)thiocarbonohydrazide methanol monosolvate

Crystal data



$$M_r = 374.46$$

Triclinic, $P\bar{1}$

$$a = 7.7223 (15) \text{ \AA}$$

$$b = 10.232 (2) \text{ \AA}$$

$$c = 12.648 (3) \text{ \AA}$$

$$\alpha = 85.938 (3)^\circ$$

$$\beta = 80.796 (3)^\circ$$

$$\gamma = 79.550 (3)^\circ$$

$$V = 969.3 (3) \text{ \AA}^3$$

$$Z = 2$$

$$F(000) = 396$$

$$D_x = 1.283 \text{ Mg m}^{-3}$$

Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$

Cell parameters from 2955 reflections

$$\theta = 3.0\text{--}28.1^\circ$$

$$\mu = 0.19 \text{ mm}^{-1}$$

$$T = 296 \text{ K}$$

Block, colourless

$$0.25 \times 0.21 \times 0.18 \text{ mm}$$

Data collection

Bruker SMART APEX CCD area-detector
diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

phi and ω scans

Absorption correction: multi-scan
(*SADABS*; Bruker, 2007)

$$T_{\min} = 0.954, T_{\max} = 0.966$$

$$4769 \text{ measured reflections}$$

$$3324 \text{ independent reflections}$$

$$2766 \text{ reflections with } I > 2\sigma(I)$$

$$R_{\text{int}} = 0.024$$

$$\theta_{\max} = 25.0^\circ, \theta_{\min} = 1.6^\circ$$

$$h = -8 \rightarrow 9$$

$$k = -9 \rightarrow 12$$

$$l = -15 \rightarrow 13$$

Refinement

Refinement on F^2

Least-squares matrix: full

$$R[F^2 > 2\sigma(F^2)] = 0.049$$

$$wR(F^2) = 0.171$$

$$S = 1.01$$

$$3324 \text{ reflections}$$

$$240 \text{ parameters}$$

$$0 \text{ restraints}$$

Primary atom site location: structure-invariant
direct methods

Secondary atom site location: difference Fourier
map

Hydrogen site location: inferred from
neighbouring sites

H-atom parameters constrained

$$w = 1/[\sigma^2(F_o^2) + (0.1P)^2 + 0.3851P]$$

where $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\max} = 0.010$
 $\Delta\rho_{\max} = 0.20 \text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.21 \text{ e \AA}^{-3}$

Extinction correction: *SHELXL97* (Sheldrick, 2008), $F_c^* = kF_c[1 + 0.001xF_c^2\lambda^3/\sin(2\theta)]^{-1/4}$
Extinction coefficient: 0.038 (7)

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
N1	0.4120 (3)	0.85311 (18)	0.10378 (15)	0.0503 (5)
H1	0.4497	0.9278	0.0942	0.060*
N2	0.3689 (3)	0.80149 (19)	0.20555 (15)	0.0524 (5)
N3	0.3445 (2)	0.66888 (17)	0.04305 (15)	0.0476 (5)
H3A	0.3322	0.6395	0.1087	0.057*
N4	0.3114 (2)	0.59365 (18)	-0.03557 (15)	0.0480 (5)
O1	0.2688 (4)	1.0468 (2)	0.44226 (17)	0.1014 (9)
O2	0.1142 (3)	0.27193 (17)	0.08315 (18)	0.0739 (6)
S1	0.43309 (9)	0.85466 (6)	-0.10545 (5)	0.0567 (3)
C1	0.3953 (3)	0.7867 (2)	0.01844 (18)	0.0438 (5)
C2	0.3709 (3)	0.8750 (3)	0.28184 (19)	0.0574 (6)
H2	0.3972	0.9601	0.2666	0.069*
C3	0.3324 (4)	0.8276 (3)	0.3932 (2)	0.0644 (7)
C4	0.3466 (6)	0.6932 (3)	0.4199 (3)	0.0963 (11)
H4	0.3800	0.6319	0.3662	0.116*
C5	0.3117 (8)	0.6491 (4)	0.5254 (3)	0.144 (2)
H5	0.3207	0.5585	0.5428	0.173*
C6	0.2634 (8)	0.7397 (5)	0.6049 (3)	0.137 (2)
H6	0.2389	0.7098	0.6760	0.164*
C7	0.2510 (6)	0.8722 (4)	0.5809 (3)	0.1038 (12)
H7	0.2214	0.9324	0.6354	0.125*
C8	0.2826 (4)	0.9174 (3)	0.4751 (2)	0.0720 (7)
C9	0.2107 (7)	1.1424 (4)	0.5220 (3)	0.1161 (15)
H9A	0.0990	1.1274	0.5620	0.174*
H9B	0.1962	1.2301	0.4885	0.174*
H9C	0.2977	1.1346	0.5695	0.174*
C10	0.2413 (3)	0.4934 (2)	0.0006 (2)	0.0487 (5)
H10	0.2192	0.4771	0.0744	0.058*
C11	0.1943 (3)	0.4031 (2)	-0.0700 (2)	0.0543 (6)
C12	0.2091 (4)	0.4269 (3)	-0.1782 (2)	0.0731 (8)
H12	0.2533	0.5019	-0.2087	0.088*
C13	0.1593 (5)	0.3415 (4)	-0.2435 (3)	0.0937 (11)

H13	0.1698	0.3585	-0.3173	0.112*
C14	0.0940 (4)	0.2309 (4)	-0.1973 (3)	0.0958 (12)
H14	0.0581	0.1741	-0.2404	0.115*
C15	0.0809 (4)	0.2032 (3)	-0.0901 (3)	0.0804 (9)
H15	0.0392	0.1268	-0.0607	0.097*
C16	0.1297 (3)	0.2888 (2)	-0.0246 (3)	0.0600 (7)
C17	0.0616 (5)	0.1520 (3)	0.1332 (4)	0.0952 (11)
H17A	-0.0618	0.1534	0.1290	0.143*
H17B	0.0787	0.1458	0.2070	0.143*
H17C	0.1326	0.0766	0.0970	0.143*
O3	0.3985 (3)	0.4521 (2)	0.22744 (16)	0.0803 (6)
H3	0.4610	0.4009	0.1842	0.121*
C18	0.2837 (6)	0.3838 (4)	0.2970 (3)	0.1132 (14)
H18A	0.3517	0.3184	0.3393	0.170*
H18B	0.2173	0.3406	0.2563	0.170*
H18C	0.2031	0.4454	0.3434	0.170*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
N1	0.0627 (11)	0.0401 (10)	0.0515 (11)	-0.0201 (8)	-0.0034 (8)	-0.0085 (8)
N2	0.0588 (11)	0.0468 (10)	0.0529 (11)	-0.0149 (9)	-0.0028 (8)	-0.0086 (9)
N3	0.0554 (10)	0.0389 (9)	0.0514 (10)	-0.0155 (8)	-0.0049 (8)	-0.0098 (8)
N4	0.0496 (10)	0.0398 (9)	0.0569 (11)	-0.0113 (8)	-0.0059 (8)	-0.0129 (8)
O1	0.172 (2)	0.0723 (14)	0.0597 (12)	-0.0446 (15)	0.0175 (13)	-0.0220 (10)
O2	0.0782 (12)	0.0486 (10)	0.1034 (16)	-0.0266 (9)	-0.0268 (11)	0.0098 (10)
S1	0.0740 (4)	0.0492 (4)	0.0532 (4)	-0.0275 (3)	-0.0076 (3)	-0.0062 (3)
C1	0.0403 (10)	0.0361 (10)	0.0555 (12)	-0.0073 (8)	-0.0049 (9)	-0.0098 (9)
C2	0.0680 (14)	0.0541 (13)	0.0526 (13)	-0.0203 (11)	-0.0022 (11)	-0.0106 (11)
C3	0.0760 (16)	0.0634 (16)	0.0527 (14)	-0.0142 (13)	-0.0024 (12)	-0.0053 (12)
C4	0.140 (3)	0.0634 (18)	0.0710 (19)	-0.0009 (19)	0.0064 (19)	-0.0020 (15)
C5	0.235 (6)	0.077 (2)	0.085 (3)	0.015 (3)	0.022 (3)	0.022 (2)
C6	0.212 (5)	0.102 (3)	0.066 (2)	0.021 (3)	0.007 (3)	0.017 (2)
C7	0.146 (3)	0.101 (3)	0.0549 (17)	-0.007 (2)	-0.0002 (19)	-0.0093 (17)
C8	0.0876 (19)	0.0748 (18)	0.0527 (14)	-0.0158 (15)	-0.0034 (13)	-0.0082 (13)
C9	0.188 (4)	0.087 (2)	0.074 (2)	-0.047 (3)	0.019 (2)	-0.0335 (19)
C10	0.0453 (11)	0.0381 (11)	0.0653 (14)	-0.0105 (9)	-0.0103 (10)	-0.0080 (10)
C11	0.0439 (11)	0.0452 (12)	0.0758 (16)	-0.0115 (9)	-0.0040 (10)	-0.0185 (11)
C12	0.0740 (17)	0.0731 (18)	0.0786 (19)	-0.0300 (14)	-0.0008 (14)	-0.0262 (15)
C13	0.098 (2)	0.109 (3)	0.083 (2)	-0.042 (2)	0.0038 (17)	-0.044 (2)
C14	0.085 (2)	0.095 (2)	0.118 (3)	-0.0436 (19)	0.0095 (19)	-0.062 (2)
C15	0.0646 (16)	0.0599 (16)	0.122 (3)	-0.0255 (13)	0.0028 (16)	-0.0369 (17)
C16	0.0412 (11)	0.0383 (12)	0.102 (2)	-0.0056 (9)	-0.0084 (12)	-0.0193 (12)
C17	0.082 (2)	0.0633 (18)	0.148 (3)	-0.0364 (16)	-0.026 (2)	0.026 (2)
O3	0.1011 (15)	0.0707 (13)	0.0658 (12)	-0.0183 (11)	0.0032 (10)	-0.0064 (10)
C18	0.152 (4)	0.103 (3)	0.084 (2)	-0.050 (3)	0.014 (2)	0.000 (2)

Geometric parameters (\AA , $^{\circ}$)

N1—C1	1.351 (3)	C7—H7	0.9300
N1—N2	1.370 (3)	C9—H9A	0.9600
N1—H1	0.8600	C9—H9B	0.9600
N2—C2	1.268 (3)	C9—H9C	0.9600
N3—C1	1.335 (3)	C10—C11	1.458 (3)
N3—N4	1.381 (2)	C10—H10	0.9300
N3—H3A	0.8600	C11—C12	1.363 (4)
N4—C10	1.269 (3)	C11—C16	1.404 (3)
O1—C8	1.350 (4)	C12—C13	1.386 (4)
O1—C9	1.419 (4)	C12—H12	0.9300
O2—C16	1.350 (4)	C13—C14	1.377 (5)
O2—C17	1.435 (3)	C13—H13	0.9300
S1—C1	1.672 (2)	C14—C15	1.356 (5)
C2—C3	1.459 (4)	C14—H14	0.9300
C2—H2	0.9300	C15—C16	1.385 (4)
C3—C4	1.382 (4)	C15—H15	0.9300
C3—C8	1.395 (4)	C17—H17A	0.9600
C4—C5	1.379 (5)	C17—H17B	0.9600
C4—H4	0.9300	C17—H17C	0.9600
C5—C6	1.378 (6)	O3—C18	1.391 (4)
C5—H5	0.9300	O3—H3	0.8200
C6—C7	1.358 (6)	C18—H18A	0.9600
C6—H6	0.9300	C18—H18B	0.9600
C7—C8	1.383 (4)	C18—H18C	0.9600
C1—N1—N2	119.98 (18)	O1—C9—H9C	109.5
C1—N1—H1	120.0	H9A—C9—H9C	109.5
N2—N1—H1	120.0	H9B—C9—H9C	109.5
C2—N2—N1	116.55 (19)	N4—C10—C11	122.0 (2)
C1—N3—N4	120.89 (19)	N4—C10—H10	119.0
C1—N3—H3A	119.6	C11—C10—H10	119.0
N4—N3—H3A	119.6	C12—C11—C16	119.1 (2)
C10—N4—N3	113.93 (19)	C12—C11—C10	122.2 (2)
C8—O1—C9	117.2 (3)	C16—C11—C10	118.7 (2)
C16—O2—C17	118.8 (3)	C11—C12—C13	121.2 (3)
N3—C1—N1	114.5 (2)	C11—C12—H12	119.4
N3—C1—S1	125.35 (17)	C13—C12—H12	119.4
N1—C1—S1	120.10 (16)	C14—C13—C12	118.8 (4)
N2—C2—C3	120.8 (2)	C14—C13—H13	120.6
N2—C2—H2	119.6	C12—C13—H13	120.6
C3—C2—H2	119.6	C15—C14—C13	121.4 (3)
C4—C3—C8	118.6 (3)	C15—C14—H14	119.3
C4—C3—C2	120.9 (3)	C13—C14—H14	119.3
C8—C3—C2	120.5 (2)	C14—C15—C16	119.9 (3)
C5—C4—C3	120.5 (3)	C14—C15—H15	120.0
C5—C4—H4	119.7	C16—C15—H15	120.0

C3—C4—H4	119.7	O2—C16—C15	123.9 (3)
C6—C5—C4	119.7 (4)	O2—C16—C11	116.5 (2)
C6—C5—H5	120.1	C15—C16—C11	119.5 (3)
C4—C5—H5	120.1	O2—C17—H17A	109.5
C7—C6—C5	120.8 (4)	O2—C17—H17B	109.5
C7—C6—H6	119.6	H17A—C17—H17B	109.5
C5—C6—H6	119.6	O2—C17—H17C	109.5
C6—C7—C8	119.8 (3)	H17A—C17—H17C	109.5
C6—C7—H7	120.1	H17B—C17—H17C	109.5
C8—C7—H7	120.1	C18—O3—H3	109.5
O1—C8—C7	124.6 (3)	O3—C18—H18A	109.5
O1—C8—C3	115.0 (2)	O3—C18—H18B	109.5
C7—C8—C3	120.4 (3)	H18A—C18—H18B	109.5
O1—C9—H9A	109.5	O3—C18—H18C	109.5
O1—C9—H9B	109.5	H18A—C18—H18C	109.5
H9A—C9—H9B	109.5	H18B—C18—H18C	109.5

Hydrogen-bond geometry (Å, °)

D—H···A	D—H	H···A	D···A	D—H···A
O3—H3···S1 ⁱ	0.82	2.80	3.534 (2)	150
O3—H3···N4 ⁱ	0.82	2.36	3.028 (3)	139
N3—H3A···O3	0.86	2.38	3.126 (3)	145
N1—H1···S1 ⁱⁱ	0.86	2.57	3.4184 (19)	169
N3—H3A···N2	0.86	2.21	2.591 (3)	106

Symmetry codes: (i) $-x+1, -y+1, -z$; (ii) $-x+1, -y+2, -z$.