
IJMMS 26:10 (2001) 643–648
 PII. S0161171201006317
<http://ijmms.hindawi.com>
 © Hindawi Publishing Corp.

STABILITY IN E -CONVEX PROGRAMMING

EBRAHIM A. YOUNESS

(Received 12 April 2000)

ABSTRACT. We define and analyze two kinds of stability in E -convex programming problem in which the feasible domain is affected by an operator E . The first kind of this stability is that the set of all operators E that make an optimal set stable while the other kind is that the set of all operators E that make certain side of the feasible domain still active.

2000 Mathematics Subject Classification. 90Cxx.

1. Introduction. The stability notion plays an important role in the mathematical programming field, it is important for solver or for the decision maker to preserve effort and time.

Stability in mathematical programming has many types, one of these types depends on making perturbation to the decision space or to the objective space or to both by a parameter. This type is called stability in parametric programming problems (see [3, 4, 5]). Other types are called internal and external stability. This kind of stability depends on the set of efficient solution for multi-objective programming problems, namely, if for each efficient solution there exist some points in decision space dominated by it (this is external stability). On the other hand, internal stability means that any efficient solution is not preferred to another efficient solution, see [6].

In this paper, the author presents a new type of stability in mathematical programming. This type is very important in composite programming [2, 8] and in E -convex programming which was presented by the author in [9]. This kind of stability is called E -stability which is discussed in this paper.

Now, we give some examples to show that there are many operators E that transform an optimal point to another optimal point.

EXAMPLE 1.1. Consider the following problem:

$$\min(x-1)^2 + (y-1)^2 \quad (1.1)$$

subject to $x, y \geq 0$. It is clear that the optimal solution is $(\bar{x}, \bar{y}) = (1, 1)$. Operators $E: \mathbb{R}^2 \rightarrow \mathbb{R}^2$, defined as

$$E(x, y) = (2x-1, 2y-1), \quad E(x, y) = (x^2, y^2), \quad E(x, y) = (xy, y), \quad (1.2)$$

map $(1, 1)$ to $(1, 1)$.

EXAMPLE 1.2. Consider the problem

$$\max\{x+y\} \quad (1.3)$$

subject to $M = \{(x, y) \in \mathbb{R}^2 : x+y \leq 8, x \leq 6, y \leq 5, x+y \geq 1\}$.

The set of optimal solutions is

$$S = \{(x, y) \in \mathbb{R}^2 : (x, y) = \lambda(3, 5) + (1 - \lambda)(6, 2), 0 \leq \lambda \leq 1\}. \quad (1.4)$$

It is clear that a transformation $E : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ which maps (x, y) to $(9 - x, 7 - y)$ assigns any point in S again to a point in S .

Also, for the solution $(3, 5)$ we can find more than one $E : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ such that $E(3, 5) \in S$, for example

$$\begin{aligned} E(x, y) &= \left(xy - x - y - 1, \frac{1}{4}(x + y) \right), \\ E(x, y) &= \left(\frac{1}{2}x^2, y - \frac{3}{2} \right), \\ E(x, y) &= \left(x^2y - 41, \frac{12y^2}{25x} \right). \end{aligned} \quad (1.5)$$

The above two examples lead to the discussion of the set of all these operators qualitatively.

2. E-stability of an optimal set. Consider the following mathematical programming problem:

$$\min f(x) \quad \text{subject to} \quad M = \{x \in \mathbb{R}^n : g_r(x) \leq 0, r = 1, 2, \dots, m\}. \quad (2.1)$$

Denote S the set of optimal solutions for problem (2.1).

DEFINITION 2.1. An E -stability of S is denoted by $D(\bar{x})$, $\bar{x} \in S$ and is defined as

$$D(\bar{x}) = \{E : \mathbb{R}^n \rightarrow \mathbb{R}^n : E(\bar{x}) \in S\}. \quad (2.2)$$

Let ξ be the space of all operators from \mathbb{R}^n to \mathbb{R}^n then $D(\bar{x})$ is a point-to-set map from S to ξ .

DEFINITION 2.2 (see [1]). A point-to-set map $F : X \rightarrow 2^Y$ is said to be closed at a point $x^0 \in X$; if for each pair of sequences $\{x^n\} \subset X$ and $\{y^n\} \subset Y$, $n = 1, 2, \dots$ with the properties $x^n \rightarrow x^0$, $y^n \in F(x^n)$ and $y^n \rightarrow y^0$ it follows that $y^0 \in F(x^0)$.

PROPOSITION 2.3. If S is closed, then a point-to-set map $D : S \rightarrow \xi$ is closed.

PROOF. Let $\{x^t\} \subset S$, $x^t \rightarrow x^0$ as $t \rightarrow \infty$ and let $\{E^t\} \subset \xi$, $E^t \rightarrow E^0$ as $t \rightarrow \infty$ such that $E^t \in D(x^t)$, then we have a sequence $f(E^t x^t)$ with $f(E^t x^t) \leq f(x)$ for each $x \in M$. Since S is closed, then $f(E^0 x^0) \leq f(x)$ for all $x \in M$. Hence $E^0 \in D(x^0)$ and D is closed. \square

PROPOSITION 2.4. If f is lower-semicontinuous, then the set $D(x^0)$, $x^0 \in S$, is closed.

PROOF. Let E^n be a sequence of $D(x^0)$ and $E^n \rightarrow E^0$ as $n \rightarrow \infty$, then $f(E^n x^0) \leq f(x)$ for each $x \in M$. Since f is lower-semicontinuous, then

$$f(E^0 x^0) = f\left(\lim_{n \rightarrow \infty} E^n x^0\right) \leq \liminf_{n \rightarrow \infty} f(E^n x^0) \leq f(x) \quad (2.3)$$

for each $x \in M$, that is, $E^0 x^0 \in S$ and $E^0 \in D(x^0)$. Hence $D(x^0)$ is closed. \square

THEOREM 2.5. *If f is lower-semicontinuous and D is upper-semicontinuous at $x^0 \in S$, then D is closed.*

PROOF. From Proposition 2.3, the set $D(x^0)$ is closed and from [1, Lemma 2.2.1] the mapping D is closed. \square

THEOREM 2.6. *If S is compact, then D is upper-semicontinuous.*

PROOF. Since S is compact, then S is closed and, from Proposition 2.3, D is closed. From [1, Lemma 2.2.3] the mapping D is upper-semicontinuous. \square

THEOREM 2.7. *Let f be linear, S convex, and M^* a dual space of M . Then a point-to-set map $D : S \rightarrow M^*$ is convex.*

PROOF. Assume that $\bar{x}, \tilde{x} \in S$ and $E \in D(\lambda\bar{x} + (1-\lambda)\tilde{x})$, $0 \leq \lambda \leq 1$, then $\lambda E\bar{x} + (1-\lambda)E\tilde{x} \in S$. Let $E \notin \lambda D(\bar{x}) + (1-\lambda)D(\tilde{x})$ for any $0 \leq \lambda \leq 1$, then $E \notin D(\bar{x}) \cup D(\tilde{x})$ and hence $E\bar{x}, E\tilde{x} \notin S$. Therefore, there exist $\hat{x}, x^* \in M$ such that

$$f(\lambda\hat{x} + (1-\lambda)x^*) < f(\lambda E\bar{x} + (1-\lambda)E\tilde{x}), \tag{2.4}$$

which is a contradiction. Then

$$D(\lambda\bar{x} + (1-\lambda)\tilde{x}) \subset \lambda D(\bar{x}) + (1-\lambda)D(\tilde{x}) \tag{2.5}$$

and hence D is convex. \square

DEFINITION 2.8. Let S be a subset of \mathbb{R}^n and $E : \mathbb{R}^n \rightarrow \mathbb{R}^n$ be an operator. The set S is called an E -convex set if and only if

$$\lambda Ex + (1-\lambda)Ey \in S, \quad \forall x, y \in S, 0 \leq \lambda \leq 1. \tag{2.6}$$

(For more details about E -convex sets see [9].)

THEOREM 2.9. *If S is an E -convex set with respect to $E \in D(\bar{x})$, $\bar{x} \in S$, then $D(\bar{x}) = \bigcap_n D(E^n \bar{x})$.*

PROOF. Let $E \in D(\bar{x})$, then $E(\bar{x}) \in S$. Let $E \notin D(E\bar{x})$, then $E(E\bar{x}) \notin S$ which contradicts the E -convexity of S . Hence $E \in D(E\bar{x})$, that is, $D(\bar{x}) \subset D(E\bar{x})$. Similarly, $D(E\bar{x}) \subset D(E^2\bar{x}) \subset \dots \subset D(E^n\bar{x})$. Thus $D(\bar{x}) = \bigcap_n D(E^n\bar{x})$. \square

THEOREM 2.10. *Let f be a strictly convex function. If $E \in D(\bar{x})$ then $E^{-1} \in D(\bar{x})$.*

PROOF. Let $E \in D(\bar{x})$, then $E(\bar{x}) \in S$. If $E^{-1} \notin D(\bar{x})$, then $E^{-1}(\bar{x}) \notin S$ and $\bar{x} \notin E(S)$. Since f is strictly convex, then \bar{x} is the unique minimal point, that is, $s = \{\bar{x}\}$ and hence $E(\bar{x}) = \bar{x}$. Thus $\bar{x} \notin E(S)$ contradicts $E(\bar{x}) = \bar{x}$. Hence $E^{-1} \in D(\bar{x})$. \square

THEOREM 2.11. *Let S be a convex cone with vertex at the origin. If $E \in D(\bar{x})$, then $aE + b \in D(\bar{x})$ for each real numbers $a, b \geq 0$, $(a, b) \neq 0$.*

PROOF. Since $E \in D(\bar{x})$, then $E(\bar{x}) \in S$. So $(aE + b)\bar{x} = aE(\bar{x}) + bI(\bar{x})$, where I is the identity map. Since S is convex cone, then for $a, b \geq 0$, $(a, b) \neq 0$, $aE(\bar{x})$ and $bI(\bar{x})$ belong to S . Thus $aE(\bar{x}) + bI(\bar{x}) \in S$ and hence $aE + b \in D(\bar{x})$. \square

THEOREM 2.12. *Let S be a convex cone with vertex at the origin. If $D(x) - \{I\}$, $x \in S$, is compact, then there exists $E \in D(x) - \{I\}$ with $E(x) = x$.*

PROOF. Suppose that $\varphi(x) = \min_{E \in D(x) - \{I\}} \|Ex - x\|$. Since $D(x) - \{I\}$ is compact, then the minimal of φ exists. Let this minimal be \bar{E} . Therefore

$$\|\bar{E}x - x\| \leq \|Ex - x\|, \quad \forall E \in D(x) - \{I\}. \quad (2.7)$$

Since S is a convex cone with vertex at the origin, then $a\bar{E} + b \in D(x) - \{I\}$ and hence

$$\|\bar{E}x - x\| \leq \|a\bar{E}x + bx - x\|, \quad \forall a, b \geq 0, (a, b) \neq 0. \quad (2.8)$$

Then for $a = b$ we have

$$\|\bar{E}x - x\| \leq \|a(\bar{E}x - x) + (2a - 1)x\| \leq a\|\bar{E}x - x\| + (2a - 1)\|x\| \quad (2.9)$$

and hence

$$\|\bar{E}x - x\| \leq \frac{2a - 1}{1 - a} \|x\|, \quad \forall a \geq 0, \quad (2.10)$$

which implies $\bar{E}x = x$ and hence the result. \square

3. E-stability set of a side. Denote $\rho(I)$ the side in the set M which is defined as

$$\rho(I) = \{x \in \mathbb{R}^n : g_i(x) = 0, i \in I = \{1, 2, \dots, r\}, g_i(x) < 0, 1 \notin I\}. \quad (3.1)$$

DEFINITION 3.1. An E -stability set of a side $\rho(I)$ is denoted by $H(\bar{x}, I)$, $\bar{x} \in S$, and is defined as

$$H(\bar{x}, I) = \{E \in D(\bar{x}) : E(\bar{x}) \in S \cap \rho(I)\}. \quad (3.2)$$

THEOREM 3.2. *If $I_1 \neq I_2$, then $H(\bar{x}, I_1) \cap H(\bar{x}, I_2) = \emptyset$.*

PROOF. Let $\hat{E} \in H(\bar{x}, I_1) \cap H(\bar{x}, I_2)$, then $\hat{E}(\bar{x}) \in S \cap \rho(I_1)$ and $\hat{E}(\bar{x}) \in S \cap \rho(I_2)$, that is,

$$\begin{aligned} g_i(\hat{E}\bar{x}) &= 0, & i \in I_1, & & g_i(\hat{E}\bar{x}) < 0, & i \notin I_1, \\ g_i(\hat{E}\bar{x}) &= 0, & i \in I_2, & & g_i(\hat{E}\bar{x}) < 0, & i \notin I_2. \end{aligned} \quad (3.3)$$

Therefore $g_s(\hat{E}\bar{x}) = 0$, $g_s(\hat{E}\bar{x}) < 0$ for at least one $S \in \{1, 2, \dots, r\}$ which is a contradiction. Hence the result. \square

THEOREM 3.3. *If S and $\rho(I)$ are closed, then the set $H(x, I)$ with $x \in S$ is closed.*

PROOF. Let E^n be a sequence in $H(x, I)$ and E^n tends to E^0 as n tends to infinity, then $E^n(x) \in S \cap \rho(I)$. Since S and $\rho(I)$ are closed, then $E^0(x) \in S \cap \rho(I)$. So $E^0 \in H(x, I)$. Hence the result. \square

THEOREM 3.4. *Let $f, g_i, i = 1, 2, \dots, m$ be convex functions and $\rho(I)$ be a convex set, then $H(x, I)$ is convex.*

PROOF. Let $E_1, E_2 \in H(x, I)$, then $E_1, E_2 \in S \cap \rho(I)$. Since f and $g_i, i = 1, 2, \dots, m$, are convex, then S is convex and hence $S \cap \rho(I)$ is convex since $\rho(I)$ is convex. Thus $\lambda E_1 + (1 - \lambda)E_2 \in S \cap \rho(I), 0 \leq \lambda \leq 1$. Therefore $H(x, I)$ is convex. \square

DEFINITION 3.5. Let $M \subseteq \mathbb{R}^n$ be locally arcwise connected at $x^* \in \bar{M}$ (closure of M). A vector $e \in \mathbb{R}^n$ is said to be tangent to M at x^* if and only if there exists a positive number ν and a continuous function $\delta x(\cdot) : (0, \nu) \rightarrow \mathbb{R}^n$ such that

- (i) $x^* + \alpha \delta x(\alpha) \in M$ for all $\alpha \in (0, \nu)$,
- (ii) $\delta x(\alpha) \rightarrow e$ as $\alpha \rightarrow 0$.

DEFINITION 3.6. For $M \subseteq \mathbb{R}^n$, with $x^* \in \bar{M}$, the set of points

$$T = \{e \in \mathbb{R}^n : e \text{ tangent to } M \text{ at } x^*\} \quad (3.4)$$

is called the tangent cone to M at x^* . (For more details see [7].)

THEOREM 3.7. *Let T be a tangent cone of $\rho(I)$ and f convex and of class C^1 . The sufficient condition to have $E \in H(x, I)$ with $x \in S$ is $(\partial f(Ex)/\partial x)e > 0$ for all nonzero vectors $e \in T$.*

PROOF. Let $(\partial f(Ex)/\partial x)e > 0$ for all nonzero vectors $e \in T$, then $E(x)$ is a proper local minimum of f . Since f is convex, then $E(x)$ is a global minimum for f on $\rho(I)$, that is, $E(x) \in S \cap \rho(I)$. Hence $E \in H(x, I)$. \square

REFERENCES

- [1] B. Bank, J. Guddat, D. Klatte, B. Kummer, and K. Tammer, *Nonlinear Parametric Optimization*, Birkhäuser Verlag, Basel, 1983. [MR 84i:90147](#).
- [2] V. Jeyakumr and X. Q. Yang, *Convex composite multiobjective non smooth programming*, *Math. Programming* **59** (1993), 325–343.
- [3] M. S. A. Osman, *Solvability of a convex program with parameters in the objective function*, *Proceeding of the Third Annual Operations Research Conf. (Egypt)*, vol. 3, Zagazig University, 1976.
- [4] ———, *Qualitative analysis of basic notions in parametric convex programming. I. Parameters in the constraints*, *Apl. Mat.* **22** (1977), no. 5, 318–332. [MR 56 #7993](#). [Zbl 383.90097](#).
- [5] ———, *Qualitative analysis of basic notions in parametric convex programming. II. Parameters in the objective function*, *Apl. Mat.* **22** (1977), no. 5, 333–348. [MR 56 #7994](#). [Zbl 383.90098](#).
- [6] Y. Sawaragi, H. Nakayama, and T. Tanino, *Theory of Multiobjective Optimization*, Academic Press, Florida, 1985. [MR 87b:90129](#). [Zbl 566.90053](#).
- [7] L. V. Thomas and J. G. Walter, *Optimality in Parametric Systems*, John Wiley & Sons, New York, 1981. [MR 83d:90001](#).

- [8] X. Q. Yang, *Second-order global optimality conditions for convex composite optimization*, Math. Programming **81** (1998), no. 3, Ser. A, 327–347. [MR 2000g:90106](#).
- [9] E. A. Youness, *E-convex sets, E-convex functions, and E-convex programming*, J. Optim. Theory Appl. **102** (1999), no. 2, 439–450. [MR 2000d:90057](#). [Zbl 937.90082](#).

EBRAHIM A. YOUNESS: DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCE, TANTA UNIVERSITY, TANTA, EGYPT



Hindawi

Submit your manuscripts at
<http://www.hindawi.com>

