

Technical University of Denmark



## Simulation and measurement of bacterial growth on low soluble phenanthrene substrate

Trapp, Stefan; Adam, I.; Rein, A.; Miltner, A.; da Socta Fulgencio, A.; Kaestner, M.

*Published in:*  
Science Across Bridges, Borders and Boundaries

*Publication date:*  
2014

*Document Version*  
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

*Citation (APA):*  
Trapp, S., Adam, I., Rein, A., Miltner, A., da Socta Fulgencio, A., & Kaestner, M. (2014). Simulation and measurement of bacterial growth on low soluble phenanthrene substrate. In Science Across Bridges, Borders and Boundaries: Programme Book Basel, Switzerland: SETAC-Europe.

## DTU Library

Technical Information Center of Denmark

---

### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

**TU228 Simulation and measurement of bacterial growth on low soluble phenanthrene substrate** S. Trapp, Danmark Tekniske

Universitet / DTU Environment; I. Adam, Helmholtz Center UFZ Leipzig / Environmental Biotechnology; A. Rein, TU Munich; A. Miltner, Helmholtz Center UFZ / Environmental Biotechnology; A. daCostaFulgencio, Helmholtz Center UFZ; M. Kaestner, Helmholtz Centre for Environmental Research UFZ / Dept Environmental Biotechnology. The metabolism of low soluble substrate is limited by dissolution and substrate availability and can hardly be determined in a common chemostat. We developed a numerical model that calculates simultaneously dissolution kinetics of such substrate, metabolism (Michaelis-Menten kinetics) and microbial growth (Monod kinetics with decay term) for the dynamic case. Experiments on the degradation of phenanthrene by and the growth of the three degrader strains *Novosphingobium pentaromativorans* US6-1, *Sphingomonas* sp. EPA505 and *Sphingobium yanoikuyae* B1 were used to determine kinetic parameters as input for the model. Phenanthrene (Pht) in acetone was added to 10 mL test vials. The nominal initial concentration of the suspensions was 10, 25, 50, 100, 200 and 400 mg/L. Pht was present as slowly dissolving microcrystals. This provided non-limiting conditions for the growth of the degrader strains over several days. Total Pht concentration and protein were tracked over 6 to 12 days. In all replicates, Pht was completely metabolized, and biomass increased rapidly, more at higher initial concentration, but decayed when Pht was depleted. The model was fitted to the test result in order to determine the rates of dissolution, metabolism and growth. The outcome shows that the three bacterial strains have similar efficiency, with  $v_{max}$ -values of 12 to 18 g bacteria dw / g substrate / d, yields of 0.21 g/g, maximum growth rates  $u_{max}$  of 2.5 to 3.8 1/d and decay rates of 0.04 to 0.05 1/d. Simulations with the model show that i) retainment in crystals, NAPL or by sequestration compete with biodegradation, since molecules remaining non-dissolved cannot be degraded; ii) the conditions for bacterial growth (i.e. dissolution flux and resulting chemical activity of substrate) are more relevant for the final state of the system (both concerning number of degraders and time-course of substrate concentrations) than initial biomass; and iii) the desorption flux regulates the turnover when the substrate source is in solid state or present in sequestered (aged) systems. Provided the equations describe correctly the kinetics, the calibrated model can be used to simulate bioavailability, biodegradation, persistence and treatment options in real systems, such as PAH-contaminated soils.