

Technical University of Denmark



Climate friendly dietary guidelines

Trolle, Ellen; Mogensen, Lisbeth; Thorsen, Anne Vibeke; Jørgensen, Michael Søgaaard

Published in:

Proceedings of the 9th International Conference on Life Cycle Assessment in the Agri-Food Sector

Publication date:

2014

[Link back to DTU Orbit](#)

Citation (APA):

Trolle, E., Mogensen, L., Thorsen, A. V., & Jørgensen, M. S. (2014). Climate friendly dietary guidelines. In R. Schenck, & D. Huizenga (Eds.), Proceedings of the 9th International Conference on Life Cycle Assessment in the Agri-Food Sector: LCA Food 2014 (pp. 1344-1351). ACLCA.

DTU Library

Technical Information Center of Denmark

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Climate friendly dietary guidelines

Ellen Trolle¹, Lisbeth Mogensen², Michael Søgaaard Jørgensen³, Anne Vibeke Thorsen¹

- 1) Division of Nutrition, National Food Institute, Technical University of Denmark
- 2) Department of Agroecology, Aarhus University, Denmark
- 3) Department of Development and Planning, Aalborg University, Denmark

Corresponding author. Ellen Trolle E-mail: eltr@food.dtu.dk

ABSTRACT

The aim of this study was to investigate how the present Danish diet could be changed in a climate friendly direction that follows the recommendations of a healthy diet.

The carbon footprint (CF) of an average Danish diet was calculated and compared to CF of a recommended healthy diet by 1) modifying the average diet according to the Danish food based dietary guidelines, 2) and adjusting to ensure an iso-energy content and a nutrient content according to the Nordic Nutrient Recommendations. Afterwards the healthy diet were changed further to reduce CF.

CF from the diet was reduced by 4%, if the healthy diet was eaten instead of the average current diet. However, if the diet was climate optimized by choosing foods with a low CF within the food groups; meat, vegetables and fruit, CF of this diet may be reduced by 23 % compared to CF of the average diet.

Keyword: Danish average diet, carbon footprint, Danish food based dietary guidelines

1. Introduction

In recent years focus has been on emission of greenhouse gases (GHG) from the food production. It has been estimated that the food sector in the developed countries contributes with up to 30 % of total GHG emission (Tukker et al. 2009). Several Climate Summits organized by United Nation (latest COP19 was held in Warsaw, Poland from 11 to 23 November 2013) have been held in order to address the climate change at an international level, which unfortunately didn't result in specific targets or actions (UN 2009; UN 2010).

The special thing about food production is that in addition to contributing to emission of CO₂ from fossil energy consumption it also contributes with the so-called non-energy-related GHG emission – in form of the nitrox oxide (N₂O) emissions related to use of fertilizers, deforestation (CO₂), and methane (CH₄) from ruminant digestion.

The diet and consequently the related food production contribute to the GHG emission. The different stages of each food items life cycle contribute to the GHG emission: the primary agricultural food production, food processing, as well as during transport and storage of food (Carlsson-Kanyama et al. 2003; Anonymous 2009; Garnett 2008; Nielsen et al. 2003). Finally, also the cooking at home contributes to the GHG.

The climate impact can be assessed from different perspectives; a production perspective i.e. the burden of the total food produced in Denmark inclusive emissions from food that is exported to other countries, or alternative from a consumer perspective – i.e. the burden of the food consumed in Denmark exclusive emissions from food that is exported and inclusive emissions from food that is produced outside Denmark. Seen from the production perspective, Denmark's agricultural production was estimated to represent 16 % of the total Danish emission of GHG (Olesen 2008). The climate impact of the Danish food consumption from

the consumer perspective was calculated to be around 2.8 tons CO₂-eqv / person/year – equivalent to approximant 15.4 mill. tons CO₂ for the total Danish population and thus approximate 25 % of all GHG from The Danes' total consumption according to IDA's Climate Plan 2050 (Anonymous 2009).

Different types of foods contribute to different degrees to the climate impact. There are big differences between the level of GHG emission from different food groups e.g. between animal products such as meat and cheese and vegetable products such as vegetables, flour and grain. Also within the various food groups there are differences, e.g. between different types of fish or meat depending on the way the products are produced. (Anonymous 2009; Olesen 2009).

The amount of GHG emitted by a produced food product is called the food's carbon footprint (CF). CF or global warming is one among several impact categories; acidification, nutrient enrichment, photochemical smog and land use etc. The CF is calculated by a life-cycle assessment (LCA), which includes GHG emission from the foods whole life cycle: agriculture, horticulture or fishing, including emissions related to the production of inputs such as fertilizers, processing, transportation and storage of food products until the food products are placed on the shelf in the supermarket. The CF of the preparation at households is typical not included. (Anonymous 2009; Mogensen et al. 2009a; Garnett 2008; Nielsen et al. 2003). Some LCA calculations also include estimations of GHG contribution derived from food waste (Mogensen et al, 2011; Anonymous 2009). Differences in the used LCA calculation methods may complicate comparisons between CF from different sources. Furthermore, production methods and energy resources change over time, and therefore influence the results of calculations.

This study (Thorsen et al 2012) was based on Danish food based dietary guidelines from 2005 (Astrup et al. 2005) including currently updates. They aim at increasing the intake of fruit and vegetables, bread and cereals (coarse or wholegrain) and fish, and a decrease in intake of fat from dairy and meat products and of sugar containing products. In 2013 the scientific evidence of the Danish food based dietary guidelines was updated. The main conclusions were that the increase of fruit and vegetables was maintained, fish intake was increased, an increased intake of wholegrain cereals was specified and the decrease in fat from dairy and meat products was emphasized. Furthermore the intake of red meat (cow, sheep, pig) and the sugar containing foods were restricted, especially sugar containing beverages.

1.1 Objectives

The aim of this study was to investigate how the present Danish diet could be changed in a climate friendly direction and at the same time following the recommendations of a healthy diet.

2. Methods

The carbon footprint (CF) of the average Danish diet estimated from the National Danish Survey on Diet and Physical Activity from 2003 2008, including 3354 adults, 18-75 years of age, hereof 47% men (Pedersen et al. 2010) was estimated. CF of an average diet was compared with CF of a diet that follows the Danish dietary recommendations. This diet was obtained by 1) modifying the average diet by scaling the food groups according to the Danish food based dietary guidelines (Astrup et al. 2005), and 2) adjusting to ensure an iso-energy content of the diets and a nutrient content that follows the Nordic Nutrient Recommendations (Nordic Council of Ministries 2004). The healthy diet was designed by using a modeling tool developed from the Danish nutrient calculating system GIES (Biltoft-Jensen et al 2008).

To assess the impact of food production on global warming the CF from each food item was calculated by use of life cycle assessment (LCA) (ISO, 2006a; ISO, 2006b). The LCA method implies that all emissions of GHG from cradle to grave are included. I.e. all GHG emissions from production and transportation at the farm, but also the GHG emissions related to processes after the food leave the farm are included. The CF from individual foods i.e. is expressed in CO₂ equivalent per unit of food produced, and the functional unit (FU) is one kg food.

The CF of food take into account the contribution from GHG emissions related to food waste from the whole food chain, from production to retail and households. As there is no Danish estimations of food waste in households, waste is estimated from a detailed English report about food waste on individual food groups, on average about 20% of purchased food ends up as edible waste, i.e. waste that could have been avoided (WRAP 2008).

The healthy diet was investigated further with the aim to reduce CF of the diet by using the “hot spot” approach. Hot spot food groups were in focus, i.e. food groups where it is easy and convenient to change between different food products in the group in order to minimize the CF. Hot spot analyzes are interesting because in that way you could help the consumer to choose more climate friendly food, while taking into account also the Nordic recommendations (Nordic Council of Ministries 2004) and the 8 Danish dietary guidelines (Astrup et al. 2005).

When the hot spot perspective is related to the dietary guidelines it is important to focus on food groups that have a high food consumption (large amount of food) and/or include foods having a large CF (high CO₂/kg food). The next step is to assess how foods of the hot spot food groups could be exchanged with similar foods within the group with lower CF.

Finally, the Danish food based dietary guidelines from 2013 (Danish dietary guidelines 2013; Tetens et al. 2013) were investigated and if relevant, complementary advises related to improving the climate impact were added.

3. Results

The estimated CF from the Danish’ average diet (2003-2008) and a modeled recommended diet that meets the dietary guidelines and the nutritional recommendations are shown in Table 1.

Table 1 comprises 14 different food groups. Both diets are scaled to an energy intake of 10 MJ/day. These food intakes also give rise to food waste in retail (2 %, but 6 % for fruits and vegetables), processing (2 %) and household (3 %). Regarding fruit and vegetables also a 20 % peel waste (inevitable waste) is included. Table 1 shows an estimate of how much the changes from the average diet of the modeled recommended diet would benefit the climate. It is seen that the largest differences obtained by a reduction of the amount of meat with approximate 50 g. This is to some extent cancelled by the contribution from the foods replacing the meat such as the food groups of fruit, vegetables, milk and fish which according to the Danish food based dietary guidelines 2005.

The calculation shows that CF from the diet was reduced by 4%, if the healthy diet was eaten instead of the average current diet. The largest reduction is a result of reduced intake of meat and beverages (beer and wine). The GHG contribution of beverages (excluding milk and juice) has declined by almost 40 %. Overall the recommended diet has a CF that was approximate 4 % lower than the current average diet

Table 1. The carbon footprint (CF) for an average Danish diet and for a modelled recommended diet (that fulfills both the dietary guidelines and the Nordic recommendations). Both diets are scaled to a daily energy intake of 10 MJ.

Food group	CF, kg CO ₂ /kg food	Intake g/person/day		Food waste, % ⁶⁾	CF form the diet G CO ₂ /person/day	
		Present average diet	“Modelled recommended diet”		Present average diet	“Modelled recommended diet”
Milk, dairy products	1,2	359	500	(7)	462	644
Cheese	11,3	38	25	(7)	455	303
Bread, rice, pasta	0,8-3,3	236	274	(37)	403	480
Vegetables	0,1 – 2,9	186	304	(38)	381	567
Potatoes	0,2	113	192	(38)	31	52
Fruit (ex. Juice)	0,4	245	271	(45)	230	260
Juice	1,0	80	50	(21)	102	63
Meat	3,6 – 19,4 ²⁾	121 ³⁾	87 ⁴⁾	(39) ⁷⁾	1599	1277
Poultry	3,4	27	31	(39) ⁷⁾	149	171
Egg	2,0	19	25	(22)	49	63
Fish	1,8 – 10,5	25	42	(18) ⁸⁾	170	292
Fats	5,1	38	32	(7)	219	173
Sugar and candy	1,0	36	23	(22)	41	29
Beverages	0,02 – 2,1	2273	1955	(7)	698	417
Total diet ¹⁾					4986	4790
					(100)	(96)
Recommended and climate friendly diet ^{1,5)}						3864 (77)

¹⁾ Beverages are included, figures in brackets: percentages related to average diet.

²⁾ Climate footprint per kilo carcass – Amount of carcass behind an intake is calculated by using a factor 1,47 for beef, 1,33 for pork and 1,38 for poultry (chicken)

³⁾ Type of meat in the present average diet: Men: 135 g meat/day: 25 % beef, 75 % pork. Women: 106 g meat/day: 28 % beef 72 % pork.

⁴⁾ Meat in the recommended diet: Men: 92 g meat/day: 30 % beef, 70 % pork. Women: 81 g meat/day: 33 % beef, 67 % pork.

⁵⁾ Climate friendly diet: The only difference from ”recommended” diet is: within the food group: fruit, vegetables and meat a climate-friendly solution is chosen e.g. For the food group vegetables carrot is chosen, for the food group fruit apple is chosen and for the food group meat the reduction is done for beef and then for pork. a reduction

⁶⁾ A larger food production is need than the food intake figures show since food waste is found in all the chains in the food . Edible food waste in household and retail: (2 %, but 6 % for fruit and vegetables) + processing (2 %). Total edible food waste is noted in parenthesis, inclusive 20 % peel/skin (inevitable waste) for fruit, vegetables and potatoes.

⁷⁾ Incl. calculation from carcass to meat.

⁸⁾ All fish products are calculated without bone like fillet and peeled shrimp

In table 2 we take a closer look at the CF for the hot spot food groups: meat, vegetables, fruit and fish.

From Table 2 it can be seen that it is possible within the different food groups to reduce the contribution to CF by choosing foods with a lower CF. In order to choose climate friendly foods it is better to choose e.g. poultry or pork instead of beef, and fruit or vegetables in season instead of food produced in a greenhouse or imported. Regarding fish a climate friendly choice would be herring or mussels instead of flatfish or shrimps.

In addition the diet was optimized to reduce CF of the diet by choosing foods with a low CF within the food groups; meat, vegetables and fruit, CF of this diet may be reduced by 23 % compared to CF of the average diet.

Table 2: Effect on the contribution to carbon footprint (CF) from different food groups dependent on choice of food item within food group

Food group	Foods item	“Modeled” recommended daily intake in gram ²⁾	Food waste, % ⁵⁾	CF from the food items kg CO ₂ /person/day	Contribution to CF from the production of the daily intake of food ¹⁾ , gram CO ₂ /day
Meat and meat products	Average diet – mix of food items within the food group	87	(39)	9,2 ⁴⁾	1277
	Beef	87	(43)	19,4 ³⁾	2966
	Pork	87	(37)	3,6 ³⁾	498
	Chicken, fresh	87	(39)	3,1 ³⁾	445
	Chicken, frozen	87	(39)	3,7 ³⁾	531
Vegetables (ex. potatoes)	Average diet – mix of food items within the food group	304	(40)	1,32 ⁴⁾	567
	Carrot	304	(40)	0,122	62
	Onion	304	(40)	0,382	195
	Greenhouse vegetables	304	(26)	2,7	1099
Fruit	Average diet – mix of food items within the food group	271	(45)	0,52 ⁴⁾	260
	Orange	271	(45)	0,7	347
	Banana	271	(45)	0,5	248
	Nuts, almonds	271	(32)	0,88	350
	Danish apple, pear	271	(45)	0,1	50
	Imported apple, pear	271	(45)	0,4	198
	Average diet – mix of food items within the food group	42	(18)	5,7 ⁴⁾	292
Fish and fish products	Average diet – mix of food items within the food group	42	(18)	1,8	90
	Herring, fillet, peeled, frozen	42	(18)	10,5	528
	Shrimp, frozen, peeled	42	(18)	3,2	161
	Codfish, fillet, frozen	42	(18)	7,8	392
	Flatfish, fillet, frozen	42	(18)		

¹⁾ Calculation of CF is based on produced amount of feed, taking into account food waste.

²⁾ Food intake is an average for men and women.

³⁾ CF of meat is given as CF per kg carcass, needed amount of carcass per kg meat intake: 1.47 for beef 1.33 for pork and 1.38 for poultry.

⁴⁾ Average CF for all type of meat in the meet group, weighted according to distribution between different types of meat, accordingly average CF for the other food groups

⁵⁾ Food waste in household, retail and processing. Includes also bones etc. (from carcass to intake of meat).

In the climate friendly diet, we have reduced the meat intake by 50 g/day, removed the beef and then reduced the pork. The intake of vegetables was increased with 300 g/day, and the vegetables with the lowest CF (such as carrots) was chosen, and regarding fruit the amount of fruit was increased by 50 g/day and fruit with the lowest CF (such as Danish apple) was chosen. The results are shown in Table 1 above and in footnote 5. The climate friendly diet did not include a climate optimized choice from the following food groups; fish and beverages.

4. Discussion

The present study showed that the climate impacts from human food consumption can be reduced by conscious food choices. Beside that minimizing food waste and choosing a more sustainable food production in relation to agriculture could reduce CF from food production further

In this study, the most recent dietary data were used for rough calculations of the climate contribution of a recommended diet compared to the current Danish diet. According to our calculation the climate contribution from the diet will be 4 % reduced, if the recommended diet is eaten instead of the average current diet. A

saving in carbon footprint in the order of 4% CO₂-eq is so small compared to the uncertainty of the data included that is not necessarily a real saving.

In addition to eating a recommended diet people would optimize their diet in a more climate friendly way by choosing foods with a low carbon footprint, especially in the food groups; meat, vegetables and fruit. Our calculations show that the CF of such a climate friendly diet including beverages would be reduced by 23 % compared to CF contribution of the average diet. Thus, a climate optimized recommended diet would provide a significant reduction of the CF (23%) as compared to a recommended diet (4%). The number (23 %) is not essential, it can be less or more, but the savings are significant compared to the recommended diet. In real life the savings from the food groups meat, fruit and vegetables probably would be less than estimated here, but on the other hand optimized choices within the other food groups is expected to provide further savings. Thus there is a great potential for reducing CF of the diet by choosing climate friendly within a recommended diet. If households further reduce their food waste, it will have a major effect on climate impacts. The total food waste in households is estimated to be around 20%, accounting for 12.5% of the CF of food production.

The calculations include food waste from the food chain, from production to retail and households. However, the calculations are based on English data from investigating food waste in households (WRAP 2008). More recent data from Danish households are needed to validate this part of the calculation.

Other studies (Mogensen et al. 2009b; Saxe et al. 2006) find that a diet following the dietary guidelines will have a slightly lesser impact on the climate all other things being equal. Another study found that the GHG emission was 27 % lower when the diet (the New Nordic) was climate optimized by choosing either less beef or by substituting all meat with legumes, dairy products and eggs (Saxe et al. 2013).

The present study is an approximation and is conducted to get an overview of the CF from the overall diet. More accurate calculations of the climate-optimized diet require more in-depth calculations that are beyond the aim of this study. For instance in the climate-friendly diet the milk intake is set to be 500 g/day, and although this could be reduced to 250 ml in accordance with the Danish dietary guidelines from 2013 (Tetens et al. 2013), it might not be in accordance with removing all of the beef.

Fruits, vegetables, cereals and potatoes, which according to the dietary guidelines should be by far the largest part of the diet, are all low in climate impact, while meat and cheese are generally high. Also vegetable oil should replace butter and hard margarines, vegetable oils generally have a lower CF. Stimulants as sweet and alcoholic drinks, sweets and cakes, which should decrease in the Danish diet, probably have a rather high CF, but the data concerning CF of this food group is weak and should be improved considerably. An increased intake of fish would by all means increase the climate impact, but may be limited by a conscious choice of fish products. Eating the recommended diet would change the diet in the direction of lower fat and higher fiber content, e.g. by reducing the intake of red meat and cheese and instead eat more coarse vegetables and fruit, bread and grains. By further choosing foods with a low carbon footprint whenever possible the climate impacts of the food consumption is reduced significantly. These qualitative advices was in line with guideline development in Sweden and the Netherlands (Fogelberg 2008; Health Council of the Netherlands 2011)

5. Conclusion

It was concluded that there is potential synergy between a healthier diet and a more climate friendly diet. There seems evidence to complement the Danish dietary guidelines 2013 with the following advice to reduce the CF from the diet:

- Eat a varied diet, not too much, and be physical active
- Eat fruit and many vegetables - *preferably free-range and in season*
- Eat more fish - *choose the climate friendly fish; herring and mussels*
- Choose whole grain
- Choose low-fat meat and meat products –*choose pork and poultry rather than beef and lamb*
- Choose low-fat milk and milk products - *restrict intake of cheese*
- Eat less saturated fat - *choose vegetable oils rather than animal fat*
- Eat foods with less salt
- Eat less sugar - from soft drinks, sweets and cakes

- Drink water - rather than sweet and alcoholic drinks
Avoid overeating and waste - will also reduce food production.

6. References

- Anonym (2009) IDA's klimaplan 2050. Projekt Future Climate. Copenhagen
- Astrup A, Andersen NL, Stender S, Trolle E (2005) Kostrådene 2005 (Dietary Guidelines 2005). Danish Institute for Food and Veterinary Research and Danish Nutritional Council. Copenhagen
- Biltoft-Jensen A, Trolle E, Christensen T, Ygil KH, Fagt S, Matthiessen J et al. (2008) Development of a recommended food intake pattern for healthy Danish adolescents consistent with the Danish dietary guidelines, nutrient recommendations and national food preferences. *J Hum Nutr Diet* 21; 451-463.
- Carlsson-Kanyama A and Engström R (2003) Fakta om maten och miljön: Konsumtionstrender, miljöpåverkan och livscykelanalyser. Naturvårdsverket rapport nr. 5348. Stockholm
- Danish dietary guidelines 2013
<http://www.foedevarestyrelsen.dk/Publikationer/Alle%20publikationer/2013200.pdf>
- Fogelberg CL (2008) På väg mot miljöanpassade kostråd - vetenskapligt underlag inför miljökonsekvensanalysen av Livsmedelsverkets kostråd. Livsmedelsverket. Uppsala
- Garnett T (2008) Cooking up a storm. Food, greenhouse gas emissions and our changing climate. . UK: Food climate Research Network. Centre for Environmental Strategy. University of Surrey.
- Health Council of the Netherlands (2011) Guidelines for a healthy diet: the ecological perspective, publication no.2011/08E. Hague: Health Council of the Netherlands.
- ISO (2006a) ISO 14040: Environmental management - Life cycle assessment - Principles and framework. International Organisation for Standardisation. Geneva
- ISO (2006b) ISO 14044: Environmental management - Life cycle assessment - Requirements and guidelines. International Organisation for Standardisation. Geneva
- Ministeriet for Fødevarer LOF (2009) Klima på bordet. Fødevareministeriets klimakogebog. Fødevareministeriet. Copenhagen
- Mogensen L, Hermansen JE, Knudsen MT (2011) Notat til Fødevareministeriet. Madspild i fødevarereproduktionen - fra primær produktion til detailled. Aarhus Universitet, Det Jordbrugsvidenskabelige Fakultet, Institut for Jordbrugsproduktion og Miljø.
- Mogensen L, Kidmose U, Hermansen JE (2009a) Baggrundsnotat til Fødevareministeriet: Fødevarers klimaaftryk, sammenhængene mellem kostpyramiden og klimapyramiden, samt opfang og effekt af fødevarerspild. Aarhus Universitet, Det Jordbrugsvidenskabelige Fakultet, Institut for Jordbrugsproduktion og Miljø og Institut for Fødevarekvalitet.
- Mogensen L, Knudsen MT, Hermansen JE (2009b) Notat til Fødevareministeriet vedrørende: Beregning af klimaaftryk for middagsretter til klimakogebog. Aarhus Universitet, Det Jordbrugsvidenskabelige Fakultet, Institut for Jordbrugsproduktion og Miljø og Institut for Fødevarekvalitet. Aarhus
- Nielsen PH, Nielsen AM, Weidema BP, Dalgaard R, Halberg N (2003) LCA food database. Available online, 31/8 2003 at www.lcafood.dk/database.
- Nordic Council of Ministers (2004) Nordic Nutrition Recommendations 2004, 4th edition. Integrating nutrition and physical activity. Nord 2004:13. Nordic Council of Ministers. Copenhagen
- Olesen JE (2008) Climate-friendly food. In: Momentum: Clima - the agricultural view. Jord og Viden tema, nr. 4; 4-8.
- Olesen JE (2009) Theme 3: Agriculture and climate change. In: Beyond Kyoto, Addressing the challenges of climate change. The 7 Aarhus Statements on climate change. 6. March 2009, Aarhus
- Pedersen AN, Fagt S, Groth M, Christensen T, Biltoft-Jensen A, Matthiessen J, Andersen NL, Kørup K, Hartkopp HB, Ygil KH, Hinsch HJ, Saxholt E, Trolle E (2010) Dietary habits in Denmark 2003-2008. Main results (in Danish). National Food Institute, Technical University of Denmark, Søborg
- Saxe H, Jensen RB, Petersen ML (2006) Fødevarers miljøeffekter. Det politiske ansvar og det personlige valg. Institut for miljøvurdering. Copenhagen
- Saxe H, Larsen TM, Mogensen L (2013) The global warming potential of two healthy Nordic diets compared with the average Danish diet. *Climate Change* (2013) 116; 249-262

Tetens I, Andersen LB, Astrup A, Gondolf UH, Hermansen K, Jakobsen MU, Knudsen VK, Mejbom H, Schwarz P, Tjønneland A, Trolle E (2013) Evidensgrundlaget for danske råd om kost og fysisk aktivitet (Danish with English summary). National Food Institute, Technical University of Denmark, Søborg

Thorsen AV, Mogensen L, Jørgensen MS, Trolle E 2012. Klimaorienterede kostråd. DTU. National Food Institute, Technical University of Denmark, Søborg

Tukker A, Bausch-Goldbohm S, Verheijden M, Koning E, Kleijn R, Wolf O, Dominguez IP (2009)

UN (2009): The United Nations Climate Change Conference in Copenhagen, 7-9 December 2009.

UN (2010): The United Nations Climate Change Conference in Cancun, COP 16/ CMP 6, 26 November 2010.

WRAP (2008): The food we waste. The Waste Resources Action Programme WRAP Report.