



The effect of production type and antimicrobial usage on the occurrence of tetracycline resistant *E. coli* in danish slaughter pig farms

Struve, Tina; Vigre, Håkan; Wingstrand, Anne; Sørensen, Anna Irene Vedel; Lundsby, Katrine Løvenbalk; Emborg, Hanne-Dorthe

Publication date:
2009

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):
Struve, T., Vigre, H., Wingstrand, A., Sørensen, A. I. V., Lundsby, K. L., & Emborg, H-D. (2009). The effect of production type and antimicrobial usage on the occurrence of tetracycline resistant *E. coli* in danish slaughter pig farms. Abstract from 12th Conference of the International Society for Veterinary Epidemiology and Economics, Durban, South Africa.

DTU Library

Technical Information Center of Denmark

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Reference: ISVEE/ 394

Date: 2009-01-29

Abstract type: Either

Abstract theme: Food safety, environmental epidemiology and occupational health

Title:

The effect of production type and antimicrobial usage on the occurrence of Tetracycline resistant *E. coli* in Danish slaughter pig farms.

Authors and Affiliation:

T. Struve (1), Vigre, H. (1), Wingstrand, A. (1), Sørensen, A.I.V. (1), Lundsby, K.L. (1), Emborg, H-D. (1)

1: National Food Institute, Technical University of Denmark, Søborg, Denmark

The Qualysafe project was initiated in 2007 to support and strengthen the sustainable production systems in Danish food production. One of the objectives of the epidemiological investigation was to find new methods to improve food safety in conventional as well as in alternative pig production systems. At nine different slaughterhouses 1500 ceacum samples were collected from slaughter pigs originating from 226 farms. One thousand samples were analyzed and one *E. coli* isolate per sample was susceptibility tested to Tetracycline. Data on management practice and health status at farm level was collected through telephone interviews. Data on antimicrobial consumption at farm level was collected from the Danish surveillance program VetStat. In total, 99 farms used Tetracycline and participated in the interview and from these farms, 411 isolates were available. We assumed that production type (organic, free range and conventional farms) was a risk factor for occurrence of antimicrobial resistance and Tetracycline usage was regarded as an intervening factor between production type and occurrence of antimicrobial resistance. Therefore, the effect of production type and Tetracycline usage was estimated in two separate models using logistic regression, taking into account the correlation of results obtained from the same farm. Among the 411 isolates, 129 was found resistant to Tetracycline (Organic: 10%, Free Range: 27 % Conventional: 39 %). Differences was seen in the consumption pattern among the farm types, with the Organic having the lowest consumption (0.14 doses /annually produced slaughter pig) while Free Range had the highest consumption (0,85 doses/annually produced slaughter pig) and Conventional farms was in between (0.67 doses/annually produced slaughter pig). The effect of Tetracycline usage was estimated using a quadratic polynomial. This showed a significant effect of Tetracycline consumption on the occurrence of Tetracycline resistance, where the occurrence of resistance increased by increased antimicrobial usage. Production type had significant effect on the occurrence of antimicrobial resistance, with the lowest occurrence in Organic production and the highest in Conventional production. When analyzing the effect of production type and usage of antimicrobials on occurrence of resistance in the same model, the usage of antimicrobials was not found to have an effect. This lack of effect can be explained by the fact that Tetracycline, at least partly, is an intervening factor between production type and the consumption. The results from our analysis showed the importance of correctly classifying different risk factors as causal and intervening factors before analyzing the effect of potential risk factors on the occurrence of antimicrobial resistance in animal production.

Presenting Author: Struve

Corresponding Author: Struve

E-mail: tstr@food.dtu.dk