

Spatial variability of cyanobacterial community composition in Sanya Bay as determined by DGGE fingerprinting and multivariate analysis

LING Juan^{1,2,3,4}, ZHANG YanYing^{1,3}, DONG JunDe^{1,3*}, WANG YouShao², HUANG Hui^{1,3}, CHEN Lei^{1,3,4}, HUANG XiaoFang^{1,3,4}, LONG LiJuan¹ & ZHANG Si¹

¹ Key Laboratory of Marine Bio-resources Sustainable Utilization, South China Sea Institute of Oceanology, Chinese Academy of Sciences, Guangzhou 510301, China;

² State Key Laboratory of Tropical Oceanography, South China Sea Institute of Oceanology, Chinese Academy of Sciences, Guangzhou 510301, China;

³ National Experiment Station of Tropical Marine Biology, Chinese Academy of Sciences, Sanya 572000, China;

⁴ Graduate University of Chinese Academy of Sciences, Beijing 100049, China

Received April 25, 2012; accepted June 7, 2012; published online September 6, 2012

The cyanobacterial communities in the surface and bottom waters of Sanya Bay were investigated on April 24 and 25, 2010. Flow cytometry showed that the total cyanobacterial abundance in the surface and bottom layers ranged from 0.7×10^4 to 2.38×10^4 cells mL^{-1} and from 1×10^4 to 1.8×10^4 cells mL^{-1} , respectively. Cyanobacterial diversity was analyzed using a molecular fingerprinting technique called denaturing gradient gel electrophoresis (DGGE), followed by DNA sequencing. The results were then interpreted through multivariate statistical analysis. Differences in the compositions of cyanobacterial communities were observed in the surface and bottom waters at the same station, with some bands obtained from both the surface and bottom layers, whereas some bands were present only in one layer. The predominant cyanobacterial species of the excised DGGE bands were related to *Synechococcus* or *Synechococcus*-like species (56.2%). Other phylogenetic groups identified included *Chroococciopsis* (6.3%), *Cyanobium* (6.3%) and some unclassified *cyanobacteria* (31.2%). A redundancy analysis (RDA) was conducted to reveal the relationships between the cyanobacterial community composition and environmental factors. Analysis results showed that the spatial variations in the cyanobacterial community composition in surface waters was significantly related to chlorophyll *a* (Chl*a*), the biochemical oxygen demand (BOD), nitrate and phosphate ($P < 0.05$). Meanwhile, the spatial variations in the bottom waters was significantly affected by nitrate, nitrite, and phosphate ($P < 0.05$). Environmental parameters could explain 99.3% and 58.3% of the variations in the surface and bottom layers, respectively.

cyanobacterial community composition, PCR-DGGE, *Synechococcus*, redundancy analysis

Citation: Ling J, Zhang Y Y, Dong J D, et al. Spatial variability of cyanobacterial community composition in Sanya Bay as determined by DGGE fingerprinting and multivariate analysis. *Chin Sci Bull*, 2013, 58: 1019–1027, doi: 10.1007/s11434-012-5424-4

Cyanobacteria occupy a wide range of environmental niches in tropical marine regions and are the most diverse photosynthetic bacteria [1]. These organisms are considered to be major N_2 -fixation microorganisms in the open ocean and are essential components of the microbial food web [2,3]. The N_2 fixation of diazotrophic picocyanobacteria is estimated to provide a global rate of 100 Tg a^{-1} in warm oligotrophic

waters [4]. The N_2 -fixation process is also important in providing the nutrient flux that support carbon loss in the deep ocean and limit the concentration of the greenhouse gas, carbon dioxide. Cyanobacteria (e.g. *Lyngbya* sp.) are also the source of bioactive compounds [5] and are capable of degrading organic pollutants, such as polycyclic aromatic hydrocarbons and xenobiotics [6].

Sanya Bay, a typical tropical bay, is located in China's Hainan Island. It consists of several tropical ecosystems

*Corresponding author (email: dongjunde@vip.163.com)

such as coral reefs, mangroves, and seagrass beds. Previous investigations suggested Sanya Bay to be a warm and oligotrophic bay, with nitrogen as the main nutrient that limits primary production [7–9]. Cyanobacteria in Sanya Bay have been proven to be a major component of total phytoplankton biomass and productivity [10]. However, systematic investigations of the cyanobacterial population are rarely conducted, and no genetic information is available. Early studies on cyanobacterial identification in Sanya Bay relied primarily on microscopic analysis and were based on morphological characteristics [11]. These conventional methods for cyanobacterial studies were time-consuming and labor intensive, and may disregard some important picocyanobacterial species because of certain limitations. As a result, polymerase chain reaction-denaturing gradient gel electrophoresis (PCR-DGGE) and DNA sequence analysis were used in the present study to survey the structure of the cyanobacterial communities using natural samples collected from Sanya Bay [12]. In addition, direct multivariate analyses such as redundancy analysis (RDA) were used to study the species-environmental relationships. High-resolution DNA fingerprinting combined with multivariate analysis has been used to investigate the variability in the marine microbial community structure in aquatic ecosystems [13].

This study provided a detailed understanding of cyanobacterial communities in the surface and bottom waters of Sanya Bay. The overall objectives were as follows: (1) to investigate the cyanobacterial community structure and the dominant cyanobacterial species; (2) to assess the spatial variability of cyanobacterial community composition in surface and bottom waters; and (3) to determine the variations of cyanobacteria in relation to physicochemical parameters of the waters. The results of this study will help to facilitate further investigation of the cyanobacterial community structure in Sanya Bay.

1 Materials and methods

1.1 Study area and sampling methods

Sanya Bay lies in the southernmost part (109°20'–109°30'E, 18°11'–18°18'N) of Hainan Island in China [13]. It is a typical tropical Bay with a water surface area of 120 km² and an average depth of 16 m. This bay consists of Dongmao, Ximao, and Luhuitou Islands, which are located at its mouth [13]. Except for Station 4 (26 m) and Station 9 (28 m), the rest of the sampling stations are located at depths of <20 m. Water samples were collected using 5 L Niskin bottles from the surface and the bottom of the 12 stations in Sanya Bay during surveys held on 24 and 25 April, 2010 (Figure 1). Water quality parameters such as temperature (Temp), salinity (*S*) and dissolved oxygen (DO) were measured *in situ* using a water quality Monitoring system (Hydrolab Corporation, USA). Nutrients such as nitrate (NO₃-N/μmol L⁻¹), nitrite (NO₂-N/μmol L⁻¹) and, silicate (SiO₄-Si/μmol L⁻¹)




Figure 1 Sampling stations in the coral reefs area of Sanya Bay.

were analyzed using a SKALAR autoanalyzer (Skalar, Breda, The Netherlands). In addition, ammonium (NH₄-N/μmol L⁻¹) and phosphorus (PO₄-P/μmol L⁻¹) were analyzed through the oxidation by hypobromite and molybdophosphoric blue, respectively, which were monitored used an Ultraviolet (UV) 1601 spectrophotometer (Shimadzu Corporation, Japan) [14]. Cyanobacterial abundance was analyzed according to the methods described in “*The Specialties for Marine Monitoring*” [14]. One liter of seawater samples was filtered through 0.22 μm pore size cellulose acetate membrane filters (Millipore, USA). After filtration, the membranes were immediately frozen in liquid nitrogen and then stored at -20°C in preparation for DNA extraction in the laboratory.

1.2 Physicochemical parameter analyses

Analysis of the physicochemical parameters was conducted according to methods prescribed by Huang et al. [15], Zhou et al. [16], and the Specification for Oceanography Survey [14].

1.3 DNA extraction, PCR amplification and DGGE

Community DNA extraction was performed according to the method prescribed by Bostrom et al. [17]. The 16S rDNA fragments were first amplified using the primers CYA359F and 23S30R [18]. PCR amplifications were performed using a PTC-2000 thermal cycler (Bio-Rad, USA). This thermal cycling was conducted according to the method used by Nübel et al. [19]. The second PCR reaction was performed using the forward primer CYA359F and an equimolar mixture of the reverse primers CYA781R (a) and (b) [19]. The reverse primer (a) amplifies the filamentous cyanobacteria, whereas the reverse primer (b) targets the unicellular cyanobacteria. A 40 bp GC clamp (CGCCCGCCGCGCGGGCGGGGCGGGGCGGGGCGGGGCGGGG) was added to primer 359F (5') to enhance the separation of the DNA bands in the DGGE gel [20]. Each PCR reaction was performed in triplicate to reduce possible intersample

PCR variations. The PCR products were then pooled and purified prior to loading on the DGGE gel. Equal amounts of the PCR products (40 μL PCR product with 7 μL loading dye) from the different stations were loaded onto an acrylamide gel (6% acrylamide, 1 mm thick). DGGE was performed using the INGENYphor U-2 system (Ingeny International BV, The Netherlands). The 100% denaturing solution consisted of 7 mol L^{-1} urea and 40% formamide (v/v). Optimal separation of cyanobacterial DNA in the samples was achieved using a 45%–70% denaturing gradient. DGGE was performed at 60°C for 17 h at a constant voltage of 100 V. After electrophoresis, the DGGE gels were stained with ethidium bromide and visualized under UV light using an AlphaImager imaging system (Alpha Innotech, USA). Individual DGGE bands of interest were excised and then reamplified with CYA359F- CYA781R (a) and (b) under previously described reaction conditions [19]. The PCR products were again loaded onto a DGGE gel to confirm the band positions. Products of the same mobility were purified, ligated into a pMD18-T cloning vector, and subsequently transformed into *Escherichia coli* DH5 α according to the manufacturer's instructions (Takara Shuzo Co, Ltd, Otsu, Japan). Positive recombinants were then submitted for sequencing with an M13 primer on an ABI3730 DNA Sequencer (USA) at the Shanghai Invitrogen Biotech Co., Ltd. The obtained sequences were analyzed against sequences in the Ribosomal Database Project (RDP) using the Classifier tool, and against GenBank sequences using the Basic Local Alignment Search Tool (BLAST) program [21,22]. Phylogenetic trees of 16S rDNA partial sequences were generated using the neighbor-joining algorithms in Mega IV software [23]. The evolutionary distances were computed using the maximum composite likelihood method [24] and expressed as the number of base substitutions per site. The level of support for the phylogenies derived from the neighbor-joining analysis was determined using 1000 bootstrap replicates [25].

1.4 Data analysis

All samples were collected in triplicate, and data were presented as the mean values. The DGGE fingerprint results for the surface waters and bottom waters communities were analyzed according to the method prescribed by Zhang et al. [26]. The band positions and intensities were both considered. Canoco for Windows 4.5 (Wageningen, The Netherlands) was used to determine the relationships between community compositions and physicochemical parameters. First, detrended correspondence analysis was performed to decide between a linear and a unimodal response model for the multivariate analysis. Given that the gradient lengths of the first axis were 2.269 and 4.388 for the surface and bottom layers, respectively, RDA ordination was used to investigate the spatial variations and the environmental factors [27–29].

2 Results

2.1 Physicochemical parameters of the ocean water

The basic physicochemical parameters investigated are summarized in Table 1. Salinity in the bottom layer station (34.30‰–34.41‰) was higher than that in the surface layer (34.16‰–34.37‰) at the same station. The lowest DO concentration in the surface waters (6.96 mg L^{-1}) and bottom waters (6.90 mg L^{-1}) appeared at Station W1, where Sanya River flows into the bay. Compared with other stations, this station also had the highest chlorophyll *a* and nutrient concentrations (phosphate, silicate, nitrite, nitrate and ammonium). The Chl*a* concentration in the surface waters was 2.93 $\mu\text{g L}^{-1}$, which was almost 15.4 times that in Station W10. The mean Chl*a* concentration gradually decreased from the coastal stations to off-shore stations. Flow cytometry showed that total cyanobacterial abundance in the surface and bottom waters ranged from 0.7×10^4 to 2.38×10^4 cells mL^{-1} and from 1×10^4 to 1.8×10^4 cells mL^{-1} , respectively.

2.2 Cyanobacterial DGGE profiles

The DGGE-bands were labelled “C”, followed by the specific number assigned to a specific 16S rDNA sequence obtained from the cyanobacterial communities (Figures 2–4). The sequence analysis results of the excised DGGE bands are summarized in Table 2. The nucleotide sequence accession numbers obtained in this study are available in the GenBank database under the accession numbers JF914949 to JF914964.


Most of the phylotypes (56.3%) were identified as *Synechococcus* or *Synechococcus*-like species. The rest of the phylogenetic groups identified included *Chroococciopsis* (6.3%), *Cyanobium* (6.3%) and some unclassified *cyanobacteria* (31.2%). The percentage similarity with the clone and its closest blast hits ranged from 93% to 99%, respectively. Sequences obtained from Sanya Bay were similar to those from other marine environments such as the oligotrophic Kuroshio Current near the outer edge of the East China Sea, the water column in the Red Sea, seawater of the East China Sea, and the South Atlantic Subtropical Gyre and so on.

2.3 Community composition in relation to water physicochemical properties

Table 3 and Figure 3 show that the first and second RDA axes account for 98.6% to 99.2% of the variations in the cyanobacterial community composition in surface waters. Meanwhile, for the bottom waters, 49.1% and 7.5% of the variations in the composition could be explained by the first two canonical axes, respectively. The significant environmental variables could explain 99.3% and 58.3% of the structural variations in the surface layer and the bottom

Table 1 Physicochemical parameters of the surface and bottom samples of Sanya Bay (values are given as mean, $n=3$)

Stations	Temp. (°C)	S (‰)	pH	DO (mg L ⁻¹)	Chl a (μg L ⁻¹)	COD (mg L ⁻¹)	BOD (mg L ⁻¹)	DIC (mg L ⁻¹)	DOC (mg L ⁻¹)	SiO ₂ -Si (mg L ⁻¹)	PO ₄ -P (mg L ⁻¹)	NO ₂ -N (mg L ⁻¹)	NO ₃ -N (mg L ⁻¹)	NH ₄ -N (mg L ⁻¹)	CA (cells mL ⁻¹)
WS1	26.32	34.16	8.12	6.96	2.93	0.98	0.52	27.77	1.291	0.105	0.009	0.004	0.024	0.017	2.37
WS2	26.39	34.34	8.13	7.25	1.47	1.13	0.50	27.68	2.196	0.098	0.006	0.003	0.010	0.014	2.06
WS3	26.45	34.32	8.06	7.06	0.41	0.33	0.22	28.20	1.187	0.099	0.003	0.002	0.006	0.014	0.88
WS4	26.47	34.37	8.11	7.11	0.42	0.65	0.20	27.85	1.076	0.086	0.003	0.002	0.009	0.016	0.88
WS5	26.84	34.24	8.15	7.32	0.30	0.73	0.07	27.57	1.005	0.091	0.004	0.002	0.005	0.016	0.70
WS6	26.30	34.36	8.08	7.15	0.56	0.36	0.28	28.10	1.060	0.114	0.004	0.001	0.006	0.014	1.66
WS7	26.37	34.35	8.06	7.11	0.67	0.91	0.08	28.21	0.822	0.098	0.003	0.002	0.007	0.013	1.62
WS8	26.59	34.32	8.12	7.11	0.84	0.22	0.10	27.77	1.027	0.090	0.003	0.002	0.007	0.013	1.49
WS9	26.40	34.33	8.06	7.13	1.28	0.62	0.01	28.20	0.701	0.099	0.005	0.001	0.006	0.016	2.23
WS10	26.78	34.28	8.13	7.16	0.19	0.29	0.16	27.74	1.159	0.094	0.004	0.003	0.009	0.014	0.80
WS11	26.69	34.23	8.12	7.10	0.50	0.29	0.10	27.71	0.773	0.088	0.003	0.001	0.006	0.015	1.32
WS12	27.38	34.19	8.11	7.21	0.65	0.87	0.07	27.84	0.740	0.105	0.004	0.002	0.009	0.014	1.45
WB1	26.20	34.30	8.11	6.90	1.17	0.84	0.08	27.84	0.602	0.103	0.006	0.003	0.014	0.018	1.81
WB2	26.04	34.38	8.08	7.21	0.60	0.76	0.35	28.10	1.562	0.103	0.004	0.002	0.007	0.016	1.50
WB3	26.11	34.46	8.09	7.03	1.05	0.44	0.21	28.01	1.071	0.103	0.002	0.001	0.006	0.015	0.95
WB4	26.10	34.32	8.13	7.10	0.42	0.76	0.08	27.68	1.032	0.094	0.002	0.002	0.008	0.014	1.04
WB5	26.50	34.35	8.10	7.36	1.25	0.87	0.03	27.93	0.999	0.086	0.002	0.002	0.007	0.015	1.00
WB6	26.21	34.41	8.13	7.11	0.70	0.33	0.06	27.68	0.817	0.097	0.002	0.001	0.006	0.017	1.30
WB7	26.07	34.34	8.07	7.26	0.77	0.80	0.19	28.18	1.098	0.086	0.002	0.001	0.007	0.014	1.09
WB8	26.42	34.35	8.13	7.11	0.22	0.33	0.16	27.68	0.971	0.101	0.002	0.002	0.008	0.015	1.61
WB9	26.26	34.36	8.08	7.20	1.09	0.65	1.06	28.10	1.032	0.101	0.003	0.001	0.007	0.018	1.23
WB10	26.46	34.33	8.13	7.13	0.55	7.64	0.32	27.68	1.021	0.084	0.003	0.001	0.006	0.013	1.32
WB11	26.40	34.34	8.11	7.31	0.59	0.55	0.02	27.84	0.701	0.097	0.002	0.001	0.005	0.013	1.40
WB12	26.55	34.33	8.12	7.43	0.85	0.47	0.16	27.77	0.833	0.103	0.003	0.001	0.006	0.016	1.31

**Figure 2** Denaturing gradient gel electrophoresis (DGGE) patterns of the amplified cyanobacterial 16S rDNA gene fragments of the bacterial communities at the sampling sites of Sanya Bay, and of the sequences of the labeled DGGE bands that were excised and sequenced.




Figure 3 Redundancy analysis (RDA) ordination biplots of DGGE band data for (a) the surface cyanobacterial communities and (b) the bottom cyanobacterial communities (the samples are indicated by the sample site location code, whereas the environmental variables are represented by arrows). The length of each arrow is correlated with the degree of relationship between the response variables.

Table 2 Summary of closest relatives to the excised and sequenced bands derived from the DGGE profile

Band No.	Accession No.	Database match with accession No. in parentheses	Origin	Identity (%)
C1	JF914949	<i>Synechococcus</i> sp. KORDI-49 (FJ497747)	Seawater of the East China Sea	97
C2	JF914950	Uncultured <i>cyanobacterium</i> Bac18 (HM485317)	Surface seawater (5 m) from North Western Mediterranean Sea	99
C3	JF914951	<i>Synechococcus</i> sp. MBIC10089 (AB058226)	Japan sea halotolerant	98
C4	JF914952	Uncultured <i>cyanobacterium</i> A8W_30 (HM057799)	Ocean water from the Yellow Sea	99
C5	JF914953	<i>Synechococcus</i> sp. KORDI-56 (FJ497734)	Seawater of the East Sea	99
C6	JF914954	Uncultured <i>Synechococcus</i> sp. 031806#44 (FJ903279)	Oligotroph Kuroshio Current near the outer edge of East China Sea	99
C7	JF914955	<i>Synechococcus</i> sp. KORDI-15 (FJ497743)	Seawater of the East Sea	99
C8	JF914956	Uncultured <i>Synechococcus</i> sp. JL-SCS-L51 (AY664001)	South China Sea	99
C9	JF914957	<i>Chroococidiopsis</i> sp. BB96.1 (AJ344555)	South Africa	93
C10	JF914958	<i>Synechococcus</i> sp. RS9920 (AY172830)	Water column in the Red Sea	99
C11	JF914959	Uncultured <i>cyanobacterium</i> Souza16sRcya18q (DQ363186)	Yucatan channel	99
C12	JF914960	Uncultured <i>cyanobacterium</i> ARTE9_416 (GU230462)	Coastal water	99
C13	JF914961	<i>Synechococcus</i> sp. KORDI-71 (FJ497742)	Seawater of the East China Sea	99
C14	JF914962	Uncultured <i>Synechococcus</i> sp. clone 031606#40 (FJ903235)	Oligotrophic region of the Kuroshio Current	99
C15	JF914963	<i>Cyanobium</i> sp. JJ22K (AM710364)	Mesotrophic and eutrophic freshwater reservoirs	99
C16	JF914964	Uncultured <i>cyanobacterium</i> Bac18 (HM485317)	Surface seawater from North Western Mediterranean Sea	99




Figure 4 Unrooted phylogenetic tree based on partial 16S rDNA sequences that represent the DGGE bands indicated in Figure 2. Bootstrap analysis was based on 1000 replicates. Bootstrap values from distance analyses are shown. Bootstrap values below 50% are not shown. The scale indicates 5% sequence divergence.

Table 3 Redundancy analysis results of surface and bottom bacterial DGGE profiles^{a)}

Axis	Eigen value	Species-environment correlation	Cumulative % variations of species	Cumulative % variations of species-environment	Sum of all canonical eigen value
Surface cyanobacterial communities					0.993
Axis 1	0.986	1	98.6	99.3	
Axis 2	0.006	0.72	99.2	99.9	
Axis 3	0.001	0.624	99.2	100.0	
Axis 4	0.000	0.55	99.3	100.0	
Bottom cyanobacterial communities					0.583
Axis 1	0.491	0.96	49.1	84.3	
Axis 2	0.075	0.618	56.6	97.1	
Axis 3	0.017	0.867	58.3	100	
Axis 4	0.023	0.000	81.3	0	

a) Monte Carlo significance tests for surface bacterial data: sum of all Eigen values, 1.000; significance of first canonical axis, F value= 81.709, $P=0.002$; significance of all canonical axes, F value=234.089, $P=0.002$. Monte Carlo significance tests for bacterial data: sum of all Eigen values, 1.000; significance of first canonical axis, F value=8.483, $P=0.002$; significance of all canonical axes: F value=4.233, $P=0.002$. F and P values were estimated using Monte Carlo permutations.

layer, respectively. These results indicated that 99.3% of all variations within cyanobacterial communities in the surface layer were caused by environmental factors, whereas these factors account for only 58.3% of the variations in the bottom layer communities. The first two axes of the RDA ordination indicated the species-environment relationships as 99.3% and 99.9% for the surface layer, and 84.3% and 97.1% for and bottom layer. According to the Monte Carlo test, the *P*-value (which indicate the significance) of the first canonical axis and all canonical axes for the surface communities were both 0.002, respectively, whereas those for the bottom communities were 0.016 and 0.032, respectively. These results revealed that the environmental variables played vital roles in driving the observed spatial variations in cyanobacterial community composition. Furthermore, the Monte Carlo analysis indicated that *Chla* ($P=0.002$; $P<0.01$), BOD ($P=0.014$; $P<0.05$), $PO_4\text{-P}$ ($P=0.002$; $P<0.01$), and $NO_3\text{-N}$ ($P=0.002$; $P<0.01$) showed a strong correlation with surface cyanobacterial community composition, whereas $NO_3\text{-N}$ ($P=0.001$; $P<0.01$), $PO_4\text{-P}$ ($P=0.002$; $P<0.01$), and $NO_2\text{-N}$ ($P=0.026$; $P<0.05$) significantly affected the spatial variations in the cyanobacterial species composition at the bottom layer. The environmental factors affecting these two layers were different. The cyanobacterial community in Station W1 was remarkably different from those in other stations.

3 Discussion

The different cyanobacterial species were not evenly distributed in all stations in the surface and bottom waters (Figure 2). Some bands were detected only in few stations. For instance, band C3 was only detected in the surface layer of Station W1. However, bands C4, C6, C7, C8, and C9 were only detected at the bottom layer of Station W7. Some bands were detected in most of the lanes, such as band C1, C5, C10, C11 and C12. In particular, band C10 was found in every lane but in varying densities. This distribution may be attributed to differences in nutrient concentrations. The distribution pattern of the surface layer was such that stations located near the Sanya River (e.g. W1 and W2) exhibit higher diversity than the outer stations (e.g. W4, W9, and W11), whereas the bottom layer showed no significant difference.

The predominant band in the DGGE gel was band C10, as indicated by the obtained sequences. The Blast query result showed that the highest similarity sequence for this band is that of *Synechococcus* sp. RS9920, which has been isolated from the water column in the Red Sea and can use nitrate as a sole N source for growth [30]. The closest relatives of the sequences C1, C5, C7 and C13 in NCBI Genbank were all from the East China Sea, whereas certain sequences such as C6 and C14 were related to sequences found in the oligotrophic region of the Kuroshio Current.

When the Kuroshio Current passes the Luzon Strait, it forms two branches: one flows into the South China Sea, while the other heads north to the East Sea along the eastern part of Taiwan. The similarities between the sequences obtained from the DGGE gel and those of the East Sea ranged from 97% to 99%. All related sequences were presumed to be from the Kuroshio Current [31,32]. Station W7 lies between Dongmao Island and Ximao Island, unlike in other stations, tourism and other human activities contributed a lot to the variations in the cyanobacterial community in this station [15]. Sequence C9 only showed 93% similarity to its closest relatives in the GenBank (Accession No. AJ344555), indicating that this sequence could be representative of a novel cyanobacterial species [33]. Sequence C15 was related to *Cyanobium* sp. JJ22K, which was isolated from mesotrophic and eutrophic freshwater reservoirs. This species could be found in both freshwater and marine ecosystems, indicating its high adaptability to different environments [34].

Most of the identified sequences were related to *Synechococcus* or *Synechococcus*-like species (Table 2). The cyanobacterial communities in both surface and bottom layers of Sanya Bay were apparently dominated by these, which were considered dominant phytoplankton in tropical and subtropical ocean ecosystems. According to Murphy et al. [35], *Synechococcus* are widely distributed in the different world ocean, with high abundance (ranging from 10^3 to 10^5 cells mL^{-1}), particularly in the coastal area [35].

Due to their capability of photosynthesis and nitrogen fixation, they were indispensable components of the biogeochemical cycle such as carbon cycle and nitrogen cycle. *Synechococcus* and *Prochlorococcus* are estimated to absorb approximately 1×10^{10} Tg carbon from the atmosphere each year, which is equivalent to two-thirds of the total carbon fixation that occurs in the oceans [36]. *Chroococcidiopsis*, a unicellular, non-heterocyst-differentiation genus, is a wide-spread cosmopolitan unicellular cyanobacteria found in many extreme environment, such as the airspaces of porous rocks, Antarctic valleys and hot deserts [37]. Another cyanobacterial genus detected in Sanya Bay was *Cyanobium*, which usually constitutes majority component of marine, brackish and freshwater picophytoplankton community

The vast difference in the cyanobacterial community composition of Station W1 could be attributed to the physicochemical property of the water column as a result of the discharge from the Sanya River (Figure 1). RDA results showed that the relationships of the surface and bottom cyanobacterial community composition with environmental factors were different. The significant factors for the spatial variations in surface waters composition were *Chla*, BOD, nitrate and phosphate, whereas those for the bottom waters were phosphate, nitrate and nitrite. The nitrate and phosphate concentration were significant environmental factors for both surface and bottom layers. This result may be ex-

plained by the rapid growth of phytoplankton in spring. At this stage, they consumed a considerable amount of phosphate in the ocean waters, thus causing phosphate becomes a limiting factor [38].

RDA ordination results implied that Chla, BOD, nitrate, and phosphate were positively correlated with cyanobacterial species in the surface layer, whereas nitrate, nitrite, and phosphate were positively correlated with the species in the bottom layer. These nutrients can promote cyanobacterial growth. In addition, previous studies in Sanya Bay showed that dissolved inorganic nutrients, particularly phosphate and inorganic nitrogen, were important environmental factors that determine the phytoplankton distribution [38,39]. The limiting factors differ in different ecosystems or within the same ecosystem for different species. Yan et al. [40] found that nitrate, DO, and silicate were the significant factors in the Three Gorges Reservoir. On the other hand, Zeng et al. [41] reported that total nitrogen and ammonium were the determinants of bacterial distribution in eutrophic lakes. Their nutrient addition experiment showed that different species responded differently to the same nutrient concentration. The results showed that the *Verrucomicrobia* density was the highest in the control group which contained low nutrient concentration and lower in the higher concentration groups, whereas the lowest abundance of *Synechococcus*-like species was observed in the control group. Hence, further investigation should be conducted to examine the relationship between the seasonal variations in cyanobacterial composition and the physiochemical environmental parameters [40–42].

4 Conclusions

In this study, flow cytometry, fingerprinting technique, and multivariate analysis were used to investigate the cyanobacterial community composition and its relationship with the environmental factors in Sanya Bay. The following conclusions have been reached:

(1) Cyanobacterial abundance in surface waters was higher than in bottom waters, ranging from 0.7×10^4 to 2.38×10^4 cells mL^{-1} , and 1×10^4 to 1.8×10^4 cells mL^{-1} , respectively.

(2) The predominant cyanobacterial species obtained from the DGGE gel were identified as *Synechococcus* or *Synechococcus*-like species (56.2%). Other phylogenetic groups identified include *Chroococciopsis* (6.3%), *Cyanobium* (6.3%), and some unclassified cyanobacteria (31.2%).

(3) RDA ordination results showed that the significant environmental factors for the surface layer were Chla ($P=0.002$; $P<0.01$), BOD ($P=0.014$; $P<0.05$), nitrate ($P=0.002$; $P<0.01$), and phosphate ($P=0.002$; $P<0.01$), whereas those for the bottom layer were nitrate ($P=0.001$; $P<0.01$), nitrite ($P=0.026$; $P<0.05$), and phosphate ($P=0.002$; $P<0.01$).

These environmental parameters accounted for 99.3% and 58.3% of the variations in the surface and bottom layers, respectively.

The authors thank all the staff of Hainan Tropical Marine Biology Research Station of the Chinese Academy of Sciences for providing support with the Sample collecting and data processing. This work was supported by the National Science and Technology Supporting Program (2009BAB44B03), the National High-tech R&D Program of China (2012AA092104), the National Natural Science Foundation of China (40776069, 40676091 and 41006069), and the National Basic Research Program of China (2010CB833800), the Knowledge Innovation Program of Chinese Academy of Sciences (KSCX2-EW-G-12), the Sanya Station Database and the Information System of CERN.

- Hoffman L. Marine cyanobacteria in tropical regions: Diversity and ecology. *Eur J Phycol*, 1999, 34: 371–379
- Capone D G. The marine nitrogen cycle. In: Kirchner D L, ed. *Microbial Ecology of the Oceans*. New York: Wiley, 2000. 455–494
- Zehr J P. Nitrogen fixation by marine cyanobacteria. *Trends Microbiol*, 2011, 19: 162–173
- Capone D G. Marine nitrogen fixation: What's the fuss? *Curr Opin Microbiol*, 2001, 4: 341–348
- Yang B, Dong J D, Yang Z H, et al. Chemical constituents of *Cyanobacteria Lyngbya* sp. from South China Sea (In Chinese). *J Trop Oceanogr*, 2007, 26: 67–69
- El-Bestawy E A, Abd El-Salam A Z, Mansy H A R. Potential use of environmental cyanobacterial species in bioremediation of lindane-contaminated effluents. *Int Biodeterior Biodegrad*, 2007, 59: 180–192
- Ke Z X, Huang L M, Tan Y H, et al. Plankton community structure and diversity in coral reefs area of Sanya Bay, Hainan Province, China. *Biod Sci*, 2011, 19: 696–701
- Shi Q, Zhao M X, Zhang Q M, et al. Estimate of carbonate production by scleractinian corals at Luhuitou fringing reef, Sanya, China. *Chin Sci Bull*, 2009, 54: 696–705
- Zhang Q M, Shi Q, Chen G, et al. Status monitoring and management strategy research of Luhuitou fringing reef of the Sanya reserve. *Chin Sci Bull*, 2006, 51(Suppl II): 71–77
- Dong J D, Zhang Y Y, Wang Y S, et al. Spatial and seasonal variations of Cyanobacteria and their nitrogen fixation rates in Sanya Bay, South China Sea. *Sci Mar*, 2008, 72: 239–251
- Dong J D, Wang H K, Zhang S, et al. Vertical distribution characteristics of seawater temperature and DIN in Sanya Bay (in Chinese). *J Tropic Oceanol*, 2002, 21: 40–47
- Muyzer G, de Waal E C, Uitterlinden A G. Profiling of complex microbial populations by denaturing gradient gel electrophoresis analysis of polymerase chain reaction amplified genes encoding for 16S rRNA. *Appl Environ Microbiol*, 1993, 59: 695–700
- Kataoka T, Hodoki Y, Koji S, et al. Tempo-spatial patterns of bacterial community composition in the western North Pacific Ocean. *J marine Syst*, 2009, 77: 197–207
- General Administration of Quality Supervision, Inspection and Quarantine of the People's Republic of China and China National Standardization Management Committee. *The Specialties for Marine monitoring (GB17378.4-1998, China)* (in Chinese). Beijing: Standards Press of China, 1998. 142–162
- Huang L M, Tan Y H, Song X Y, et al. The status of the ecological environment and a proposed protection strategy in Sanya Bay, Hainan Island, China. *Mar Pollut Bull*, 2003, 47: 180–186
- Zhou W, Li T, Cai C H, et al. Spatial and temporal dynamics of phytoplankton and bacterioplankton biomass in Sanya Bay, northern South China Sea. *J Environ Sci-China*, 2009, 21: 595–603
- Bostrom K H, Simu K, Hagstrom A, et al. Optimization of DNA extraction for quantitative marine bacterioplankton community analysis. *Limnol Oceanogr Meth*, 2004, 2: 365–373

- 18 Lepère C, Wilmotte A, Meyer B. Molecular diversity of *Microcystis* strains (Cyanophyceae, Chroococcales) based on 16S rDNA sequences. *Syst Geogr Pl*, 2000, 70: 275–283
- 19 Nübel U, Garcia-Pichel F, Muyzer G. PCR primers to amplify 16S rRNA genes from cyanobacteria. *Appl Environ Microb*, 1997, 63: 3327–3332
- 20 Muyzer G, Smalla K. Application of denaturing gradient gel electrophoresis (DGGE) and temperature gradient gel electrophoresis (TGGE) in microbial ecology. *Anton Leeuw Int J G*, 1998, 73: 127–141
- 21 Maidak B L, Cole J R, Parker C T, et al. A new version of the RDP (Ribosomal Database Project). *Nucleic Acids Res*, 1999, 27: 171–173
- 22 Altschul S F, Madden T L, Schaeffer A A, et al. Gapped BLAST and PSI-BLAST: A new generation of protein database search programs. *Nucleic Acids Res*, 1997, 25: 3389–3402
- 23 Tamura K, Dudley J, Nei M, et al. MEGA4: Molecular evolutionary genetics analysis (MEGA) software version 4.0. *Mol Biol Evol*, 2007, 24: 1596–1599
- 24 Tamura K, Nei M, Kumar S. Prospects for inferring very large phylogenies by using the neighbor-joining method. *Proc Natl Acad Sci USA*, 2004, 101: 11030–11035
- 25 Saitou N, Nei M. The neighbor-joining method: A new method for reconstructing phylogenetic trees. *Mol Biol Evol*, 1987, 4: 406–425
- 26 Zhang Y Y, Dong J D, Yang Z H, et al. Phylogenetic diversity of nitrogen-fixing bacteria in mangrove sediments assessed by PCR-DGGE. *Arch Microbiol*, 2008, 190: 19–28
- 27 Zhang J C, Zeng G M, Chen Y N, et al. Effects of physico-chemical parameters on the bacterial and fungal communities during agricultural waste composting. *Bioresource Technol*, 2011, 102: 2950–2956
- 28 Lepš J, Šmilauer P. *Multivariate Analysis of Ecological Data Using CANOCO*. Cambridge: Cambridge University Press, 2003. 43–75
- 29 Luan Q S, Sun J, Song S Q, et al. Canonical correspondence analysis of summer phytoplankton community and its environment in the Yangtze River estuary China (in Chinese). *J Plant Ecol*, 2007, 31: 445–450
- 30 Fuller N J, Marie D, Partensky F. Clade-specific 16S ribosomal DNA oligonucleotides reveal the predominance of a single marine *Synechococcus* clade throughout a stratified water column in the Red Sea. *Appl Environ Microbiol*, 2003, 2430–2443
- 31 Choi D H, Noh J H. Phylogenetic diversity of *Synechococcus* strains isolated from the East China Sea and the East Sea. *FEMS Microbiol Ecol*, 2009, 69: 439–448
- 32 Liang W D, Yang Y J, Tang T Y, et al. Kuroshio in the Luzon Strait. *J Geophys Res*, 2008, 113: C08048
- 33 Jezberova J. Phenotypic diversity and phylogeny of picocyanobacteria in mesotrophic and eutrophic freshwater reservoirs investigated by a cultivation-dependent polyphasic approach. The Dissertation for the Doctoral Degree. Ceske Budejovice: University of South Bohemia, 2006
- 34 Rohwer F, Breitbart M, Jara J, et al. Diversity of bacteria associated with the Caribbean coral *Montastraea franksi*. *Coral Reefs*, 2001, 20: 85–91
- 35 Murphy L S, Haugen E M. The distribution and abundance of phototrophic ultraplankton in the North Atlantic. *Limnol Oceanogr*, 1985, 30: 47–58
- 36 Iturriaga R, Mitchell B G. Chroococcoid cyanobacteria: A significant component in the food web dynamics of the open ocean. *Mar Ecol Prog Ser*, 1986, 28: 291–297
- 37 Fewer D, Friedl T, Budel B. *Chroococciopsis* and heterocyst-differentiating cyanobacteria are each other's closest living relatives. *Mol Phylogenet Evol*, 2002, 23: 82–90
- 38 Zhang Y Y, Dong J D, Ling J, et al. Phytoplankton distribution and their relationship to environmental variables in Sanya Bay, South China Sea. *Sci Mar*, 2010, 74: 783–792
- 39 Yang Z H, Dong J D, Wu M L, et al. Structure characteristics of phytoplankton community at Sanya Bay (in Chinese). *J Trop Oceanogr*, 2007, 26: 62–66
- 40 Yan Q Y, Yu Y H, Feng W S, et al. Plankton community composition in Three Gorges Reservoir region revealed by PCR-DGGE and its relationships with environmental factors. *J Environ Sci*, 2008, 20: 732–738
- 41 Zeng J, Yang L Y, Du H W, et al. Bacterioplankton community structure in a eutrophic lake in relation to water chemistry. *World J Microbiol Biotechnol*, 2009, 25: 763–772
- 42 Sipura J, Haukka K, Helminen H, et al. Effect of nutrient enrichment on bacterioplankton biomass and community composition in mesocosms in the Archipelago Sea, northern Baltic. *J Plankton Res*, 2005, 27: 1261–1272

Open Access This article is distributed under the terms of the Creative Commons Attribution License which permits any use, distribution, and reproduction in any medium, provided the original author(s) and source are credited.