

Poster presentation

Dopamine mediated dynamical changes in the striatum: a numerical study

Krisztina Szalicsnyó* and László Müller

Address: Biophysics Department, Computational Neuroscience Group, Particle and Nuclear Physics Institute of the Hungarian Academy of Sciences, Budapest, Hungary

Email: Krisztina Szalicsnyó* - szali.@sunserv.kfki.hu

* Corresponding author

from Eighteenth Annual Computational Neuroscience Meeting: CNS*2009
Berlin, Germany. 18–23 July 2009

Published: 13 July 2009

BMC Neuroscience 2009, **10**(Suppl 1):P150 doi:10.1186/1471-2202-10-S1-P150This abstract is available from: <http://www.biomedcentral.com/1471-2202/10/S1/P150>

© 2009 Szalicsnyó and Müller; licensee BioMed Central Ltd.

The striatum is a part of the basal ganglia, which are a group of nuclei in the brain associated with motor control, cognition and learning. In this study we examined the consequences of the dopamine modulation in a small striatal network. We employed point neuron models to analyze the conductance based dopaminergic changes. The model is built from the following elements: tonically active neuron (cholinergic interneuron) (TAN), dopaminergic neuron (DAN), medium spiny neuron (MSN) and fast spiking interneuron (FSN). TANs are able to fire in the absence of synaptic inputs and respond to sensory stimuli and sensorimotor learning by transiently suppressing their firing activity [1]. This pause is dopamine signal sensitive, but the neurophysiological mechanism of the dopaminergic influence is under debate. We analyzed the robustness of the TAN subthreshold oscillations and demonstrated how they are affected by dopaminergic modulation [1]. The TAN-DAN interaction is reciprocal and precisely timed [2]. TAN pause responses co-occur with the DAN bursts and both influence the activities of the MSN neurons and the feed-forward FSN neurons. Our aim was to examine the dynamic interactions in this network and study the effects of the dopaminergic/cholinergic time-dependent modulations [3].

Our results predict that the dopamine mediated effects (through D1 and D2 receptors) are able to switch the TANs between stable oscillatory and fixed-point behaviors [1]. The results suggest that the MSN neurons exhibit dynamical sub-threshold hysteresis without showing

static hysteresis and this bi-stability is dopamine dependent [4]. We further predict that different dopamine receptors (D(1) and D(2)) mediate opposing dynamical effects on these cell types (small network) and we suggest that these opposing effects act on different timescales.

Our work seeks to more deeply understand the details of the striatal small network dynamics and give predictions for the possible dynamical consequences of the dopamine depleted states, where the cortico-striatal coupling is weakened and the striatal firing thresholds are reduced [5,6].

We thank to Peter Simon for his help and useful discussions. KSz was supported by the Eötvös Fellowship. This work was further supported by the EU Sixth Framework programme grant no.: IST-4-027819-IP, ICEA).

References

1. Szalicsnyó K, Müller L: **Dopamine induced switch in the sub-threshold dynamics of the striatal cholinergic interneurons: a numerical study.** *J Theor Biol* 2009, **256**:547-60.
2. Tan CO, Bullock D: **A dopamine-acetylcholine cascade: simulating learned and lesion-induced behavior of striatal cholinergic interneurons.** *J Neurophysiol* 2008, **100**:2409-2421.
3. Gragg SJ: **Meaningful silences: how dopamine listens to the Ach pause.** *Trends in Neurosci* 2006, **2**:125-131.
4. Gruber AJ, Solla SA, Surmeier DJ, Houk JC: **Modulation of striatal single units by expected reward: a spiny neuron model displaying dopamine-induced bistability.** *J Neurophysiol* 2003, **90**:1095-1114.
5. van Albada SJ, Robinson PA: **Mean-field modeling of the basal ganglia-thalamocortical system. I. Firing rates in healthy and parkinsonian states.** *J Theor Biol* 2008 in press.

6. van Albada SJ, Gray RT, Drysdale PM, Robinson PA: **Mean-field modeling of the basal ganglia-thalamocortical system. II. Dynamics of parkinsonian oscillations.** *J Theor Biol* 2008 in press.

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp

