



Animal (2008), 2:12, pp 1802–1808 © The Animal Consortium 2008
doi:10.1017/S1751731108003157

Effect of concentrate percentage on ruminal pH and time-budget in dairy goats

M. Desnoyers[†], C. Duvaux-Ponter, K. Rigalma, S. Roussel, O. Martin and S. Giger-Reverdin

UMR INRA-AgroParisTech Physiologie de la Nutrition et Alimentation, 16 rue Claude Bernard, 75005 Paris, France

(Received 21 February 2008; Accepted 2 August 2008; First published online 16 September 2008)

The aim of this study was to compare rumen pH and time-budget in eight mid-lactation goats receiving two diets in a cross-over design (low-concentrate diet (L): 30% and high-concentrate diet (H): 60% concentrate). Feeding H increased daily intake ($4.3 \pm 0.08\%$ v. $4.7 \pm 0.08\%$ of body weight for L and H, respectively) and daily milk production (3.01 ± 0.130 v. 3.50 ± 0.130 kg/day of 3.5% fat-corrected milk for L and H, respectively). It decreased milk fat and inverted the fat-to-protein ratio (1.07 ± 0.054 v. 0.94 ± 0.054 for L and H, respectively). As suggested by the percentage of time spent with rumen pH below 6.0 ($23.4 \pm 6.60\%$ v. $39.9 \pm 5.88\%$ for L and H, respectively), H was more acidogenic than L. When offered H instead of L, goats spent less time eating (298 ± 17.5 v. 265 ± 17.5 min for L and H, respectively) and ruminating (521 ± 21.0 v. 421 ± 21.0 min for L and H, respectively) but more time resting (352 ± 27.1 v. 459 ± 21.1 min for L and H, respectively) over a 24-h period. They also tended to spend more time drinking (20 ± 2.9 v. 25 ± 2.9 min for L and H, respectively; $P = 0.08$) when offered H rather than L. These differences in activities were mainly observed during the first hours following feeding. When offered H, goats adapted their feeding behaviour around the feedings, which allowed them to limit the physiological disturbances potentially inducible by H and to increase milk production, without experiencing too much acidosis.

Keywords: acidosis, animal behaviour, dairy goat, rumen pH

Introduction

A major negative consequence of feeding high-concentrate diets to high-producing ruminants is the occurrence of subacute ruminal acidosis. Acidosis is usually defined as a decrease in rumen pH below a threshold value of 6.0, but its severity is related to the frequency and duration of alterations in rumen pH. Acute acidosis is defined by long bouts of rumen pH below 5.0 and subacute acidosis by mean rumen pH below 6.0 and short bouts of rumen pH between 5.5 and 5.0 (Nocek, 1997; Oetzel, 2000). For Sauvant *et al.* (1999), a mean rumen pH of 6.25 corresponds to around 4 h spent below 6.0 and could thus be used as a threshold to define the occurrence of subacute acidosis. Subacute ruminal acidosis is one of the major concerns of current ruminant nutrition because it is poorly detected in herds and it has many consequences, such as decreased milk production, decreased efficiency of milk production, premature culling and increased mortality (Krause and Oetzel, 2005). Subacute acidosis is frequently studied from digestive or metabolic points of view (Braun *et al.*, 1992; Martin *et al.*, 2006; Peyraud and Apper-Bossard, 2006) but

behavioural aspects are very seldom taken into account. On a daily basis, goats adapt their feeding behaviour depending on the composition of the diet offered, especially in terms of the number of meals and meal layout during the day, as shown by Abijaoudé *et al.* (2000a and 2000b). Thus, goats might adapt their daily time-budget according to the percentage of concentrate in the diet.

The aim of this study was to determine adaptations in goat behaviour due to high-concentrate diets, by comparing, in the same animals, feed intake, milk production, rumen pH and time-budget with two diets differing in their concentrate percentage.

Material and methods

Diets, animals and experimental design

Eight fistulated dairy goats (Saanen and Alpine) in mid-lactation (79 ± 5.6 days in milk), averaging 60 ± 4.9 kg body weight (BW) and producing 3.1 ± 0.60 kg milk per day at the start of the experiment, were used. The experiment was conducted under the guidelines given by the French Agriculture Ministry. Animals were assigned to two groups, which were balanced according to goat parturition date,

[†] E-mail: marion.desnoyers@agroparistech.fr

Table 1 Composition and analysis of two experimental diets containing either a low (L) or a high (H) percentage of concentrate

Diet	L	H
Composition		
Concentrate (% DM) ^a	30.0 ^b	60.0 ^c
Grass hay (% DM)	46.6	26.6
Sugar beet pulp (% DM)	23.4	13.4
Dry matter (%)	58	67
Net energy (MJ/kg DM)	6.28	6.78
PDIN (g/kg DM)	100	136
PDIE (g/kg DM)	105	123
Analysis (% DM)		
NDF	48.2	37.9
ADF	24.0	16.7
ADL	2.4	1.3

DM=dry matter.

^a20% wheat, 20% barley, 20% oats, 35% soyabean meal, 3% mineral and vitamin premix and 2% molasses.

^bComposition of the mineral and vitamin premix (per kg of premix): 702 g CaCO₃, 119 g dicalcium phosphate, 88 g NaCl, 21.21 g ZnSO₄·7H₂O, 1.04 g/kg MnSO₄·H₂O, 0.02 g CoSO₄, 0.05 g KI, 0.02 g Na₂SeO₃, 400 000 IU Vitamin A, 100 000 IU Vitamin D, 2000 IU Vitamin E, 63.33 g corn starch.

^cComposition of the mineral and vitamin premix (per kg of premix): 702 g CaCO₃, 88 g/kg NaCl, 6.82 g ZnSO₄·7H₂O, 1.04 g MnSO₄·H₂O, 0.01 g CoSO₄, 0.03 g KI, 0.01 g Na₂SeO₃, 200 000 IU Vitamin A, 50 000 IU Vitamin D, 1000 IU Vitamin E, 199.72 g corn starch.

BW, milk production the previous year and breed. They were housed in 2 m × 1 m individual pens throughout the experimental period, with free access to water.

Diet composition is described in Table 1. Energy and nitrogen values of the feed and the diet were calculated according to the additive method from the INRA tables (Baumont *et al.*, 2007). Goats were given two total mixed rations: high-concentrate diet (H) containing 60% concentrate and low-concentrate diet (L) containing 30% concentrate. Diets were offered in a cross-over design of two 4-week periods separated by 3 days of transition diet. The concentrate part of the diets was composed of wheat (20%), barley (20%), oats (20%), soyabean meal (35%) and a vitamin and mineral mixture. Forage was a mixture of two-thirds chopped grass hay and one-third ensiled sugar beet pulp on a dry matter basis. Goats were fed *ad libitum* and quantities offered were adjusted weekly to ensure 10% refusals. Feed was offered individually twice a day after milking, in the proportion of two-thirds in the afternoon and one-third in the morning, according to the time interval between milkings.

Experimental measurements

Experimental measurements were performed during all the weeks of the cross-over design. Animals were weighed weekly. Offered feed and refusals were individually weighed daily, which allowed the assessment of dry matter intake (DMI) using the theoretical dry matter percentage of the diet. Daily DMI per kg of BW was calculated using the BW of the previous week. Day-to-day variability in DMI (cvDMI) was assessed using the coefficient of variation calculated as the absolute value of the difference in DMI

between two successive days divided by the mean value of these 2 days. Raw milk yield (RMV) was recorded daily and milk composition was analysed once a week on milk from two consecutive milkings. Fat-corrected milk (35 g/kg basis) (FCM) was calculated according to the formula of Sauvant *et al.* (2007), and day-to-day variability in milk yield (cvRMV) was assessed using the coefficient of variation calculated as the absolute value of the difference in RMV between two successive days divided by the mean value of these 2 days. Energy and nitrogen balances were calculated according to the formula published by Sauvant *et al.* (2007).

Rumen pH was continuously monitored by indwelling pH probes. Goats were accustomed, for 1 week before the start of the experiment, to wear a coat covering their back and maintained by individually adjusted elastic straps. Rumen pH was measured every minute by a self-cleaning pH probe (accuracy = 0.01 pH; Fisher Bioblock Scientific, France) placed in the rumen through the ruminal cannulae and linked to a portable device (Easy Log EL-2; Omega Engineering Inc.) (Brossard *et al.*, 2003) placed in one of the coat pockets. A 300 g weight was attached to each probe to reduce its mobility in the rumen. Mean daily pH and percentage of time when the pH was below 6.0 were calculated.

Time-budget of the goats was analysed using four video cameras recording two goats at a time. They were fitted to the ceiling above the individual pens and linked to a quad splitter allowing cyclic sequences to be recorded with a time-lapse video recorder. Nycthemeral activity of the goats was recorded during the last two weekends of each period. Twenty-four hours, from the end of the afternoon milking to the start of the next afternoon milking, were analysed during each of these weekends by scan sampling of 5 s every 2 min using The Observer software (version 5.0, 2004; Noldus Information Technology, Wageningen, The Netherlands). The calculated variables were total time spent standing, lying, ruminating, eating and idling, latency for rumination after feeding and mean duration of bouts for all these behaviours. The variables were analysed throughout the day and by 2-h intervals (12 intervals). Posture and activities were recorded independently.

Statistical analyses

Data were analysed using a repeated measures analysis of variance (ANOVA) and statistical analyses were carried out by the mixed-model procedure of SAS (version 9.1, 2002; SAS Institute Inc., Cary, NC, USA), using the following model:

$$Y_{ijkt} = \alpha_i + \beta_t + \gamma_j + \delta_k + (\gamma\delta)_{jk} + e_{ijkt},$$

where α_i is the random effect of the goat; β_t , γ_j and δ_k are, respectively, the fixed effect of time (day or week number according to the parameter), diet (H or L) and experimental period, $(\gamma\delta)_{jk}$ is the interaction between diet and experimental period and e_{ijkt} is the residual error.

Milk production analysis used DMI as a covariable in the model.

Table 2 Effect of feeding a low- (L: 30%) or high- (H: 60%) concentrate diet in a cross-over design on intake and production parameters

	Diet		s.e. ^a	p ^b
	L	H		
DMI (% of BW/day)	4.31	4.66	0.084	***
DMI variation coefficient ^c	0.22 (0.05)	0.25 (0.06)	0.010	**
Body weight (kg)	62.5	61.9	0.48	ns
Raw milk yield (kg/day)	3.08	3.56	0.044	***
3.5% Fat Corrected Milk (kg/day)	3.01	3.50	0.130	***
RMY variation coefficient ^c	0.25 (0.06)	0.24 (0.06)	0.015	ns
Milk fat (%)	3.43	3.08	0.146	*
Milk protein (%)	3.20	3.29	0.033	ns
Milk fat/protein ratio	1.07	0.94	0.054	*
Energy intake (MJ/day) ^d	16.9	19.4	0.35	***
Energy balance (MJ/day) ^d	2.1	3.9	0.50	***
Nitrogen intake (g PDI/d) ^d	269	354	6.1	***
Nitrogen balance (g PDI/d) ^d	75	148	8.4	***

DMI = dry matter intake; RMY = ram milk yield.

^aStandard error of the difference of least square mean.

^b†: 0.05 ≤ P < 0.10, ns: P ≥ 0.10.

^cSquare root transformed data with backtransformed data in brackets.

^dCalculated from the INRA Tables (Baumont *et al.*, 2007).

Nycthemeral kinetics of rumen pH were analysed using a repeated measures ANOVA using the following model:

$$Y_{ijk} = \alpha_i + \beta_j + (\alpha\beta)_{ij} + e_{ijk},$$

where α_i is the random effect of the time after feeding (4-min intervals), β_t is the fixed effect of the diet (H or L) and $(\alpha\beta)_{ij}$ is the interaction between diet and time after feeding.

When assumptions of homogeneity of variance and normal distribution of the residuals were not confirmed (cvDMI and cvRMY), a square root transformation was performed before carrying out the analysis. All data are presented as least square means (lsmean) ± standard errors (s.e.) except when otherwise stated.

Results

One of the four goats that started with H suffered from an acute bout of acidosis during the first experimental period. Its milk production decreased abruptly and did not increase thereafter, even while receiving the L diet. This goat was removed from all the calculations.

Body weight, intake and milk production

BW, intake and milk production results are presented in Table 2. BW was not influenced by the diet. Increasing the percentage of concentrate in the diet increased DMI, RMY, FCM and cvDMI. cvRMY was not influenced by the concentrate percentage. Milk protein percentage was not influenced by the diet but fat percentage decreased when the percentage of concentrate increased in the diet. Fat-to-protein ratio was lower for H than for L and inverted between the two diets. Energy and nitrogen balances were positive for the two diets but were almost doubled for H compared to L.

Rumen pH and acidosis

Mean daily pH was lower for H than for L (6.09 ± 0.071 v. 6.25 ± 0.072 , respectively; $P = 0.003$). The percentage of time when the pH was below 6.0 increased with the concentrate percentage ($23.4 \pm 6.60\%$ v. $39.9 \pm 5.88\%$ for L and H, respectively; $P = 0.002$).

Figure 1 represents the nycthemeral kinetics of rumen pH. The interaction between diet and time after feeding, when analysed by 4-min intervals, was not significant. This indicates that the pattern of averaged diurnal rumen pH was similar between treatments and the curve for H was below that of the curve for L by 0.23 pH units ($P < 0.001$).

Nycthemeral activity

Daily total durations and behavioural bout mean durations are presented in Table 3. Increasing the percentage of concentrate in the diet tended to decrease the total duration of intake and decreased the mean duration of intake bouts. The total time spent drinking tended to increase with H compared to L. The total time spent idling was longer



Figure 1 Nycthemeral kinetics of rumen pH in seven mid-lactation dairy goats offered a low- (L: 30%) or high- (H: 60%) concentrate diet (arrows represent feeding).

when goats received H than when they received L but the mean duration of idling bouts was not influenced by the diet. Increasing the percentage of concentrate in the diet decreased total time spent ruminating and mean duration of rumination bouts. The total time spent lying or standing and the mean duration of lying and standing bouts were

not influenced by the diet. Latency before the start of the first rumination bout was shorter when the animals were offered H than when they were offered L after the afternoon, but not after the morning feeding (Table 3).

The time spent eating and ruminating during each of the 12 intervals is presented in Figure 2. Goats spent less time eating with H than with L only during the two intervals including feeding (0730 to 0930 h and 1530 to 1730 h) and during the first interval after the afternoon feeding (1730 to 1930 h). No differences were observed during the remaining intervals. When offered H, goats spent less time ruminating after the morning feeding (0930 to 1330 h), during the evening (1930 to 2330 h) and during the early morning (0530 to 0730 h), but spent more time ruminating during the interval including the afternoon feeding (1530 to 1730 h). No differences were observed during the remaining intervals.

The time spent drinking and idling during each of the 12 intervals is not presented as a figure. The time spent drinking was higher with H than with L during two intervals (1930 to 2130 h; 166 ± 43.7 v. 43 ± 44.3 s for H and L, respectively; $P = 0.04$ and 0130 to 0330 h; 28 ± 7.7 v. 0 ± 7.7 s for H and L, respectively; $P = 0.019$). No differences were found during all the other intervals.

The time spent idling tended to be increased by H compared to L during the first three intervals (0730 to 1330 h; 7 ± 3.0 v. 2 ± 3.1 min; $P = 0.097$; 37 ± 6.8 v. 22 ± 7.0 min; $P = 0.076$ and 1530 to 0330 h; 51 ± 7.6 v. 39 ± 8.0 min, $P = 0.095$, for H and L, respectively), and was increased during the afternoon feeding and the following interval (1530 to 1930 h; 15 ± 3.1 v. 2 ± 3.1 min, $P = 0.01$ and 42 ± 7.5 v. 20 ± 7.9 min, $P = 0.01$, for H and L, respectively). No differences were found during the remaining intervals.

Table 3 Behavioural parameters over 24 h in dairy goats offered a low- (L: 30%) or high- (H: 60%) concentrate diet

	Diet		s.e. ^a	P ^b
	L	H		
Total duration of postures (min/day)				
Standing	536	532	33.7	ns
Lying	828	844	36.3	ns
Total duration of activities (min/day)				
Intake	298	265	17.5	†
Rumination	521	421	21.0	***
Idling	352	459	27.1	**
Drinking	20	25	2.9	†
Mean duration of postures (min/bout)				
Standing	20	24	2.4	ns
Lying	26	18	5.8	ns
Mean duration of activities (min/bout)				
Intake	13	10	1.3	*
Rumination	18	13	1.3	*
Idling	13	12	0.7	ns
Rumination latency (min) ^c				
Afternoon feeding	110	71	13.0	*
Morning feeding	80	77	9.3	ns

^aStandard error of the difference of least square mean.

^b†: $0.05 \leq P < 0.10$; ns: $P \geq 0.10$.

^cLatency of the first rumination bout after feeding.




Figure 2 Nycthemeral kinetics of eating and ruminating in mid-lactation dairy goats offered a low- (L: 30%) or high- (H: 60%) concentrate diet. Interval starting times are indicated below. Arrows represent feeding. †: $0.05 \leq P < 0.10$; NS: $P \geq 0.10$.




Figure 3 Nycthemeral kinetics of standing and lying in mid-lactation dairy goats offered a low- (L: 30%) or high- (H: 60%) concentrate diet. Interval starting times are indicated below. Arrows represent feeding. †: $0.05 \leq P < 0.10$; NS: $P \geq 0.10$.

Figure 3 shows for each interval the time spent standing and lying. During the interval including the afternoon feeding (1530 to 1730 h) increasing the percentage of concentrate in the diet tended to decrease the time spent standing. During the next interval (1730 to 1930 h) goats spent less time standing and more time lying when offered H than when offered L, while the opposite was observed for the next two intervals (1930 to 2330 h). No differences were found during the other intervals.

Discussion

Feeding a high-concentrate diet increased DMI, which agrees with the low rumen fill effect of concentrate compared to roughage (Jarrige *et al.*, 1995). Feeding a high-concentrate diet also increased milk production, which agrees with other experiments performed in goats (Kawas *et al.*, 1991) and cows (Manson and Leaver, 1988). This is also in accordance with the higher nitrogen and energy content of the H diet and the two-fold increases in nitrogen and energy balance in H compared to L. Changes in milk composition were in accordance with data from Kawas *et al.* (1991) in goats, Manson and Leaver (1988) in cows and Susin *et al.* (1995) in ewes although another experiment did not show any changes in milk composition or DMI when high-concentrate diets based on barley silage were offered to dairy cows (Maekawa *et al.*, 2002).

Goats did not exhibit any clear symptoms of acidosis. Although the higher variability in day to day DMI observed with H could be a sign of subacute acidosis (Nocek, 1997; Owens *et al.*, 1998; Martin *et al.*, 2006), DMI was increased for H compared to L, which is not in accordance with the

literature, showing a decrease in DMI with acute acidosis (Owens *et al.*, 1998; Oetzel, 2000). The difference in the percentage of time spent with a pH below 6.0 between the two diets was similar to that observed by Keunen *et al.* (2002) after the induction of subacute acidosis in dairy cows, which confirms that the animals fed H were suffering from subacute acidosis. However, the daily mean pH obtained with both diets in the present experiment are very close to the threshold of 6.25 proposed by Sauvant *et al.* (1999) to characterize subacute acidosis and they both induced at least 4 h of rumen pH below 6.0. Thus, according to the definition of Sauvant *et al.* (1999) both diets induced some subacute acidosis but it is reasonable to suggest that H was more acidogenic than L. However, because the two diets not only differed in their acidogenic capacity but also in their fibre, energy and protein content, it was difficult to determine the consequences of acidosis compared to those due to diet composition.

Regardless of the diet offered, the goats spent more time ruminating than eating, which is in accordance with results obtained in lactating goats (Kawas *et al.*, 1991) and cows (Deswysen *et al.*, 1993; Maekawa *et al.*, 2002), but not with the results of Rapetti *et al.* (2005) and Abijaoudé *et al.* (2000b) in lactating goats. The total time spent eating and ruminating was shorter when the goats were offered H than L, which is in accordance with the literature on goats and cows (Kawas *et al.*, 1991; Abijaoudé *et al.*, 2000b; Maekawa *et al.*, 2002) and confirms that the total time spent chewing (intake and rumination) decreases when the proportion of NDF decreases in the diet (Maekawa *et al.*, 2002). When a physiological parameter changes too much, homeostatic regulation mechanisms try to maintain it within physiological limits (Sauvant, 1992). Thus, the higher concentrate percentage of the H diet

probably induced a faster and more intense fermentation in the rumen, which is known to induce satiety due to the increase in rumen lactic acid (Buéno, 1975), propionic acid (Allen, 2000), osmotic pressure (Rémond *et al.*, 1995) or volatile fatty acids (Forbes, 2007). The shorter total chewing time observed with H might have limited the total amount of saliva and buffers secreted (Sauvant *et al.*, 2006), which could have increased the risk of acidosis for the animals offered H. Both the increase in the fermentation processes and the decrease in buffers from chewing could explain the lower rumen pH observed with H compared to L, but the difference was however lower than expected. Goats tended to spend more time drinking when offered H than when offered L. This might be due to the increase in osmolarity of the rumen content and also to the lower water content of the diet H compared to L. This result, however, has to be confirmed because scan sampling techniques are less precise for measuring behaviours of short duration, like drinking, than those of long duration, like feeding, standing or lying (Mitlöhner *et al.*, 2001) and because the differences in drinking time observed during our experiment are of the same magnitude (2 min) as the scan sampling interval.

The mean durations of behavioural bouts (except standing) were shorter when the goats were offered H than when they were offered L, even if it was not always significant. No data were found for comparison in the literature. The shorter mean durations observed for intake and rumination bouts might be related to the shorter total intake and rumination duration observed during 24 h. Shorter intake bouts might have resulted from earlier satiety with H than with L as already discussed. With high-concentrate diets, the more relevant factor inducing satiety is the production of volatile fatty acids by ruminal fermentation, whereas in more fibrous diets, the main factor inducing satiety is the physical repletion of the rumen (see review by Forbes, 2007). The major satiety factor was probably VFA production for both L and H diets, but their production was probably faster and higher with H than with L, inducing earlier satiety signals and shorter mean durations of intake bouts. These shorter mean intake bouts allowed the daily repartition of chewing activities to increase during the day, which may have limited metabolic disorders, such as a decrease in pH. In turn, this might explain the small difference in rumen pH observed between the two diets. This increased daily repartition of intake has already been observed with a high-concentrate diet (Abijaoudé *et al.*, 2000b). These shorter activity bouts may also highlight a greater activity level of the goats offered H. Some goats were actually very nervous, searching for straw, salt or something else to eat, particularly during milking when they were outside their individual pens. Sawyer (1998) showed that excessive activity might be a sign of discomfort or pain, but more observations are needed to confirm that high-concentrate diets or subacute acidosis can cause discomfort or pain.

The majority of rumination occurred during the night and in the morning, which is in accordance with the review by Beauchemin *et al.* (1990). However, regardless of the diet offered, the goats spent at least a little time ruminating during

all the intervals, which could also have minimized fluctuations in rumen pH (Beauchemin *et al.*, 1990). The decrease in the total time spent ruminating when goats received H was due to the decreased rumination time during the morning and the evening but not during the main night rumination period. The nycthemeral pattern of rumination was thus not really influenced by the diet. Goats offered H tended to spend more time resting during the morning while goats offered L spent more time ruminating, but the differences in resting time were of a lower magnitude than those for rumination.

During the early morning (0530 to 0730 h) intake increased for both diets compared to the night intervals even without any feeding occurring, which agrees with data from Dulphy and Faverdin (1987). The majority of intake occurred during the day (0530 to 2130 h) with two intake peaks after milking and feeding (0800 and 1600 h), as has already been described (Beauchemin *et al.*, 1990; DeVries *et al.*, 2003). During the two intervals including feeding and during the first interval following the afternoon feeding, goats spent less time eating with H than with L. As it has already been discussed, this might be due to earlier satiety and thus a more rapid end to the meal due to higher fermentation rates with H than with L. These shorter main meals are in accordance with previous observations with high-concentrate diets (Abijaoudé *et al.*, 2000a).

The time spent idling can be interpreted as resting periods, even if it was recorded during both standing and lying, because secondary behaviours (self-grooming, interaction with the environment, observation of the environment, etc.) were recorded separately and therefore were not included in this idling time. Idling was longer when goats received H than when they received L during the intervals following feeding, which is in accordance with previous observations on intake and rumination. It also shows that there is a period without any activity between the end of a meal and the start of rumination, which agrees with results from Dulphy and Faverdin (1987). This idling bout seemed to be longer with H than with L, but as for meal determination we only assessed idling bouts and could not cluster them into a longer resting bout separating the end of the meal from the start of the first rumination bout.

Therefore, nycthemeral pattern was mainly influenced by the diet during the hours following feedings. When offered H, goats adapted their feeding behaviour, according to metabolic signals, which probably led to limited ruminal disturbances and which might explain why mean daily rumen pH of goats offered H remained above 6.0.

The time spent standing and lying was only influenced by the diet after the afternoon feeding (1730 to 2330 h). As it has already been suggested, it is certainly due to the two intake periods, separated by resting and ruminating, observed with H, while goats receiving L performed a longer main meal, followed by resting and rumination.

High individual variability was observed during this experiment and seems to be typical of experiments dealing with subacute acidosis. For example, goat susceptibility to acidosis is highly variable: some goats probably did not suffer from

subacute acidosis with the H diet while others certainly did. Thus, to progress further in the analysis, results have to be interpreted individually, according to the detection of acidosis bouts and not only according to the percentage of concentrate in the diet.

Conclusion

Increasing the percentage of concentrate in the diet from 30% to 60% increased DMI and milk production and decreased milk fat content. Both diets led to short periods of time where rumen pH was below 6.0 but H induced lower mean rumen pH and was therefore more acidogenic than L. We can conclude from this experiment that goats fed a high-concentrate diet changed their nycthemeral activity patterns, particularly during the hours following feeding. They appeared to eat and ruminate more often but during shorter periods of time, which is probably due to satiety control. This adaptation could have limited the digestive disorders due to the rapid intake of a large amount of highly fermentable carbohydrates with the H diet. This adaptation might explain the small differences observed in rumen pH with the two diets.

Acknowledgements

We would like to thank J. Tessier, A. Aymard, M. Langot, G. Durandet, A. Severin and J. Kloster from the experimental farm at Grignon for the care of the goats and their technical assistance during the experiment. We thank C. Martin, L. Brossard and M. Fabre for their help in using pH probes. We also thank D. Sauvaut, P. Morand-Fehr and A. Ponter for their advice and rereading the manuscript.

References

Abijaoudé JA, Morand-Fehr P, Tessier J, Schmidely P and Sauvaut D 2000a. Diet effect on the daily feeding behaviour, frequency and characteristics of meals in dairy goats. *Livestock Production Science* 64, 29–37.

Abijaoudé JA, Morand-Fehr P, Tessier J, Schmidely P and Sauvaut D 2000b. Influence of forage: concentrate ratio and type of starch in the diet on feeding behaviour, dietary preferences, digestion, metabolism and performance of dairy goats in mid lactation. *Animal Science* 71, 359–368.

Allen MS 2000. Effects of diet on short-term regulation of feed intake by lactating dairy cattle. *Journal of Dairy Science* 83, 1598–1624.

Baumont R, Dulphy JP, Sauvaut D, Tran G, Meschy F, Aufrère J, Peyraud JL and Champciaux P 2007. Les tables de la valeur des aliments. Alimentation des bovins, ovins et caprins. Besoin des animaux-Valeurs des aliments. Tables INRA 2007. Institut National de la Recherche Agronomique (INRA), Paris, France, pp. 181–286.

Beauchemin KA, Kachanosky RG, Schaalje GB and Buchanan-Smith JG 1990. Characterizing rumination patterns of dairy cows using spectral analysis. *Journal of Animal Science* 68, 3163–3170.

Braun U, Rihs T and Schefer U 1992. Ruminal lactic acidosis in sheep and goats. *The Veterinary Record* 130, 343–349.

Brossard L, Fabre M, Martin C and Michalet-Doreau B 2003. Validation of continuous ruminal pH measurements by indwelling probes (poster). Conference of Gastrointestinal Function, Chicago, p. 25.

Buño L 1975. Rôle de l'acide DL-lactique dans le contrôle de l'ingestion alimentaire chez le mouton. *Annales de Recherches Vétérinaires* 6, 325–335.

Deswysen AG, Dutilleul P, Godfrin JP and Ellis WC 1993. Nycterohemeral eating and ruminating patterns in heifers fed grass or corn-silage – analysis by finite Fourier-transform. *Journal of Animal Science* 71, 2739–2747.

DeVries TJ, von Keyserlingk MAG and Beauchemin KA 2003. Short communication: diurnal feeding pattern of lactating dairy cows. *Journal of Dairy Science* 86, 4079–4082.

Dulphy JP and Faverdin P 1987. L'ingestion alimentaire chez les ruminants: modalités et phénomènes associés. *Reproduction, Nutrition, Development* 27, 129–155.

Forbes JM 2007. A personal view of how ruminant animals control their intake and choice of food: minimal total discomfort. *Nutrition Research Reviews* 20, 132–146.

Jarrige R, Dulphy JP, Faverdin P, Baumont R and Demarquilly C 1995. Activités d'ingestion et de rumination. In *Nutrition des ruminants domestiques. Ingestion et digestion* (ed. M Journet), pp. 123–181. Institut National de la Recherche Agronomique (INRA), Paris, France.

Kawas JR, Lopes J, Danelon DL and Lu CD 1991. Influence of forage-to-concentrate ratio on intake, digestibility, chewing and milk production of dairy goats. *Small Ruminant Research* 4, 11–18.

Keunen JE, Plaizier JC, Kyriazakis L, Duffield TF, Widowski TM, Lindinger MI and McBride BW 2002. Effects of a subacute ruminal acidosis model on the diet selection of dairy cows. *Journal of Dairy Science* 85, 3304–3313.

Krause KM and Oetzel GR 2005. Inducing subacute ruminal acidosis in lactating dairy cows. *Journal of Dairy Science* 88, 3633–3639.

Maekawa M, Beauchemin KA and Christensen DA 2002. Effect of concentrate level and feeding management on chewing activities, saliva production, and ruminal pH of lactating dairy cows. *Journal of Dairy Science* 85, 1165–1175.

Manson FJ and Leaver JD 1988. The influence of concentrate amount on locomotion and clinical lameness in dairy cattle. *Animal Production* 47, 185–190.

Martin C, Brossard L and Doreau M 2006. Mécanismes d'apparition de l'acidose ruminale latente et conséquences physiopathologiques et zootechniques. *Productions Animales* 19, 93–108.

Mitlöhrer FM, Morrow-Tesch JL, Wilson SC, Dailey JW and McGlone JJ 2001. Behavioral sampling techniques for feedlot cattle. *Journal of Animal Science* 79, 1189–1193.

Nocek JE 1997. Bovine acidosis: implications on laminitis. *Journal of Dairy Science* 80, 1005–1028.

Oetzel GR 2000. Clinical aspects of ruminal acidosis in dairy cattle. In the 33rd Annual Conference, American Association of Bovine Practitioners, Rapid City, South Dakota, USA, 14pp.

Owens FN, Secrist DS, Hill WJ and Gill DR 1998. Acidosis in cattle: a review. *Journal of Animal Science* 76, 275–286.

Peyraud JL and Apper-Bossard E 2006. L'acidose latente chez la vache laitière. *Productions Animales* 19, 79–92.

Rapetti L, Bava L, Tamburini A and Crovetto GM 2005. Feeding behaviour, digestibility, energy balance and productive performance of lactating goats fed forage-based and forage-free diets. *Italian Journal of Animal Science* 4, 71–83.

Rémond B, Brugère H, Poncet C and Baumont R 1995. Le contenu du réticulorumen. In *Nutrition des ruminants domestiques, ingestion et digestion* (ed. YRR Jarrige, C Demarquilly, MH Farce and M Journet), pp. 253–298. Institut National de la Recherche Agronomique (INRA), Paris, France.

Sauvaut D 1992. Systemic modeling in nutrition. *Reproduction, Nutrition, Development* 32, 217–230.

Sauvaut D, Meschy F and Mertens D 1999. Les composantes de l'acidose ruminale et les effets acidogènes des rations. *Productions Animales* 12, 49–60.

Sauvaut D, Giger-Reverdin S and Meschy F 2006. Le contrôle de l'acidose ruminale latente. *Productions Animales* 19, 69–78.

Sauvaut D, Giger-Reverdin S and Meschy F 2007. Alimentation des caprins. Alimentation des bovins, ovins et caprins. Besoin des animaux-Valeurs des aliments. Tables INRA 2007. Institut National de la Recherche Agronomique (INRA), Paris, France, pp. 137–148.

Sawyer DC 1998. Pain control in small-animal patients. *Applied Animal Behaviour Science* 59, 135–146.

Susin I, Loerch SC and McClure KE 1995. Effects of feeding a high-grain diet at a restricted intake on lactation performance and rebreeding of ewes. *Journal of Animal Science* 73, 3199–3205.