

Research Article

**New Criteria for Meromorphic Multivalent
Alpha-Convex Functions****Muhammad Arif,¹ Maslina Darus,² Fazal Ghani,¹ and Saeed Islam¹**¹ Department of Mathematics, Abdul Wali Khan University Mardan, Khyber Pakhtunkhwa, Pakistan² School of Mathematical Sciences, Faculty of Science and Technology, Universiti Kebangsaan Malaysia, 43600 Bangi, Selangor, MalaysiaCorrespondence should be addressed to Maslina Darus; maslina@ukm.my

Received 9 March 2013; Revised 14 July 2013; Accepted 15 July 2013

Academic Editor: Zhihua Zhang

Copyright © 2013 Muhammad Arif et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

The aim of the present paper is to obtain sufficient condition for the class of meromorphic p -valent alpha convex functions of order ξ and then to study mapping properties of the newly defined integral operators. Many known results appeared as special consequences of our work.**1. Introduction**Let $\Sigma_p(n)$ denote the class of meromorphic functions $f(z)$ normalized by

$$f(z) = z^{-p} + \sum_{k=n}^{\infty} a_k z^{k-p+1}, \quad (p \in \mathbb{N}), \quad (1)$$

which are analytic and p -valent in the punctured unit disk $\mathbb{U}^* = \{z : 0 < |z| < 1\}$. In particular, $\Sigma_1(n) = \Sigma(n)$, $\Sigma_p(1) = \Sigma_p$, and $\Sigma_1(1) = \Sigma$. For λ which is real with $|\lambda| < \pi/2$, $\alpha \geq 0$, $0 \leq \xi < p$, and $n, p \in \mathbb{N}$, we denote by $\Sigma\mathcal{S}_p(\lambda, n, \xi)$, $\Sigma\mathcal{C}_p(\lambda, n, \xi)$, and $\Sigma\mathcal{M}_p(\lambda, n, \alpha, \xi)$, the subclasses of $\Sigma_p(n)$ consisting of all meromorphic p -valent functions of the form (1) which are defined, respectively, by

$$\begin{aligned} & -\operatorname{Re} e^{i\lambda} \frac{zf'(z)}{f(z)} > \xi \cos \lambda, \quad (z \in \mathbb{U}^*), \\ & -\operatorname{Re} e^{i\lambda} \frac{(zf'(z))'}{f'(z)} > \xi \cos \lambda, \quad (z \in \mathbb{U}^*), \\ & -\operatorname{Re} e^{i\lambda} \left\{ (1-\alpha) \frac{zf'(z)}{f(z)} + \alpha \frac{(zf'(z))'}{f'(z)} \right\} \\ & > \xi \cos \lambda, \quad (z \in \mathbb{U}^*). \end{aligned} \quad (2)$$

Making $\lambda = 0$, $n = 1$ in (2), we get the well-known subclasses of Σ_p consisting of meromorphic p -valent functions which are starlike, convex, and alpha convex of order ξ ($0 \leq \xi < p$), respectively. For details of the classes defined by (2) and related topics, we refer the reader to the work of Aouf and Hossen [1], Aouf and Srivastava [2], Ali and Ravichandran [3], Goyal and Prajapat [4], Joshi and Srivastava [5], Liu and Srivastava [6], Raina and Srivastava [7], Xu and Yang [8], and Owa et al. [9].

For $f(z) \in \Sigma$, Wang et al. [10] and Nehari and Netanyahu [11] introduced and studied the subclass $\Sigma_N(\tau)$ of Σ consisting of functions $f(z)$ satisfying

$$-\operatorname{Re} \frac{(zf'(z))'}{f'(z)} < \tau, \quad (\tau > 1, z \in \mathbb{U}^*). \quad (3)$$

We now extended this concept to define a subclass $\Sigma\mathcal{N}_p(\lambda, n, \alpha, \tau)$ of $\Sigma_p(n)$ consisting of functions $f(z)$ of the form (1) satisfying

$$-\operatorname{Re} e^{i\lambda} \left((1-\alpha) \frac{zf'(z)}{f(z)} + \alpha \frac{(zf'(z))'}{f'(z)} \right) < \tau \cos \lambda, \quad (\tau > p, z \in \mathbb{U}^*). \quad (4)$$

For $\alpha = 0$ and $\alpha = 1$ in (4), we obtain the classes $\Sigma\mathcal{N}\mathcal{C}_p(\lambda, n, \tau)$ and $\Sigma\mathcal{N}\mathcal{S}_p(\lambda, n, \tau)$ of $\Sigma_p(n)$, respectively, studied by Arif [12]; also see [13, 14].

Integral operators for different classes of analytic, univalent, and multivalent functions in the open unit disk are studied by various authors; see [15–21]. We now define the following general integral operator of meromorphic p -valent functions:

$$G_m(z) = I_{m,p}(\delta, \alpha_j; f_j(z)) = \left\{ \frac{\delta}{z^{p+1}} \int_0^z t^{\delta-1} \prod_{j=1}^m (t^p (f_j(t)))^{\alpha_j} dt \right\}^{1/\delta} \quad (5)$$

For $\delta = 1$, we obtain the integral operator $I_{m,p}(f_j(z))$ studied recently in [22, 23], and, further for $p = 1$, we obtain the integral operator introduced and studied by Mohammed and Darus [24].

Sufficient conditions were studied by various authors for different subclasses of analytic and multivalent functions; for some of the related work see [25–27]. The object of the present paper is to obtain sufficient conditions for the class $\Sigma\mathcal{M}_p(\lambda, n, \alpha, \xi)$ and then study mapping properties of the integral operator given by (5). We also consider some special cases of our results which lead to various interesting corollaries and relevance of some of these results with other known results which are also mentioned.

We will assume throughout our discussion, unless otherwise stated, that λ is real with $|\lambda| < \pi/2$, $0 \leq \xi < p$, $\tau > p$, $p, n \in \mathbb{N}$, $\alpha_j > 0$ for $j \in \{1, \dots, m\}$, $\delta > 0$, $\alpha \geq 0$, and

$$J_\alpha(f) = (1 - \alpha) \frac{zf'(z)}{f(z)} + \alpha \frac{(zf'(z))'}{f'(z)}. \quad (6)$$

To obtain our main results, we need the following lemmas.

Lemma 1 (see [27]). *If $q(z) \in \Sigma(n)$ with $n \geq 1$ and satisfies the condition*

$$|z^2 q'(z) + 1| < \frac{n}{\sqrt{n^2 + 1}} \quad (z \in \mathbb{U}^*), \quad (7)$$

then

$$q(z) \in \Sigma\mathcal{S}(n). \quad (8)$$

Lemma 2 (see [28]). *Let $\psi : \mathbb{C}^2 \rightarrow \mathbb{C}$ satisfy the following condition:*

$$\text{Re} \{ \psi(is, t) \} \leq 0, \quad \left(s, t \in \mathbb{R}; t \leq -\frac{|a + is|^2}{2}, \text{Re}(a) > 0 \right). \quad (9)$$

If the function $h(z) = a + h_1z + h_2z^2 + \dots$ is analytic in $\mathbb{U} = \mathbb{U}^ \setminus \{0\}$ and*

$$\text{Re} \{ \psi(h(z), zh'(z)) \} > 0, \quad (z \in \mathbb{U}), \quad (10)$$

then

$$\text{Re} h(z) > 0, \quad (z \in \mathbb{U}). \quad (11)$$

2. Sufficiency Criteria for the Class

$$\Sigma\mathcal{M}_p(\lambda, n, \alpha, \xi)$$

In this section we establish a new sufficiency criteria for the subclass $\Sigma\mathcal{M}_p(\lambda, n, \alpha, \xi)$ of $\Sigma_p(n)$.

Theorem 3. *If $f(z) \in \Sigma_p(n)$ satisfies*

$$\left| \left\{ \left(z^p f(z) \left(\frac{-zf'(z)}{pf(z)} \right)^\alpha \right)^{e^{i\lambda}/(p-\xi) \cos \lambda} e^{i\lambda} J_\alpha(f) + \xi \cos \lambda + ip \sin \lambda \right\} + (p - \xi) \cos \lambda \right| < \frac{n}{\sqrt{n^2 + 1}} (p - \xi) \cos \lambda \quad (z \in \mathbb{U}^*), \quad (12)$$

then $f(z) \in \Sigma\mathcal{M}_p(\lambda, n, \alpha, \xi)$, where $J_\alpha(f)$ is given by (6).

Proof. Let us set a function $q(z)$ by

$$q(z) = \frac{1}{z} \left(z^p f(z) \left(\frac{-zf'(z)}{pf(z)} \right)^\alpha \right)^{e^{i\lambda}/(p-\xi) \cos \lambda} = \frac{1}{z} + \frac{\alpha e^{i\lambda} a_n b_n}{(p - \xi) \cos \lambda} z^n + \dots \quad (13)$$

for $f(z) \in \Sigma_p(n)$. Then clearly (13) shows that $q(z) \in \Sigma(n)$.

Logarithmic differentiating of (13) gives

$$\frac{q'(z)}{q(z)} = \frac{e^{i\lambda}}{(p - \xi) \cos \lambda} \times \left[(1 - \alpha) \frac{f'(z)}{f(z)} + \alpha \frac{(zf'(z))'}{zf'(z)} + \frac{p}{z} \right] - \frac{1}{z} \quad (14)$$

which further implies

$$\left| z^2 q'(z) + 1 \right| = \left| \left(z^p f(z) \left(\frac{-zf'(z)}{pf(z)} \right)^\alpha \right)^{e^{i\lambda}/(p-\xi) \cos \lambda} \frac{e^{i\lambda}}{(p - \xi) \cos \lambda} \times [J_\alpha(f) + \xi \cos \lambda + ip \sin \lambda] + 1 \right|. \quad (15)$$

Thus using (12), we get

$$\left| z^2 q'(z) + 1 \right| \leq \frac{n}{\sqrt{n^2 + 1}}, \quad (z \in \mathbb{U}^*). \quad (16)$$

Therefore by Lemma 1, we have $q(z) \in \Sigma\mathcal{S}(n)$.

From (14), we can write

$$\frac{zq'(z)}{q(z)} = \frac{1}{(p-\xi)\cos\lambda} \left[e^{i\lambda} J_\alpha(f) + \xi \cos\lambda + ip \sin\lambda \right]. \tag{17}$$

Since $q(z) \in \Sigma\mathcal{S}(n)$, it implies that $\operatorname{Re}(-zq'(z)/q(z)) > 0$. Therefore, we get

$$\begin{aligned} & \frac{1}{(p-\xi)\cos\lambda} \left[-\operatorname{Re} e^{i\lambda} J_\alpha(f) - \xi \cos\lambda \right] \\ & = \operatorname{Re} \left(-\frac{zq'(z)}{q(z)} \right) > 0 \end{aligned} \tag{18}$$

or

$$-\operatorname{Re} e^{i\lambda} J_\alpha(f) > \xi \cos\lambda. \tag{19}$$

And therefore $f(z) \in \Sigma\mathcal{M}_p(\lambda, n, \alpha, \xi)$.

By taking $\alpha = 0$ and $\alpha = 1$ in Theorem 3, we obtain Corollaries 4 and 5, respectively, proved by Arif [12]. \square

Corollary 4. *If $f(z) \in \Sigma_p(n)$ satisfies*

$$\begin{aligned} & \left| (z^p f(z))^{e^{i\lambda}/(p-\xi)\cos\lambda} \left\{ e^{i\lambda} \frac{zf'(z)}{f(z)} + \xi \cos\lambda + ip \sin\lambda \right\} \right. \\ & \left. + (p-\xi)\cos\lambda \right| < \frac{n(p-\xi)\cos\lambda}{\sqrt{n^2+1}} \quad (z \in \mathbb{U}^*), \end{aligned} \tag{20}$$

then $f(z) \in \Sigma\mathcal{D}_p(\lambda, n, \xi)$.

Corollary 5. *If $f(z) \in \Sigma_p(n)$ satisfies*

$$\begin{aligned} & \left| \left(\frac{z^{p+1} f'(z)}{-p} \right)^{e^{i\lambda}/(p-\xi)\cos\lambda} \right. \\ & \left. \times \left\{ e^{i\lambda} \left(\frac{zf''(z)}{f'(z)} + 1 \right) + \xi \cos\lambda + ip \sin\lambda \right\} + (p-\xi)\cos\lambda \right| \\ & < \frac{n}{\sqrt{n^2+1}} (p-\xi)\cos\lambda, \quad (z \in \mathbb{U}^*), \end{aligned} \tag{21}$$

then $f(z) \in \Sigma\mathcal{C}_p(\lambda, n, \xi)$.

Remarks. We note that by simple computation (13) gives

$$f(z) = \left[-\frac{p}{\alpha} \int_0^z \frac{(tq(t))^{(p-\xi)\cos\lambda/\alpha e^{i\lambda}}}{t^{p/\alpha+1}} dt \right]^\alpha. \tag{22}$$

By taking suitable meromorphic starlike function for $q(z)$ in (22) such as $q(z) = (1-z^n)^2/z$, which satisfies the inequality of Lemma 1, we can conclude that the function f of (22) is the subclass of a meromorphic function.

3. Some Properties of the Integral Operator

$G_m(z)$

In this section, we discuss some mapping properties of the integral operator $G_m(z)$.

Theorem 6. *For $j \in \{1, \dots, m\}$, let $f_j(z) \in \Sigma_p(n)$ and satisfy (20). If*

$$\sum_{j=1}^m \alpha_j < \frac{(p+1)(2-\delta)}{(p-\xi)}, \quad 0 < \delta < 2, \tag{23}$$

then $G_m(z) \in \Sigma\mathcal{N}_p(\lambda, n, \delta, \eta)$ with $\eta > p$, and $G_m(z)$ is given by (5).

Proof. From (5), we obtain

$$\begin{aligned} & \delta z^{p+1} G_m^{\delta-1}(z) G'_m(z) + (p+1) z^p G_m^\delta(z) \\ & = \delta z^{\delta-1} \prod_{j=1}^m (z^p f_j(z))^{\alpha_j}. \end{aligned} \tag{24}$$

Dividing both sides by $z^p G_m^{\delta-1}(z)$, we have

$$\begin{aligned} & \delta z G'_m(z) + (p+1) G_m(z) \\ & = \delta z^{\delta-p-1} G_m^{1-\delta}(z) \prod_{j=1}^m (z^p f_j(z))^{\alpha_j}. \end{aligned} \tag{25}$$

Differentiating again logarithmically, we have

$$\begin{aligned} & \frac{\delta z G''_m(z) + (p+\delta+1) G'_m(z)}{\delta z G'_m(z) + (p+1) G_m(z)} \\ & = (\delta-p-1) \frac{1}{z} + (1-\delta) \frac{G'_m(z)}{G_m(z)} \\ & \quad + \sum_{j=1}^m \alpha_j \left(\frac{f'_j(z)}{f_j(z)} + \frac{p}{z} \right). \end{aligned} \tag{26}$$

Now by simple computation, we get

$$\begin{aligned} & \left(1 - \frac{1}{\delta} \right) \frac{z G'_m(z)}{G_m(z)} + \frac{1}{\delta} \frac{(z G'_m(z))'}{G'_m(z)} \\ & = \frac{1}{\delta} \sum_{j=1}^m \alpha_j \left(\frac{z f'_j(z)}{f_j(z)} + p \right) \\ & \quad - \frac{1}{\delta} (2+2p-\delta) + \frac{G_m(z)}{z G'_m(z)} \\ & \quad \times \left[(p+1) \sum_{j=1}^m \alpha_j \left(\frac{z f'_j(z)}{f_j(z)} + p \right) + (p+1)(\delta-p-1) \right], \end{aligned} \tag{27}$$

or equivalently we have

$$\begin{aligned}
 & -e^{i\lambda} \left\{ \left(1 - \frac{1}{\delta} \right) \frac{zG'_m(z)}{G_m(z)} + \frac{1}{\delta} \frac{(zG'_m(z))'}{G'_m(z)} \right\} \\
 &= \frac{1}{\delta} \sum_{j=1}^m \alpha_j \left(-e^{i\lambda} \frac{zf'_j(z)}{f_j(z)} - pe^{i\lambda} \right) \\
 &+ \frac{1}{\delta} (2 + 2p - \delta) e^{i\lambda} + \frac{G_m(z)}{zG'_m(z)} \\
 &\times \left[(p + 1 - \delta) - \sum_{j=1}^m \alpha_j \left(\frac{zf'_j(z)}{f_j(z)} + p \right) \right] (p + 1) e^{i\lambda}.
 \end{aligned} \tag{28}$$

By taking real part on both sides, we obtain

$$\begin{aligned}
 & -\operatorname{Re} e^{i\lambda} \left\{ \left(1 - \frac{1}{\delta} \right) \frac{zG'_m(z)}{G_m(z)} + \frac{1}{\delta} \frac{(zG'_m(z))'}{G'_m(z)} \right\} \\
 &= \frac{1}{\delta} \sum_{j=1}^m \alpha_j \left(-\operatorname{Re} e^{i\lambda} \frac{zf'_j(z)}{f_j(z)} - p \cos \lambda \right) \\
 &+ \frac{1}{\delta} (2 + 2p - \delta) \cos \lambda + \operatorname{Re} \frac{G_m(z)}{zG'_m(z)} \\
 &\times \left[(p + 1 - \delta) - \sum_{j=1}^m \alpha_j \left(\frac{zf'_j(z)}{f_j(z)} + p \right) \right] (p + 1) e^{i\lambda},
 \end{aligned} \tag{29}$$

which further implies that

$$\begin{aligned}
 & -\operatorname{Re} e^{i\lambda} \left\{ \left(1 - \frac{1}{\delta} \right) \frac{zG'_m(z)}{G_m(z)} + \frac{1}{\delta} \frac{(zG'_m(z))'}{G'_m(z)} \right\} \\
 &\leq \frac{1}{\delta} \sum_{j=1}^m \alpha_j \left(-\operatorname{Re} e^{i\lambda} \frac{zf'_j(z)}{f_j(z)} - p \cos \lambda \right) \\
 &+ \frac{1}{\delta} (2 + 2p - \delta) \cos \lambda \\
 &+ \left| \frac{G_m(z)}{zG'_m(z)} \right| \left[(p + 1 - \delta) \right. \\
 &\quad \left. - \sum_{j=1}^m \alpha_j \left(\frac{zf'_j(z)}{f_j(z)} + p \right) \right] (p + 1) e^{i\lambda}.
 \end{aligned} \tag{30}$$

Let

$$\begin{aligned}
 \eta &= \left| \frac{G_m(z)}{zG'_m(z)} \left[(p + 1 - \delta) \right. \right. \\
 &\quad \left. \left. - \sum_{j=1}^m \alpha_j \left(\frac{zf'_j(z)}{f_j(z)} + p \right) \right] (p + 1) e^{i\lambda} \right| \\
 &+ \frac{1}{\delta} (2 + 2p - \delta) \\
 &+ \frac{1}{\delta} \sum_{j=1}^m \alpha_j \left(-\frac{1}{\cos \lambda} \operatorname{Re} e^{i\lambda} \frac{zf'_j(z)}{f_j(z)} - p \right).
 \end{aligned} \tag{31}$$

Clearly we have

$$\begin{aligned}
 \eta &> \frac{1}{\delta} (2 + 2p - \delta) \\
 &+ \frac{1}{\delta \cos \lambda} \sum_{j=1}^m \alpha_j \left(-\operatorname{Re} e^{i\lambda} \frac{zf'_j(z)}{f_j(z)} - p \cos \lambda \right).
 \end{aligned} \tag{32}$$

Then by using (23) and Theorem 3 with $\alpha = 0$, we obtain

$$\eta > \frac{1}{\delta} \left[\sum_{j=1}^m \alpha_j (\xi - p) + (2 + 2p - \delta) \right] > p. \tag{33}$$

Therefore $G_m(z) \in \Sigma \mathcal{N}'_p(\lambda, n, \delta, \eta)$ with $\eta > p$.

Making $\delta = 1$ in Theorem 6, we have the following. \square

Corollary 7. For $j \in \{1, \dots, m\}$, let $f_j(z) \in \Sigma_p(n)$ and satisfy (20). If

$$\sum_{j=1}^m \alpha_j < \frac{(p + 1)}{(p - \xi)}, \tag{34}$$

then $I_{m,p}(f_j(z)) \in \Sigma \mathcal{N}\mathcal{C}_p(\lambda, n, \eta)$ with $\eta > p$.

Theorem 8. For $j \in \{1, \dots, m\}$, let $f_j(z) \in \Sigma \mathcal{S}_p(0, n, \xi)$. If

$$\sum_{j=1}^m \alpha_j < \frac{p + 1 - \delta}{p - \xi}, \tag{35}$$

then $G_m(z) \in \Sigma \mathcal{S}_p(0, n, \rho)$, where $0 < \rho \leq p$ with $\rho = p + 1 - \delta + (\xi - p) \sum_{j=1}^m \alpha_j$.

Proof. From (26), we obtain

$$\begin{aligned}
 & \frac{\delta (zG''_m(z)/G'_m(z) + 1) + (p + 1) zG'_m(z)}{\delta (zG'_m(z)/G_m(z) + (p + 1)) G_m(z)} \\
 &= (\delta - p - 1) + (1 - \delta) \frac{zG'_m(z)}{G_m(z)} \\
 &+ \sum_{j=1}^m \alpha_j \left(\frac{zf'_j(z)}{f_j(z)} + p \right).
 \end{aligned} \tag{36}$$

Let

$$-e^{i\lambda} \frac{zG'_m(z)}{G_m(z)} = q(z) = ((p - \rho) p(z) + \rho) \cos \lambda + i \sin \lambda, \tag{37}$$

such that $q(z)$ is analytic in \mathbb{U}^* with $q(0) = p$. Then (36) can be written as

$$q(z) + \frac{zq'(z) e^{i\lambda}}{(p+1) e^{i\lambda} - \delta q(z)} = \frac{(p+1-\delta) e^{i\lambda}}{\delta} + \frac{1}{\delta} \sum_{j=1}^m \alpha_j \left(-e^{i\lambda} \frac{zf'_j(z)}{f_j(z)} - p e^{i\lambda} \right). \tag{38}$$

Taking real part on both sides, we have

$$\begin{aligned} \operatorname{Re} \left(q(z) + \frac{zq'(z) e^{i\lambda}}{(p+1) e^{i\lambda} - \delta q(z)} \right) &= \frac{p+1-\delta}{\delta} \cos \lambda \\ &+ \frac{1}{\delta} \sum_{j=1}^m \alpha_j \left(-\operatorname{Re} e^{i\lambda} \frac{zf'_j(z)}{f_j(z)} - p \cos \lambda \right) \\ &> \frac{1}{\delta} \left(p+1-\delta - (p-\xi) \sum_{j=1}^m \alpha_j \right) \cos \lambda > 0, \end{aligned} \tag{39}$$

where we have used (35) and the assumption that $f_j(z) \in \Sigma_{\mathcal{S}_p}(\lambda, n, \xi)$. Let us put

$$\psi(u, v) = u + \frac{v e^{i\lambda}}{(p+1) e^{i\lambda} - \delta u}. \tag{40}$$

Then, for $s, t \in \mathbb{R}$ such that $t \leq -(p^2 + s^2)/2$, we have

$$\begin{aligned} \operatorname{Re}(\psi(is, t)) &= \operatorname{Re} \left\{ is + \frac{t e^{i\lambda}}{(p+1) e^{i\lambda} - i\delta s} \right\} \\ &= \frac{[(p+1) - \delta s \sin \lambda] (p^2 + s^2)}{2 [(p+1)^2 \cos^2 \lambda + \delta^2 s^2 (1 - (p+1) \sin \lambda)^2]} < 0. \end{aligned} \tag{41}$$

Thus using Lemma 2, we conclude that $\operatorname{Re} q(z) > 0$, which is equivalent to

$$-\operatorname{Re} e^{i\lambda} \frac{zG'_m(z)}{G_m(z)} > \rho \cos \lambda. \tag{42}$$

That is, $G_m(z) \in \Sigma_{\mathcal{S}_p}(\lambda, n, \rho)$. □

Acknowledgment

The work presented here was partially supported by LRGS/TD/2011/UKM/ICT/03/02.

References

- [1] M. K. Aouf and H. M. Hossen, "New criteria for meromorphic p -valent starlike functions," *Tsukuba Journal of Mathematics*, vol. 17, no. 2, pp. 481–486, 1993.
- [2] M. K. Aouf and M. H. Srivastava, "A new criteria for meromorphically p -valent convex functions of order α ," *Mathematical Sciences Research Hot-Line*, vol. 1, no. 8, pp. 7–12, 1997.
- [3] R. M. Ali and V. Ravichandran, "Classes of meromorphic α -convex functions," *Taiwanese Journal of Mathematics*, vol. 14, no. 4, pp. 1479–1490, 2010.
- [4] S. P. Goyal and J. K. Prajapat, "A new class of meromorphic multivalent functions involving certain linear operator," *Tamsui Oxford Journal of Mathematical Sciences*, vol. 25, no. 2, pp. 167–176, 2009.
- [5] S. B. Joshi and H. M. Srivastava, "A certain family of meromorphically multivalent functions," *Computers & Mathematics with Applications*, vol. 38, no. 3-4, pp. 201–211, 1999.
- [6] J. L. Liu and H. M. Srivastava, "A linear operator and associated families of meromorphically multivalent functions," *Journal of Mathematical Analysis and Applications*, vol. 259, no. 2, pp. 566–581, 2001.
- [7] R. K. Raina and H. M. Srivastava, "A new class of meromorphically multivalent functions with applications to generalized hypergeometric functions," *Mathematical and Computer Modelling*, vol. 43, no. 3-4, pp. 350–356, 2006.
- [8] N. Xu and D. Yang, "On starlikeness and close-to-convexity of certain meromorphic functions," *Journal of the Korea Society of Mathematical Education B*, vol. 10, no. 1, pp. 566–581, 2003.
- [9] S. Owa, H. E. Darwish, and M. K. Aouf, "Meromorphic multivalent functions with positive and fixed second coefficients," *Mathematica Japonica*, vol. 46, no. 2, pp. 231–236, 1997.
- [10] Z. G. Wang, Y. Sun, and Z. H. Zhang, "Certain classes of meromorphic multivalent functions," *Computers & Mathematics with Applications*, vol. 58, no. 7, pp. 1408–1417, 2009.
- [11] Z. Nehari and E. Netanyahu, "On the coefficients of meromorphic schlicht functions," *Proceedings of the American Mathematical Society*, vol. 8, pp. 15–23, 1957.
- [12] M. Arif, "On certain sufficiency criteria for p -valent meromorphic spiralike functions," *Abstract and Applied Analysis*, vol. 2012, Article ID 837913, 10 pages, 2012.
- [13] Z. G. Wang, Z. H. Liu, and R. G. Xiang, "Some criteria for meromorphic multivalent starlike functions," *Applied Mathematics and Computation*, vol. 218, no. 3, pp. 1107–1111, 2011.
- [14] Z. G. Wang, Z. H. Liu, and A. Cățaș, "On neighborhoods and partial sums of certain meromorphic multivalent functions," *Applied Mathematics Letters*, vol. 24, no. 6, pp. 864–868, 2011.
- [15] B. A. Frasin, "New general integral operators of p -valent functions," *Journal of Inequalities in Pure and Applied Mathematics*, vol. 10, no. 4, article 109, 2009.
- [16] M. Arif, K. I. Noor, and F. Ghani, "Some properties of an integral operator defined by convolution," *Journal of Inequalities and Applications*, vol. 2012, article 13, 6 pages, 2012.
- [17] K. I. Noor and M. Arif, "Mapping properties of an integral operator," *Applied Mathematics Letters*, vol. 25, no. 11, pp. 1826–1829, 2012.
- [18] K. I. Noor, M. Arif, and A. Muhammad, "Mapping properties of some classes of analytic functions under an integral operator," *Journal of Mathematical Inequalities*, vol. 4, no. 4, pp. 593–600, 2010.

- [19] N. Breaz, V. Pescar, and D. Breaz, "Univalence criteria for a new integral operator," *Mathematical and Computer Modelling*, vol. 52, no. 1-2, pp. 241–246, 2010.
- [20] B. A. Frasin, "Convexity of integral operators of p -valent functions," *Mathematical and Computer Modelling*, vol. 51, no. 5-6, pp. 601–605, 2010.
- [21] B. A. Frasin, "Some sufficient conditions for certain integral operators," *Journal of Mathematical Inequalities*, vol. 2, no. 4, pp. 527–535, 2008.
- [22] A. Mohammed and M. Darus, "Some properties of certain integral operators on new subclasses of analytic functions with complex order," *Journal of Applied Mathematics*, vol. 2012, Article ID 161436, 9 pages, 2012.
- [23] A. Mohammed and M. Darus, "The order of starlikeness of new p -valent meromorphic functions," *International Journal of Mathematical Analysis*, vol. 6, no. 27, pp. 1329–1340, 2012.
- [24] A. Mohammed and M. Darus, "Starlikeness properties of a new integral operator for meromorphic functions," *Journal of Applied Mathematics*, vol. 2011, Article ID 804150, 8 pages, 2011.
- [25] M. Arif, I. Ahmad, M. Raza, and K. Khan, "Sufficient condition of a subclass of analytic functions defined by Hadamard product," *Life Science Journal*, vol. 9, no. 4, pp. 2487–2489, 2012.
- [26] M. Arif, M. Raza, S. Islam, J. Iqbal, and F. Faizullah, "Some sufficient conditions for spirallike functions with argument properties," *Life Science Journal*, vol. 9, no. 4, pp. 3770–3773, 2012.
- [27] H. Al-Amiri and P. T. Mocanu, "Some simple criteria of starlikeness and convexity for meromorphic functions," *Mathematica*, vol. 37, no. 60, pp. 11–21, 1995.
- [28] S. S. Miller and P. T. Mocanu, *Differential Subordinations: Theory and Applications*, vol. 225 of *Monographs and Textbooks in Pure and Applied Mathematics*, Marcel Dekker, New York, NY, USA, 2000.



Hindawi

Submit your manuscripts at
<http://www.hindawi.com>

