

1 **Distinct roles of DBHS family members in the circadian transcriptional feedback**
2 **loop.**

3
4 Elzbieta Kowalska¹, Jürgen A. Ripperger^{2§}, Christine Muheim^{1§}, Bert Maier³, Yasuyuki
5 Kurihara⁴, Archa H. Fox⁵, Achim Kramer³, and Steven A. Brown^{1*}

6
7 ¹Institute of Pharmacology and Toxicology, University of Zurich, Winterthurerstrasse
8 190, 8057 Zurich, Switzerland.

9 ²Department of Biology, Unit for Biochemistry, University of Fribourg, Chemin de Musée
10 5, 1700 Fribourg, Switzerland

11 ³Laboratory of Chronobiology, Institute of Medical Immunology, Charite
12 Universitätsmedizin, Berlin, Germany

13 ⁴Department of Natural Environment and Information, Yokohama National University,
14 Tokiwadai 79-7, Hodogaya-ku, Yokohama 240-8501 Japan

15 ⁵Western Australian Institute for Medical Research and Centre For Medical Research,
16 University of Western Australia, Crawley 6009 Australia.

17
18 *Address correspondence to steven.brown@pharma.uzh.ch

19 §These authors contributed equally to this publication.

20
21 Key words: DBHS, NOPS, NONO, SFPQ, PSPC1, Paraspeckle, Transcription,
22 Coregulator

23
24 **Factors interacting with core circadian clock components are essential to achieve**
25 **transcriptional feedback necessary for metazoan clocks. Here we show that all**
26 **three members of the *Drosophila* Behavior Human Splicing (DBHS) family of**
27 **RNA-binding proteins play a role in the mammalian circadian oscillator,**
28 **abrogating or altering clock function when overexpressed or depleted in cells.**
29 **Although these proteins are members of so-called nuclear paraspeckles,**
30 **depletion of paraspeckles themselves via silencing of the structural non-coding**
31 **RNA (ncRNA) *Neat1* did not affect overall clock function, suggesting that**
32 **paraspeckles are not required for DBHS-mediated circadian effects. Instead, we**
33 **show that the proteins bound to circadian promoter DNA in a fashion that**
34 **required the PERIOD (PER) proteins, and potently repressed E box-mediated**
35 **transcription but not CMV promoter-mediated transcription when exogenously**

36 **recruited. Nevertheless, mice with one or both copies of these genes deleted**
37 **show only small changes in period length or clock gene expression *in vivo*. Data**
38 **from transient transfections show that each of these proteins can either repress**
39 **or activate depending on the context. Taken together, our data suggest that all of**
40 **the DBHS family members serve overlapping or redundant roles as**
41 **transcriptional cofactors at circadian clock-regulated genes.**

42

43 **INTRODUCTION**

44 The circadian oscillator governs diurnal timing for most aspects of mammalian
45 physiology (7). Its mechanism is cell-autonomous, and consists of interlocked feedback
46 loops of circadian transcription, translation, and protein modification. In one loop, the
47 CLOCK/NPAS2 and BMAL1/ARNTL transcriptional activators drive expression of the
48 Period (*Per1* and *Per2*) and Cryptochrome (*Cry1* and *Cry2*) gene families, whose
49 products subsequently multimerize and repress their own transcription. In a second
50 loop, the transcriptional repressor REV-ERB α , whose transcription is also driven by
51 CLOCK and BMAL1, represses the expression of *Bmal1* itself (7). Beyond these
52 "dedicated" clock genes, a large number of other factors are necessary to the circadian
53 clock or for its regulation of physiology – kinases and phosphatases, chromatin
54 modifying factors, and other proteins (25, 33). We have shown previously that the
55 NONO protein in mammalian cells (or its ortholog NON-A in flies) plays such a role by
56 modulating PER-mediated transcriptional repression via unknown mechanisms (4).
57

58 NONO (also known as p54nrb in humans) has two RNA-binding (RRM) domains and
59 has been shown to regulate a variety of processes outside the circadian clock (38).
60 These include transcriptional activation and repression (16, 26), pre-mRNA processing
61 (19), and RNA transport in neurons (18). For example, NONO has been shown to
62 regulate the transcriptional activation of the TORC family of growth and metabolic
63 factors via recruitment of the RNA polymerase II (1). In an apparently unrelated nuclear
64 function, it also mediates the nuclear retention of edited RNAs in nuclear paraspeckles,
65 which are thought to be RNA holding structures (30). These structures contain the
66 NONO, SFPQ, and PSPC1 proteins, as well as the scaffolding ncRNA *Neat1* (3). Both
67 SFPQ and PSPC1 share significant structural and functional similarity to NONO, and for
68 this reason all three proteins have been grouped into the DBHS (*Drosophila* Behavior
69 Human Splicing) family of nuclear factors. Nevertheless, to date only NONO (4) and
70 SFPQ (10) have been implicated in the circadian clock mechanism.

71

72 Herein, we show that all three DBHS factors play important roles in the circadian clock
73 by binding directly to the promoter of the *Rev-erba* clock gene in circadian and PER
74 protein-dependent fashion. In addition, although overexpression or silencing of any one
75 of them influences clock period and amplitude in cells, depletion of paraspeckles
76 themselves has no effect on the circadian oscillator. Mice deficient for two of these
77 proteins show circadian phenotypes, albeit less prominent than *in vitro*. We therefore
78 suggest that all three proteins play redundant roles in circadian transcriptional
79 modulation.

80

81

82 **METHODS**

83

84 **Animal husbandry**

85 Chimeric mice were obtained from *Nono* genetrapped (*Nono^{gt}*) embryonic stem (ES)
86 cells (C57/BL6J genotype) via standard blastocyst injection of the ES clone YHA266
87 into SV129 mice by the University of California, Davis. Individual chimeric mice were
88 back-crossed 4-10 generations against C57/BL6J. The same procedure was chosen to
89 obtain *Pspc1^{gt/gt}* and *Sfpq^{gt/+}* mutant mice, using ES clones RRS358 and BC0256,
90 respectively. Individual chimeric mice were back-crossed 2-4 generations against
91 C57/BL6J. All experiments were performed by comparing wildtype and mutant
92 littermates. Animal housing and experimental procedures are in agreement with
93 veterinary law of the canton of Zurich.

94 **Animal activity measurements**

95 For period measurements of *Nono^{gt}* mice, 24 mice of each genotype were habituated to
96 a controlled 12:12 light-dark (LD) cycle in the presence of running wheels for 2 weeks,
97 and then kept in constant dim red light for an additional two weeks. Data recording and
98 period analysis was performed using the Clocklab software package (Actimetrics).
99 Period measurements of *Pspc1^{gt/gt}* and *Sfpq^{gt/+}* mice were performed identically except
100 that 6 mice of each genotype were used, and measurements were performed twice on
101 each mouse. For skeleton photoperiod measurements, the same mice were given 1
102 hour of normal room light at each LD transition of a normal day, and otherwise kept in
103 constant dim red light. Running wheel activity was measured as in period experiments,

104 but plotted as the sum of activities of all the mice over a 24-hour day using the Clocklab
105 software.

106 **Plasmids**

107 The bioluminescence reporter construct *pBmal1-Luciferase* has been described
108 previously (27). Overexpression of NONO, SFPQ, and PSPC1 (tagged with the myc
109 epitope) were achieved using the plasmids described in (21). Plasmids expressing
110 PER1 and PER2 proteins tagged with the FLAG epitope were a gift of T. Wallach
111 (Kramer lab, Charite Universitätsmedizin, Berlin). To create GAL4 fusion constructs,
112 the same constructs were obtained as EntryTM vectors from NITE (the Japanese
113 Bioresource Information Center), and recombined into a DestinationTM vector
114 (Invitrogen) containing the GAL4 DNA-binding domain (aa 1-93). This vector was made
115 by cloning PCRed recombination sites from pEF-DEST51 (Invitrogen) into pSCT-
116 GALVP80 (gift of W. Schaffner, University of Zurich). The *Neat1* overexpression vector
117 is described in (6). RNAi vectors against NONO have been described previously in (4).
118 Vectors targeting SFPQ and PSPC1 were purchased from Open Biosystems (clone
119 numbers RRM3981 – 98064499 TRCN0000102241 and RMM3981 – 98064691
120 TRCN0000102470, respectively). p4xEbox-*luc* is described in (4). pGAL4-Ebox-*luc* was
121 made by inserting a multimerized 5xGAL4 site (cut from pFR-*luc*, Invitrogen) upstream
122 of the E boxes in p4xEbox-*luc*. pGAL4-CMV-*luc* was made by inserting the same
123 fragment the same distance upstream relative to the transcription start site of the CMV
124 promoter.

125

126

127 Primary cell isolation and culture

128 Primary adult dermal fibroblasts (ADFs) were taken from a 0.5cm piece of mouse tail
129 that was cut into several small pieces by using a razor blade. Digestion occurred in
130 1.8ml DMEM containing 20% FBS, 1% penicillin/streptomycin and 1% amphotericin B
131 supplemented with 0.7 units liberase blendzyme (Roche), at 37°C and 5% CO₂ for eight
132 hours. After centrifugation in 1x PBS the pellet was resuspended in DMEM containing
133 20% FBS, 100U/ml penicillin, 100ug/ml streptomycin and 2.5ug/ml amphotericin B and
134 kept at 37°C and 5% CO₂. The day after, medium was exchanged and remaining tail
135 pieces were removed. Another medium exchange was done three days later. After a
136 week the medium was exchanged for medium without amphotericin B. ADFs were
137 cultured at 37°C and 5% CO₂ in DMEM supplemented with 20% FBS and 1%
138 penicillin/streptomycin.

139 Transient transfections

140 For *p4xE-box* luciferase reporter transfection studies in NIH3T3 cells, lipofectamine LTX
141 with PLUS reagent (Invitrogen) was used according to the manufacturer's instructions,
142 cultivating cells in 24-well plates and transfecting them with a total of 850ng DNA of
143 which 50ng were the promoter luciferase reporter construct. Varying amounts of
144 plasmid were "balanced" by the addition of pcDNA3.1 to a total of 800ng. Cells were
145 harvested after 60 hours by washing once with 1x PBS and extracting luciferase with a
146 luciferase assay kit (Promega) and normalizing against amount of total protein in each
147 extract (measured by Coomassie staining compared to a bovine serum albumin
148 standard curve). Transfections in primary cells were performed identically, except that
149 twice the amount of cells was used for each reaction.

150 Lentiviral infections

151 Measurements were conducted in U2OS cells stably transfected with a circadian
152 *Bmal1-luciferase* reporter, and then infected with Open Biosystems RNAi lentivectors
153 (pGIPZ), as described previously (24).

154 Measurement of circadian bioluminescence in cultured cells

155 After transfection or infection as described above, circadian rhythms in cell populations
156 were synchronized with dexamethasone, and then measured for 3-5 days via real-time
157 luminometry in normal culture medium lacking phenol red but supplemented with 0.2mM
158 luciferin and 25mM HEPES, as described previously (e.g. (27)). Data were analyzed
159 using the Lumicycle Analysis program (Actimetrics).

160 cDNA production and quantitative real-time PCR

161 RNA was extracted as described in (42). 500ng of total RNA was transcribed to cDNA
162 with SuperScript II (Invitrogen) using random hexamer primers according to
163 manufacturer's instructions. For quantitative real-time PCR 20ng of cDNA was used and
164 single transcript levels of genes were detected by Taqman probes used with the
165 Taqman PCR mix protocol (Roche) using the AB7900 thermocycler. Primers used for
166 detection of NOPS transcripts are: *Nono* sense TGC GCT TCG CCT GTC A, antisense
167 GCA GTT CGT TCG ACA GTA CTG, probe FAM-AGT GCA CCC TTA CAG TCC GCA
168 ACC TT-TAMRA; *Pspc1* sense GAA CTA TAC CTG GCC CAC CAA T, antisense ACT
169 GCG CC ATTA TCT GGT ATC A, probe FAM-ATA TTT GCA GCT CCT TCT GGT CCC
170 ATG –TAMRA; *Sfpq* sense TTT GAA AGA TGC AGT GAA GGT GTT; antisense CCT
171 GCT TCA CCA CCT TCT TGA, probe FAM-TCC TAC TGA CAA CGA CTC CTC GCC
172 CA-TAMRA. Primers for detection of circadian genes and GAPDH can be found in (31).

173

174 Protein Extraction and western blotting

175 For *in vitro* immunoprecipitation a 10cm culture dish of HEK 293T cells were co-
176 transfected with each 5ug of NONO-myc, SPFQ-myc or PSCP-myc together with 5ug
177 PER1-FLAG or PER2-FLAG, via polyethyleneimine transfection (JetPEI, Polyplus)
178 following manufacturer's instructions. The cells were harvested 24h later by rinsing with
179 PBS and resuspending in a total of 100ul of lysis buffer as described previously for liver
180 nuclei in (23). Extracts were stored in 500ul aliquots in -80°C until usage. Liver nuclei
181 were prepared by sucrose cushion centrifugation as described in (23), then extracted
182 exactly as for cells. Western blotting was performed using standard procedures
183 (*Current Protocols in Molecular Biology*, Wiley). Equal loading and size detection using
184 protein ladder was verified by Ponceau-S staining of membranes prior to probing.

185 Immunohistochemistry

186 Immunohistochemistry was performed according to the protocols described at
187 <http://www.pharma.uzh.ch/research/neuromorphology/researchareas/neuromorphology/Protocol>
188 [s/protocol_immuno.pdf](#). Substrates were either brains collected in isopentane at -20°C
189 and cryostatically sliced, or cells grown on glass coverslips, rinsed with PBS, and fixed
190 5' at room temperature in PBS/4%paraformaldehyde.

191 Antibodies

192 Polyclonal antibodies against NONO, SFPQ, PSCP1, and PER2 were produced from
193 rabbits by Charles River Laboratories using bacterially-overexpressed proteins.
194 Antibody from each serum was immunopurified over a column whose resin consisted of
195 the relevant antigen covalently coupled to Affygel 10 (BioRad). Anti-PSCP1 is

196 described in (12). For detection in Co-IP experiments primary anti-MYC antibody
197 (Roche, Cat N°11667149001) was diluted at 1:2000, primary anti-FLAG antibody
198 (Sigma, F3167) 1:2000, primary anti-NONO antibody at 1:2000, primary anti-PSPC1 at
199 1:1000, primary anti-SPFQ antibody at 1:2000, primary anti-PER2 antibody at 1:1000.
200 The probing of the secondary antibody was done at 1:10'000 for IRDye 680 Goat Anti-
201 Mouse IgG (Licor, 926-32220) and 1:10'000 for IRDye 800 Goat Anti-Rabbit IgG (Licor,
202 926-33210). For immunoprecipitations, primary anti-cMYC antibody was diluted at
203 1:500, primary anti-FLAG antibody at 1:500, primary anti-NONO antibody for IP at
204 1:100, primary anti-SFPQ antibody for IP at 1:100, primary anti-PSCP1 antibody for IP
205 at 1:100 and primary anti-PER2 antibody for IP at 1:100.

206 **Immunoprecipitation**

207 Immunoprecipitation was performed using standard procedures with the below
208 mentioned adjustments (*Current Protocols in Molecular Biology*, Wiley). Extracts were
209 pre-cleared by incubation the crude extracts with protein-A beads (Calbiochem, Cat. N°
210 IP06) and 0.1%BSA for 1h at 4°C. 500ug of pre-cleared extract were bound for 2h to
211 antibody with Co-IP buffer. The Antibody-protein complex was then incubated for 1h
212 with protein-A beads. The beads were washed gently with Co-IP buffer (without
213 protease inhibitor mix) and denatured for 15min at 65°C with 2xSDS sample buffer
214 containing beta-mercaptoethanol. Equal amounts of IP reactions were loaded on a 7%
215 (overexpression in cells IP) or 9% (liver nuclei extracts IP) SDS PAGE gel together with
216 1/10 of the IP amounts of pre-cleared extract as input. The protein gel and blotting was
217 performed as described in the *Western blotting and immunohistochemistry* section
218 above.

219

220 Chromatin Immunoprecipitation

221 Chromatin from mouse liver and tissue culture cells was obtained as described
222 previously (34). Equal amounts of precleared chromatin were incubated overnight at 4°C
223 with 1 ul of anti-NONO antibody or anti-PER2 antibody. The capture of the DNA:protein
224 complexes, the washing conditions and the purification of the DNA fragments prior to
225 qPCR as well the control antibodies have been described previously (36). The region-
226 specific primer/probe pairs are listed in Supplementary Methods.

227 Paraspeckle Quantification

228 For paraspeckle detection, after immunodetection of PSPC1 as described above, cells
229 were analyzed with a LSM710 Zeiss confocal microscope. Pictures taken were with 40x
230 (NA1.3), and the pinhole was kept at 1AU or 0.8 to 0.9um. Nuclei were manually
231 detected using ImageJ software routines (<http://rsbweb.nih.gov/ij/index.html>). Speckles
232 were determined by subtracting background nucleoplasmic PSPC1 protein staining, and
233 thereafter counting remaining pixel clusters in nuclei. The total amount of paraspeckles
234 per cell was estimated by counting all pixels brighter than 140 (arbitrary units) with spot
235 sizes between 0.25-10 squaremicrometers. Nuclei smaller than 200 pixels or 100
236 squaremicrometers as well as dividing cells were excluded. The averaged number of
237 speckles was normalized to mean area and compared to the control transfected cells
238 (hairpin NEAT-S).

239

240

241

242 **RESULTS**243 *NONO-deficient mice show significant changes in circadian period*

244 To better understand the function of NONO in the circadian clock and in mammalian
245 physiology, we obtained NONO-deficient mice from ES cells bearing a genetrap in the
246 intron preceding the *Nono* translational start site (Fig S1A, B). In wildtype mice, NONO
247 is expressed in most tissues including the suprachiasmatic nuclei in the brain. *Nono*^{gt}
248 mice showed no expression of *Nono* mRNA or protein in all tissues examined (Fig 1A,
249 B). These mice showed a twenty-minute reduction in circadian behavioral period when
250 placed into constant dark conditions (Fig 1C). This reduction in period length was highly
251 significant, but it was nevertheless far less dramatic compared to a *Drosophila*
252 hypomorphic *nonA* strain that we observed previously to become arrhythmic (4).
253 Hence, we suspected that in mammals the lack of *Nono* may be compensated by other
254 factors.

255

256 *NONO binds to the circadian promoter of the Rev-erba gene*

257 Next, we verified the relevance of NONO *in vivo* by looking for its presence at the
258 promoters of clock genes. Since we showed previously that NONO interacted with PER
259 proteins (4), we guessed that it ought to be found at PER-regulated clock genes.
260 Chromatin immunoprecipitation experiments confirmed that this was indeed the case:
261 NONO interacted with the promoter of the *Rev-Erb α* gene in circadian fashion, sharing
262 the same kinetics as the PER1 protein (Fig 1D, top). This interaction was considerably
263 reduced but surprisingly not absent in *Nono*^{gt} mice, which completely lack NONO
264 transcript and protein (Fig 1D, bottom). Equivalent results were seen for the *Dbp*

265 promoter (Fig 2A), and no binding was observed at the promoter of the antiphase
266 *Bmal1* gene (Fig 2B). Based upon the residual binding observed at the *Rev-Erb α* and
267 *Dbp* promoters, we considered the possibility that NONO might be redundant with
268 homologous DBHS-family factors with which our antibody might weakly cross-react.
269 Conserved domain analysis (with the Conserved Domain Architecture Tool CDART,
270 (14)) showed that the other two known DBHS proteins, PSPC1 and SFPQ, shared both
271 high homology with NONO and a similar domain architecture (Fig S1C, D). In addition,
272 SFPQ was recently shown to play a role in circadian transcriptional repression (10).
273 Therefore, we speculated that all three proteins might have similar functions in the
274 circadian oscillator.

275

276 Overexpression or silencing of DBHS proteins interferes with circadian function

277 To test this idea, we transfected vectors expressing each of the three proteins into
278 cultured cells together with a luciferase reporter under control of the circadian *Rev-Erb α*
279 gene promoter. After synchronizing circadian clocks in these transfected cells with
280 dexamethasone (2), we monitored reporter bioluminescence in real time.

281 Overexpression of any of the three proteins in human U2OS fibroblasts perturbed
282 circadian rhythmicity (Fig 3A).

283

284 We next undertook loss-of-function experiments based upon RNA interference (RNAi),
285 in which U2OS human osteosarcoma cells containing an integrated *Bmal1-luciferase*
286 reporter were infected with lentiviruses expressing shRNAs targeting *Pspc1* or *Sfpq*.
287 RNAi hairpins against SFPQ dampened circadian oscillations dramatically (Fig 3B)

288 similar to what was observed previously for NONO (4), but those against PSPC1
289 lengthened it and somewhat dampened amplitude (Fig 3C). Measurement of *Sfpq* and
290 *Pspc1* RNA levels in these cells showed that these hairpins reduced expression of *Sfpq*
291 7-fold, and *Pspc1* 2.5-fold (Fig 3D,E).

292

293 To insure that the effects that we observed were not cell type-specific, identical
294 experiments were conducted using NIH-3T3 mouse fibroblasts. Again, overexpression
295 of any of the three proteins perturbed circadian rhythmicity (Fig 4A). Suppression of
296 circadian rhythmicity was also seen in NIH3T3 cells transiently transfected with the
297 circadian *Rev-Erba* promoter reporter together with RNAi hairpins targeting *Pspc1* or
298 *Sfpq* (Fig 4B,C). In this case, immunofluorescence experiments showed that these
299 hairpins reduced expression of SFPQ 2-fold, and PSPC1 10-fold (Fig 4D).

300

301 Depletion of paraspeckles does not perturb overall circadian clock function

302 Since the three NONO-related proteins are also the three known members of nuclear
303 paraspeckles, we speculated that the paraspeckle itself might serve a circadian role.
304 This subnuclear domain requires the nuclear noncoding RNA *Neat1*, probably as a
305 scaffold, and depletion of *Neat1* has been shown to eliminate paraspeckles themselves
306 (5, 6). By transiently transfecting shRNAs complementary to *Neat1* into U2OS cells, we
307 were able to reduce *Neat1* levels (Fig 4E) and thereby deplete paraspeckles, measured
308 by counting the number of punctate PSPC1 foci (Fig 5A,B). However, cotransfection of
309 the circadian *Bmal1-luc* reporter showed that the circadian clock retained normal period
310 length in these paraspeckle-depleted cells (Fig 5C), making it unlikely that paraspeckles

311 *per se* play a significant role in the circadian oscillator. Therefore, it is probable that
312 nucleoplasmic, non-paraspeckle-associated pools of NONO, SFPQ, and PSPC1
313 proteins were responsible for the circadian effects that we have documented.

314

315 *DBHS proteins bind to clock promoter DNA and repress clock gene transcription*

316 Since NONO can bind to circadian clock gene promoters *in vivo* (Fig 1, 2), it was logical
317 to imagine that the other DBHS factors might do the same. Indeed, similarly to NONO,
318 SFPQ and PSPC1 could also be immunoprecipitated at the *Rev-Erb α* promoter in a
319 circadian fashion in liver nuclear extracts (Fig 6A).

320 All three proteins are present at clock gene promoters with the same temporal profile as
321 PER proteins, suggesting a co-repressor function. Moreover, both NONO and SFPQ
322 were identified as PER-interacting proteins, and show interactions with PER1 and PER2
323 in various contexts ((4), (10), and Fig 6B-D). However, actual functions of these
324 proteins are less clear. For example, we have shown previously that NONO can
325 *antagonize* PER-mediated transcriptional repression when transfected into immortalized
326 cells (4). In fact, the same is seen with SFPQ and to a lesser extent with PSPC1 (Fig
327 S2A-B), whereas overexpression of NEAT does not influence BMAL1/CLOCK mediated
328 transcriptional activation (Fig S2C). In other reports, NONO and SFPQ have been
329 reported by different investigators as either transcriptional coactivators or corepressors
330 (16, 26), and SFPQ has been shown to act as a transcriptional repressor in the
331 circadian clock (10).

332

333 Within the circadian clock, we favor a repressive role of these factors because of the
334 following experiments. First, when we transfected primary mouse fibroblasts with an E-
335 box-driven luciferase reporter, together with the transcriptional activators CLOCK and
336 BMAL1 and either NONO, PSPC1, or SFPQ, both NONO and SFPQ repressed
337 CLOCK-BMAL-mediated transcription from the reporter, and to a lesser extent PSPC1,
338 which was initially activating and then repressing at higher concentrations (Fig 7A).
339 When equivalent transfections were performed using fibroblasts from *Per1^{brdm/brdm} /*
340 *Per2^{brdm/brdm}* mice that lack functional PER proteins and circadian clocks (44),
341 repression was no longer observed, but instead weak activation (Fig 7A). Similarly, in
342 *Per1^{brdm/brdm} / Per2^{brdm/brdm}* mice no circadian immunoprecipitation of NONO was
343 observed at the *Rev-Erb α* promoter (Fig 7B). Therefore, PER recruits NONO and
344 presumably the other family members too.

345

346 Secondly, to confirm that DBHS factors are repressors at circadian promoters, we
347 designed a hybrid GAL4 DNA-binding domain-E-box-luciferase reporter and fusions of
348 NONO, PSPC1, and SFPQ with the GAL4 DNA-binding domain in order to enable their
349 direct recruitment to DNA independently of PER proteins. When the GAL4DBD-Ebox-
350 luciferase construct was transfected into primary mouse fibroblasts together with the
351 GAL4-VP16 transcriptional activator, strong activation was observed, demonstrating the
352 functionality of the construct. When GAL4-NONO, -PSPC1, and -SFPQ were
353 cotransfected into primary mouse fibroblasts together with the GAL4 DNA-binding
354 domain-E-box-luciferase reporter, all three proteins strongly repressed CLOCK-BMAL-

355 mediated transcription (Fig 7C), though they had no statistically significant effect when
356 similarly recruited to the constitutively active CMV promoter (Fig 7D).

357

358 Importance of DBHS proteins to circadian behavior

359 Finally, in order to verify the relevance of these factors to the circadian clock *in vivo*, we
360 obtained mice with genetrap-based inactivations of *Pspc1* and *Sfpq*, to match the
361 *Nono^{gt}* mouse described earlier in this paper. Homozygous *Pspc1*-genetrapped mice
362 showed fivefold reduction in *Pspc1* transcript levels in multiple tissues (Fig 8A), and no
363 detectable levels of PSPC1 protein in liver nuclear extracts (Fig 8C). Although the *Sfpq*
364 genetrap was homozygous lethal, heterozygous mice showed up to twofold reduction in
365 both RNA and protein (Fig 8B, D). When tested for circadian wheel-running behavior,
366 these *Sfpq^{gt/+}* mice also showed a trend toward shortening of period similar to that of
367 *Nono^{gt}* in some animals (Fig 9A, B), as well as altered entrainment in a minimal-light
368 "skeleton" photoperiod in all animals (Fig S3A-C). *Pspc1^{gt/gt}* mice showed no
369 abnormalities (Fig 9A-B, Fig S3).

370

371 Consistent with the proposed repressive role of these factors, at the gene expression
372 level, *Rev-Erb α* RNA showed modestly increased expression in liver extracts from all
373 three knockouts at the time (CT8-12) that coincides with binding of NONO and PER2
374 (Fig 9C). Interestingly, its timing coincides with the peak of *Rev-Erb α* expression levels
375 and the beginning of their decline, but not with maximum repression. Hence, it is
376 possible that these factors are associated with the establishment of repression but not
377 its maintenance. Similar but smaller gene expression effects were seen upon *Per2*

378 transcript levels, but the expression of other clock genes remained mostly unchanged
379 (Fig S4).

380

381 **DISCUSSION**

382 Because of their homologies, shared functions, and abilities to interact with one another,
383 the three factors NONO, PSpC1, and SFPQ have recently been classified by multiple
384 authors as a family of proteins: the NOPS family (for NO_{no} and PS_{pc}1, (40)) or DBHS
385 family (for Drosophila Behavior Human Splicing, (3)). Our data and that of others point
386 to another important role of these proteins within the circadian oscillator. We initially
387 isolated NONO as a PER-interacting protein (4), and Duong *et al.* recently isolated
388 SFPQ in the same way (10). Here we present data that all three DBHS proteins likely
389 play overlapping roles within the circadian clock.

390

391 **Nuclear paraspeckles and the circadian clock**

392 All three DBHS proteins are part of nuclear paraspeckles (13), subnuclear bodies
393 probably involved in splicing and RNA storage. Nevertheless, our data suggests that
394 the paraspeckle *per se* is not important for circadian function: depletion of these
395 nuclear bodies by targeting the structural ncRNA *Neat1* (6) has no effect upon the
396 circadian clock, nor does transfection of this ncRNA into cells alter E-box-mediated
397 transcription. Instead, our results imply that DBHS proteins likely exists in at least two
398 nuclear pools. One of these pools is present in paraspeckles, and appears to play no
399 role so far in the circadian clock, though it may be important for nuclear retention of

400 edited RNAs as reported by others (5, 30, 43). A second pool is nucleoplasmic, and
401 could be in part responsible for the transcriptional roles reported for DBHS proteins.

402

403 *DBHS proteins as transcription factors*

404 Besides their roles in nuclear paraspeckles, the previously reported functions of DBHS
405 proteins have ranged widely. They have been implicated in splicing (17, 29) and axonal
406 transport of RNA (18). They are players in the regulation of pre-mRNA processing and
407 transcription termination (19), and in the DNA damage response (32, 35). NONO has
408 also been characterized as a nonclassical carbonic anhydrase (20). In addition,
409 however, all have been implicated in transcription. In some cases they have been
410 implicated as activators (1, 16, 21), and in other cases as repressors (8, 26, 39), even
411 for the same gene (37). Interestingly, a mechanism has been proposed in both cases:
412 whereas NONO and SFPQ can interact directly with the RNA polymerase II CTD in a
413 way that might explain transcriptional activation (11), SFPQ has been proposed to
414 recruit the mSIN3A histone deacetylase to promote repression (10, 26). For the
415 circadian clock we originally identified NONO as an antagonist of PER-mediated
416 repression (4), and others demonstrated PER-mediated repression by SFPQ (10).
417 Apparently the roles of DBHS protein depend on the cellular context. In this paper
418 alone, we show that transient transfection of SFPQ into U2OS cells can activate
419 transcription from E box reporters, but its transfection into mouse primary fibroblasts
420 represses it. To try to resolve the role of these factors within the circadian clock, we
421 therefore created GAL4 fusion proteins to unambiguously recruit these factors to
422 promoters. All three had no effect on the CMV promoter, but strongly repressed

423 transcription mediated by the circadian transcription factors CLOCK and BMAL1 at a
424 circadian E-box promoter. With this experiment, we show a) that the effects of these
425 factors upon transcription are context-specific, and b) that they are likely to be
426 repressors in the circadian context.

427

428 *Overlapping functions of DBHS proteins in the circadian clock*

429 One possible explanation for our results and those of others suggesting activation or
430 repression is overlapping function of related genes: if two repressors have different
431 repressive potentials, for example, then titrating increasing exogenous amounts of the
432 weaker results in an increase in transcription as the endogenous stronger one is
433 displaced. In this paper, we present considerable evidence for overlapping functions of
434 the three DBHS proteins within the circadian clock. Depletion or overexpression of all
435 three unambiguously affects circadian function in cells and in cellular transcription
436 assays, but depletion of any one in mice results in only small circadian phenotypes, and
437 generation of double DBHS mutants -- or even complete knockouts of *Sfpq* -- is
438 complicated by embryonic lethality. Moreover, we and others have shown by chromatin
439 immunoprecipitation that all three DBHS proteins can bind directly to clock promoters or
440 clock-controlled promoters in circadian fashion *in vivo* and in cells (10, 15, 26). At least
441 for circadian function, it is likely that this binding requires PER proteins. SFPQ and
442 NONO were identified as PER-interacting proteins, and clearly immunoprecipitate with
443 them as shown here and elsewhere (4, 10). Moreover, we show here that in PER-
444 deficient mice, binding of NONO to circadian promoters is no longer observed. Thus,
445 we think it is likely that PER proteins recruit DBHS proteins to clock-controlled genes to

446 control and orchestrate PER-mediated transcriptional repression. The degree of this
447 repression could be precisely controlled by the mix of these factors recruited.
448 Functional redundancy of this family of proteins is also highlighted by the recently
449 published crystal structure of a NONO-PSPC1 complex (28). Not only do these proteins
450 probably form obligate heterodimers, which would suggest a role for multiple DBHS
451 family members within the circadian clock, but their structure also allows for possible
452 higher-order oligomers, which might provide an ideal platform for the recruitment of
453 other factors that have been found associated with these factors in various contexts.

454

455 *DBHS proteins as orchestrators of circadian physiology*

456 Although we have shown clear roles of DBHS proteins in a cellular context, the
457 circadian behavioral phenotypes of DBHS protein-deficient mice were relatively minor.
458 As discussed above, functional redundancy could account for this lack of phenotype. In
459 addition, however, the uniquely coupling of SCN cells into a network renders them more
460 resilient to the effects of mutation (22). Therefore, it is also possible that more severe
461 circadian effects of DBHS proteins might occur in peripheral tissues.

462

463 Indeed, it is likely that considerable further circadian physiology directed by DBHS
464 proteins remains to be elucidated. Mice deficient in these factors show a spectrum of
465 unique phenotypes, ranging from embryonic lethality (*Sfpq*) to neurological phenotypes
466 (*Nono*). *Pspc1* protein is strongly regulated in circadian fashion although the other two
467 factors are not (Fig 8 and data not shown). Moreover, the E-box is a standard motif for
468 orchestrating clock-controlled physiology (34), and directs circadian transcription at

469 thousands of promoters (41). DBHS factor binding has been observed at multiple clock-
470 regulated promoters containing this motif, including prolactin (15), progesterone (9),
471 *Rev-Erb α* (Fig 1B), and androgen receptor (8). Through their interaction with PER
472 proteins, we show here that DBHS factors play an important role directly in the circadian
473 oscillator. Binding to clock gene promoters and modulating transcriptional repression,
474 they regulate a portion of the transcriptional feedback which is the hallmark of metazoan
475 circadian clocks.

476

477 **ACKNOWLEDGEMENTS**

478 We thank Robert Dallmann for a critical reading of this manuscript, and W. Schaffner
479 (U. Zurich) for donation of Gal4 vectors. This work was supported by the Swiss
480 National Science Foundation and the University of Zurich Fonds der Akademischen
481 Nachwuchses. Further support to S.A.B. was provided by the Neurosciences Center
482 Zurich and the Molecular Life Sciences programs.

483

484

485 **REFERENCES**

486

- 487 1. **Amelio, A. L., L. J. Miraglia, J. J. Conkright, B. A. Mercer, S. Batalov, V.**
488 **Cavett, A. P. Orth, J. Busby, J. B. Hogenesch, and M. D. Conkright.** 2007. A
489 coactivator trap identifies NONO (p54nrb) as a component of the cAMP-signaling
490 pathway. *Proc Natl Acad Sci U S A* **104**:20314-9.
- 491 2. **Balsalobre, A., S. A. Brown, L. Marcacci, F. Tronche, C. Kellendonk, H. M.**
492 **Reichardt, G. Schutz, and U. Schibler.** 2000. Resetting of circadian time in
493 peripheral tissues by glucocorticoid signaling. *Science* **289**:2344-7.
- 494 3. **Bond, C. S., and A. H. Fox.** 2009. Paraspeckles: nuclear bodies built on long
495 noncoding RNA. *J Cell Biol* **186**:637-44.

- 496 4. **Brown, S. A., J. Ripperger, S. Kadener, F. Fleury-Olela, F. Vilbois, M.**
497 **Rosbash, and U. Schibler.** 2005. PERIOD1-associated proteins modulate the
498 negative limb of the mammalian circadian oscillator. *Science* **308**:693-6.
- 499 5. **Chen, L. L., and G. G. Carmichael.** 2009. Altered nuclear retention of mRNAs
500 containing inverted repeats in human embryonic stem cells: functional role of a
501 nuclear noncoding RNA. *Mol Cell* **35**:467-78.
- 502 6. **Clemson, C. M., J. N. Hutchinson, S. A. Sara, A. W. Ensminger, A. H. Fox, A.**
503 **Chess, and J. B. Lawrence.** 2009. An architectural role for a nuclear noncoding
504 RNA: NEAT1 RNA is essential for the structure of paraspeckles. *Mol Cell* **33**:717-
505 26.
- 506 7. **Dibner, C., U. Schibler, and U. Albrecht.** 2010. The mammalian circadian
507 timing system: organization and coordination of central and peripheral clocks.
508 *Annu Rev Physiol* **72**:517-49.
- 509 8. **Dong, X., J. Sweet, J. R. Challis, T. Brown, and S. J. Lye.** 2007.
510 Transcriptional activity of androgen receptor is modulated by two RNA splicing
511 factors, PSF and p54nrb. *Mol Cell Biol* **27**:4863-75.
- 512 9. **Dong, X., C. Yu, O. Shynlova, J. R. Challis, P. S. Rennie, and S. J. Lye.** 2009.
513 p54nrb is a transcriptional corepressor of the progesterone receptor that
514 modulates transcription of the labor-associated gene, connexin 43 (Gja1). *Mol*
515 *Endocrinol* **23**:1147-60.
- 516 10. **Duong, H. A., M. S. Robles, D. Knutti, and C. J. Weitz.** 2011. A molecular
517 mechanism for circadian clock negative feedback. *Science* **332**:1436-9.
- 518 11. **Emili, A., M. Shales, S. McCracken, W. Xie, P. W. Tucker, R. Kobayashi, B. J.**
519 **Blencowe, and C. J. Ingles.** 2002. Splicing and transcription-associated
520 proteins PSF and p54nrb/nonO bind to the RNA polymerase II CTD. *Rna* **8**:1102-
521 11.
- 522 12. **Fox, A. H., C. S. Bond, and A. I. Lamond.** 2005. P54nrb forms a heterodimer
523 with PSP1 that localizes to paraspeckles in an RNA-dependent manner. *Mol Biol*
524 *Cell* **16**:5304-15.
- 525 13. **Fox, A. H., and A. I. Lamond.** 2010. Paraspeckles. *Cold Spring Harb Perspect*
526 *Biol* **2**:a000687.
- 527 14. **Geer, L. Y., M. Domrachev, D. J. Lipman, and S. H. Bryant.** 2002. CDART:
528 protein homology by domain architecture. *Genome Res* **12**:1619-23.
- 529 15. **Guillaumond, F., B. Boyer, D. Becquet, S. Guillen, L. Kuhn, J. Garin, M.**
530 **Belghazi, O. Bosler, J. L. Franc, and A. M. Francois-Bellan.** 2011. Chromatin
531 remodeling as a mechanism for circadian prolactin transcription: rhythmic NONO
532 and SFPQ recruitment to HLTF. *Faseb J* **25**:2740-56.
- 533 16. **Ishitani, K., T. Yoshida, H. Kitagawa, H. Ohta, S. Nozawa, and S. Kato.** 2003.
534 p54nrb acts as a transcriptional coactivator for activation function 1 of the human
535 androgen receptor. *Biochem Biophys Res Commun* **306**:660-5.

- 536 17. **Kameoka, S., P. Duque, and M. M. Konarska.** 2004. p54(nrb) associates with
537 the 5' splice site within large transcription/splicing complexes. *Embo J* **23**:1782-
538 91.
- 539 18. **Kanai, Y., N. Dohmae, and N. Hirokawa.** 2004. Kinesin transports RNA:
540 isolation and characterization of an RNA-transporting granule. *Neuron* **43**:513-
541 25.
- 542 19. **Kaneko, S., O. Rozenblatt-Rosen, M. Meyerson, and J. L. Manley.** 2007. The
543 multifunctional protein p54nrb/PSF recruits the exonuclease XRN2 to facilitate
544 pre-mRNA 3' processing and transcription termination. *Genes Dev* **21**:1779-89.
- 545 20. **Karhumaa, P., S. Parkkila, A. Waheed, A. K. Parkkila, K. Kaunisto, P. W.**
546 **Tucker, C. J. Huang, W. S. Sly, and H. Rajaniemi.** 2000. Nuclear
547 NonO/p54(nrb) protein is a nonclassical carbonic anhydrase. *J Biol Chem*
548 **275**:16044-9.
- 549 21. **Kuwahara, S., A. Ikei, Y. Taguchi, Y. Tabuchi, N. Fujimoto, M. Obinata, S.**
550 **Uesugi, and Y. Kurihara.** 2006. PSPC1, NONO, and SFPQ are expressed in
551 mouse Sertoli cells and may function as coregulators of androgen receptor-
552 mediated transcription. *Biol Reprod* **75**:352-9.
- 553 22. **Liu, A. C., D. K. Welsh, C. H. Ko, H. G. Tran, E. E. Zhang, A. A. Priest, E. D.**
554 **Buhr, O. Singer, K. Meeker, I. M. Verma, F. J. Doyle, 3rd, J. S. Takahashi,**
555 **and S. A. Kay.** 2007. Intercellular coupling confers robustness against mutations
556 in the SCN circadian clock network. *Cell* **129**:605-16.
- 557 23. **Lopez-Molina, L., F. Conquet, M. Dubois-Dauphin, and U. Schibler.** 1997.
558 The DBP gene is expressed according to a circadian rhythm in the
559 suprachiasmatic nucleus and influences circadian behavior. *Embo J* **16**:6762-71.
- 560 24. **Maier, B., S. Wendt, J. T. Vanselow, T. Wallach, S. Reischl, S. Oehmke, A.**
561 **Schlosser, and A. Kramer.** 2009. A large-scale functional RNAi screen reveals
562 a role for CK2 in the mammalian circadian clock. *Genes Dev* **23**:708-18.
- 563 25. **Masri, S., and P. Sassone-Corsi.** 2010. Plasticity and specificity of the circadian
564 epigenome. *Nat Neurosci* **13**:1324-9.
- 565 26. **Mathur, M., P. W. Tucker, and H. H. Samuels.** 2001. PSF is a novel
566 corepressor that mediates its effect through Sin3A and the DNA binding domain
567 of nuclear hormone receptors. *Mol Cell Biol* **21**:2298-311.
- 568 27. **Nagoshi, E., C. Saini, C. Bauer, T. Laroche, F. Naef, and U. Schibler.** 2004.
569 Circadian gene expression in individual fibroblasts: cell-autonomous and self-
570 sustained oscillators pass time to daughter cells. *Cell* **119**:693-705.
- 571 28. **Passon, D. M., M. Lee, O. Rackham, W. A. Stanley, A. Sadowska, A.**
572 **Filipovska, A. H. Fox, and C. S. Bond.** 2012. Structure of the heterodimer of
573 human NONO and paraspeckle protein component 1 and analysis of its role in
574 subnuclear body formation. *Proc Natl Acad Sci U S A* **109**:4846-50.

- 575 29. **Patton, J. G., E. B. Porro, J. Galceran, P. Tempst, and B. Nadal-Ginard.**
576 1993. Cloning and characterization of PSF, a novel pre-mRNA splicing factor.
577 *Genes Dev* **7**:393-406.
- 578 30. **Prasanth, K. V., S. G. Prasanth, Z. Xuan, S. Hearn, S. M. Freier, C. F.**
579 **Bennett, M. Q. Zhang, and D. L. Spector.** 2005. Regulating gene expression
580 through RNA nuclear retention. *Cell* **123**:249-63.
- 581 31. **Preitner, N., F. Damiola, L. Lopez-Molina, J. Zakany, D. Duboule, U.**
582 **Albrecht, and U. Schibler.** 2002. The orphan nuclear receptor REV-ERB α
583 controls circadian transcription within the positive limb of the mammalian
584 circadian oscillator. *Cell* **110**:251-60.
- 585 32. **Proteau, A., S. Blier, A. L. Albert, S. B. Lavoie, A. M. Traish, and M. Vincent.**
586 2005. The multifunctional nuclear protein p54^{nrb} is multiphosphorylated in
587 mitosis and interacts with the mitotic regulator Pin1. *J Mol Biol* **346**:1163-72.
- 588 33. **Reischl, S., and A. Kramer.** 2011. Kinases and phosphatases in the mammalian
589 circadian clock. *FEBS Lett.*
- 590 34. **Ripperger, J. A., and U. Schibler.** 2006. Rhythmic CLOCK-BMAL1 binding to
591 multiple E-box motifs drives circadian Dbp transcription and chromatin
592 transitions. *Nat Genet* **38**:369-74.
- 593 35. **Salton, M., Y. Lerenthal, S. Y. Wang, D. J. Chen, and Y. Shiloh.** 2010.
594 Involvement of matrin 3 and SFPQ/NONO in the DNA damage response. *Cell*
595 *Cycle* **9**.
- 596 36. **Schmutz, I., J. A. Ripperger, S. Baeriswyl-Aebischer, and U. Albrecht.** 2010.
597 The mammalian clock component PERIOD2 coordinates circadian output by
598 interaction with nuclear receptors. *Genes Dev* **24**:345-57.
- 599 37. **Sewer, M. B., V. Q. Nguyen, C. J. Huang, P. W. Tucker, N. Kagawa, and M. R.**
600 **Waterman.** 2002. Transcriptional activation of human CYP17 in H295R
601 adrenocortical cells depends on complex formation among p54^(nrb)/NonO,
602 protein-associated splicing factor, and SF-1, a complex that also participates in
603 repression of transcription. *Endocrinology* **143**:1280-90.
- 604 38. **Shav-Tal, Y., and D. Zipori.** 2002. PSF and p54^(nrb)/NonO--multi-functional
605 nuclear proteins. *FEBS Lett* **531**:109-14.
- 606 39. **Song, K. S., K. Kim, K. C. Chung, J. H. Seol, and J. H. Yoon.** 2008. Interaction
607 of SOCS3 with NonO attenuates IL-1 β -dependent MUC8 gene expression.
608 *Biochem Biophys Res Commun* **377**:946-51.
- 609 40. **Staub, E., P. Fizev, A. Rosenthal, and B. Hinzmann.** 2004. Insights into the
610 evolution of the nucleolus by an analysis of its protein domain repertoire.
611 *Bioessays* **26**:567-81.
- 612 41. **Ueda, H. R., S. Hayashi, W. Chen, M. Sano, M. Machida, Y. Shigeyoshi, M.**
613 **Iino, and S. Hashimoto.** 2005. System-level identification of transcriptional
614 circuits underlying mammalian circadian clocks. *Nat Genet* **37**:187-92.

- 615 42. **Xie, W. Q., and L. I. Rothblum.** 1991. Rapid, small-scale RNA isolation from
616 tissue culture cells. *Biotechniques* **11**:324, 326-7.
- 617 43. **Zhang, Z., and G. G. Carmichael.** 2001. The fate of dsRNA in the nucleus: a
618 p54(nrb)-containing complex mediates the nuclear retention of promiscuously A-
619 to-I edited RNAs. *Cell* **106**:465-75.
- 620 44. **Zheng, B., U. Albrecht, K. Kaasik, M. Sage, W. Lu, S. Vaishnav, Q. Li, Z. S.**
621 **Sun, G. Eichele, A. Bradley, and C. C. Lee.** 2001. Nonredundant roles of the
622 mPer1 and mPer2 genes in the mammalian circadian clock. *Cell* **105**:683-94.
- 623
- 624
- 625

626 **FIGURE LEGENDS**

627

628 **Fig 1. (A)** NONO RNA expression measured by qPCR in various tissues taken from
629 wildtype (black) and *Nono*^{gt} animals (grey, not detectable). Inset, NONO protein
630 measured in liver nuclear extract from the same animals, as well as in unrelated C57-
631 Bl6J mice (Bl6/J). **(B)** NONO protein expression in brain coronal sections from wildtype
632 and *Nono*^{gt} animals, visualized by immunohistochemistry using a polyclonal anti-NONO
633 antibody. Arrows from left to right show principal areas of NONO expression in wildtype
634 mouse brain: suprachiasmatic nuclei, hippocampus, and neocortex. **(C)** Wheel-running
635 activity of wildtype and *Nono*^{gt} mice in 12:12 light-dark cycles (LD, arrow) and in
636 constant darkness (DD, arrow). Darkness is indicated by grey shading. N=23. **(D)**
637 Chromatin immunoprecipitation of NONO (black bars) and PER1 (striped bars) at the
638 *Rev-Erb α* promoter in liver nuclei harvested at different circadian times (CT) of day in
639 constant darkness. CT0 = beginning of subjective day. Top panel, wildtype mice.
640 Bottom panel, *Nono*^{gt} mice. (N=3 experiments, shown +/- standard deviation, **
641 corresponds to a p-Value of ≤ 0.01).

642

643 **Fig 2. (A)** Chromatin immunoprecipitation of NONO (black bars) and PER2 (striped
644 bars) at the *Dbp* promoter in liver nuclei harvested from wildtype mice (top panel) and
645 *Nono*^{gt} mice (bottom panel) at different times of day in constant darkness. N=3
646 experiments, +/-SD. **(B)** Identical experiments for the *Bmal1* promoter.

647

648 **Fig 3. (A)** Bioluminescence from U2OS cells transiently transfected with the *Rev-Erb α -*
649 *luciferase* circadian reporter and constructs expressing either NONO, SFPQ, or PSPC1.
650 Data shown is detrended and expressed in arbitrary units relative to mean expression.
651 Solid black line, cells transfected with the empty vector. Dashed line, cells
652 overexpressing NONO. Dashed and dotted line, cells overexpressing PSPC1. Dotted
653 line, cells overexpressing SFPQ. **(B)** Bioluminescence from U2OS cells containing an
654 integrated *Bmal1-luciferase* circadian reporter, infected with viruses expressing two
655 different RNAi hairpins targeting the *Sfpq* gene, and then clock-synchronized with
656 dexamethasone. Data shown is detrended and expressed in arbitrary units relative to
657 mean expression. Solid black line, scrambled-sequence shRNA. Grey line, shRNA3.
658 Dashed line, shRNA5. **(C)** Similar experiment with RNAi constructs targeting *Pspc1*.
659 Dashed line, shRNA3. Grey line, shRNA4. **(D,E)** Transcript levels of *Pspc1* (D) and *Sfpq*
660 (E) in U2OS cells infected with lentiviruses expressing the indicated RNAi targeting
661 vectors used in Figure 2. N=3 +/- SE.

662

663 **Fig 4. (A)** Bioluminescence from 3T3 cells transiently transfected with the *Rev-Erb α -*
664 *luciferase* circadian reporter and constructs expressing either NONO, SFPQ, or PSPC1.
665 Data shown is detrended and expressed in arbitrary units relative to mean expression.
666 Solid black line, wildtype cells. Dashed line, cells overexpressing NONO. Dashed and
667 dotted line, cells overexpressing PSPC1. Dotted line, cells overexpressing SFPQ. **(B,C)**
668 Bioluminescence from 3T3 cells transiently transfected with the *Rev-Erb α -luciferase*
669 circadian reporter and RNAi constructs targeting either *Pspc1* (B) or *Sfpq* (C). After
670 synchronization with dexamethasone, cultures were measured 3 days. Data are shown

671 detrended and expressed in arbitrary units relative to mean expression. Solid black
672 line, wildtype cells. Dashed lines, duplicate plates of cells expressing an *Sfpq*- or
673 *Pspc1*-targeting vector. **(D)** Quantification of depletion of SFPQ and PSPC1 protein
674 from experiments above. Relative repression from 3T3 cells cotransfected with a GFP-
675 expressing plasmid and a plasmid expressing an RNAi interference construct targeting
676 *Sfpq* or *Pspc1*. Averages shown are from 10 cells each, +/- SE. Mean fluorescence is
677 expressed in arbitrary units. **(E)** Quantification (+/- SE; n=2 independent experiments,
678 performed in triplicate) of *Neat1* levels for two different RNAi constructs (R and B), as
679 well as a scrambled hairpin (S) used in Fig 4, quantified from RNA of bulk-transfected
680 cells (black bars), or from cells cotransfected with a GFP-expressing plasmid and then
681 FACS-sorted to isolate GFP-expressing cells (grey bars).

682

683 **Fig 5 (A)** Immunofluorescence from cells transfected with a plasmid expressing GFP
684 and an RNAi interference construct targeting *Neat1* (*Neat-R*). Top panel, red filter
685 (PSPC1 protein). Bottom panel, same cells, green filter (GFP). White arrow,
686 paraspeckle in transfected cell; yellow arrow, paraspeckle in untransfected cell. Size
687 bar, 10µm. **(B)** Quantification (+/- SD) of paraspeckles per cell for two different RNAi
688 constructs (R and B), as well as a scrambled hairpin (S), quantified by immunostaining
689 similarly to (A), N= 12 cells (*Neat1-R*), 24 (*Neat1-B*), 18 (*Neat1-S*). Significance from
690 Student t-test, *<0.05, **<0.01. **(C)** Period length of circadian reporter expression for
691 U2oS cells cotransfected with the hairpins described in (A) and the *Bmal-luciferase*
692 circadian reporter. (N=6 per sample, no significant differences as determined by
693 Student t-test).

694

695 **Fig 6. (A)** Chromatin immunoprecipitation of the indicated proteins at the *Rev-*
696 *Erb α* promoter in liver nuclei harvested at different circadian times (CT) of day in
697 constant darkness. N=4, +/- SD, expressed relative to timepoint of minimum binding.
698 Ctrl reactions used an unrelated antibody raised at the same time in the same species
699 (anti-PAR-BZIP). **(B)** Immunoprecipitations from whole-cell extracts from 293T cells
700 cotransfected with myc-tagged NONO and Flag-tagged PER1 (left panel) or PER2 (right
701 panel). For each panel, left lane is 1/10 input, middle is immunoprecipitate with anti-
702 Myc antibody, and right lane is precipitation with IgG. Subsequent to
703 immunoprecipitation, all blots are probed with both anti-myc and anti-FLAG antibodies.
704 **(C)** Identical experiments performed with whole-cell extracts from 293T cells
705 cotransfected with myc-tagged SFPQ and Flag-tagged PER1 (left panel) or PER2 (right
706 panel). **(D)** Mouse liver nuclear extracts from ZT16 were immunoprecipitated with anti-
707 PER2 and probed with anti-NONO or anti-SFPQ. Left lane 1/10 input, right lane IP.

708

709 **Fig 7. (A)** Bioluminescence measured after transient transfection of mouse primary
710 fibroblasts from wildtype (black bars) or *per1^{brdm/brdm}/per2^{brdm/brdm}* double mutant
711 animals (striped bars) transfected with an *E-box-luciferase* reporter, and vectors
712 expressing CLOCK and BMAL proteins, and NONO, SFPQ, or PSPC1 as indicated.
713 N=3 experiments in duplicate, +/- SE, for all of figure **(B)** Chromatin
714 immunoprecipitation of NONO in wildtype (WT, black bars) or PER-deficient (p1/p2 mut,
715 striped bars) mice at the *Rev-Erb α* promoter in liver nuclei harvested at different
716 circadian times (CT) of day in constant darkness +/- SD . CT0 = beginning of

717 subjective day. N=3 **(C)** Transient transfection of fibroblasts with a *GAL4 DNA-binding*
718 *domain-Ebox-luciferase* reporter and vectors expressing GAL4-NONO, -PSPC1, -
719 SFPQ, or -VP16. Black bars, no exogenous activator added. Striped bars, vectors
720 expressing CLOCK and BMAL1 also added. **(D)** Identical experiments showing no
721 statistically significant effects using a *GAL4-CMV-luciferase* reporter without exogenous
722 activator.

723

724 **Fig 8. (A)** *Pspc1* RNA levels measured by qPCR from different tissues of genetrapped
725 mice (striped bars) and wildtype littermates (black bars). For parts A-B, N=2 mice per
726 measurement, measured 4x in duplicate, +/- SE. **(B)** *Sfpq* RNA levels measured by
727 qPCR from different tissues of genetrapped mice (striped bars) and wildtype littermates
728 (black bars). **(C)** PSPC1 protein levels in liver nuclear extracts harvested at different
729 times of day from wildtype and genetrapped animals kept in darkness. Top panel,
730 western blot probed with anti-PSPC1. Bottom panel, Ponceau-S staining of filter to
731 show equal loading. **(D)** SFPQ protein levels in liver nuclear extracts harvested at
732 different times of day from wildtype and genetrapped animals kept in darkness. Top
733 panel, western blot probed with anti-SFPQ. Bottom panel, Ponceau-S staining of filter
734 to show relative loading.





735



736 **Fig 9. (A)** Left, wheel-running activity of wildtype, *Pspc1^{gt/gt}*, and *Sfpq^{gt/+}* mice in 12:12
737 LD (arrow) and in constant darkness (DD). Darkness is indicated by grey shading. **(B)**
738 Period lengths of twelve mice of each genotype, together with wildtype littermates. No
739 significant differences for either *Pspc1* or *Sfpq* using Student t-test. **(C)** *Rev-Erb α* RNA






740 expression from *Nono*, *Sfpq*, and *Pspc1*-genetrapped mice (striped bars) and wildtype
741 littermates (black bars), measured by quantitative RT-PCR from liver extracts harvested
742 at different circadian times (CT) of day from mice in constant darkness. N=2 mice per
743 time point, RNA measured 4x in technical duplicate. Data shown is +/- SE.






744

745






















A. Transfection: *Neat1-R* + *GFP*










A.**B.****C.**

Ponceau red Staining

**D.**

Ponceau red Staining



A.**B.****C.**