

ID Design 2012/DOOEL Skopje
Open Access Macedonian Journal of Medical Sciences. 2015 Sep 15; 3(3):500-505.
<http://dx.doi.org/10.3889/oamjms.2015.086>
eISSN: 1857-9655
Public Health



Respiratory Symptoms and Lung Function in Never-Smoking Male Workers Exposed To Hardwood Dust

Dragana Bislimovska^{1*}, Sunchica Petrovska¹, Jordan Minov²

¹Department of Medical and Experimental Physiology with Anthropology, Medical Faculty, Ss Cyril and Methodius University of Skopje, Skopje, Republic of Macedonia; ²Institute of Occupational Health of Republic of Macedonia, WHO Collaborating Center, Skopje, Republic of Macedonia

Abstract

BACKGROUND: Results from many studies suggest that workplace exposure to organic dust may lead to adverse respiratory effects in exposed workers.

AIM: In order to assess the respiratory effects of the workplace exposure to hardwood dust we performed a cross-sectional study of never-smoking male workers employed in parquet manufacture and never-smoking male office workers as a control.

MATERIAL AND METHODS: We performed a cross-sectional study including 37 never-smoking male workers employed in parquet manufacture and an equal number of never-smoking male office workers studied as a control. Evaluation of examined subjects included completion of a questionnaire for respiratory symptoms in the last 12 months and baseline spirometry performed according to the actual recommendations.

RESULTS: We found a higher prevalence of respiratory symptoms in parquet manufacturers than in office workers with significant difference for cough and phlegm. Majority of the respiratory symptoms in the parquet manufacturers were work-related. The mean values of all spirometric parameters with exception of forced ventilatory capacity (FVC) were significantly lower in the parquet manufacturers as compared to their mean values in the office workers. We found close relationship between both the prevalence of respiratory symptoms and the reduction of spirometric parameters in the parquet manufacturers and the duration of the workplace exposure to wood dust.

CONCLUSION: Our data suggest that workplace exposure to hardwood dust may lead to adverse respiratory effects indicating the need of adequate preventive measures in order to protect the respiratory health of exposed workers.

Citation: Bislimovska D, Petrovska S, Minov J. Respiratory Symptoms and Lung Function in Never-Smoking Male Workers Exposed To Hardwood Dust. *OA Maced J Med Sci.* 2015 Sep 15; 3(3):500-505. <http://dx.doi.org/10.3889/oamjms.2015.086>

Key words: respiratory symptoms; questionnaire; spirometry; wood dust; workplace exposure.

***Correspondence:** Dr. Dragana Bislimovska, Department of Medical and Experimental Physiology with Anthropology, Medical Faculty, Ss Cyril and Methodius University of Skopje, 50 Divizija 6, Skopje, Republic of Macedonia. E-Mail: draganamk@yahoo.com

Received: 03-Jul-2015; **Revised:** 07-Jul-2015; **Accepted:** 13-Jul-2015; **Online first:** 16-Jul-2015

Copyright: © 2015 Dragana Bislimovska, Sunchica Petrovska, Jordan Minov. This is an open access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Competing Interests: The authors have declared that no competing interests exist.

Introduction

Results from a number of studies indicated that the exposure to organic dusts, such as cotton, flax, wood, paper, grain and other agricultural dusts, may lead to respiratory impairment and development of chronic lung disease, independently of any effect due to smoking [1-5].

Wood dust is created when machines are used to cut or shape wood materials. Industries with a risk of wood dust exposure include sawmills,

dimension mills, secondary wood products manufacture (parquet, doors, windows, etc.), furniture manufacture, cabinet makers, and carpenters. During their working activities woodworkers may be exposed to different exposure types, i.e. hardwood or softwood dust, dust from natural wood or wood-based composites, pure wood dust or wood dust containing adhesives, paints and other chemicals, etc [6, 7]. Woodworking at the global level is dominantly done by men. For example, with 98.5% of the workers being male, carpentry was the fourth male-dominated occupation in the USA in 1999, and there were about 1.5 million positions in 2006 [8].

There is evidence that workplace exposure to wood dust may cause adverse health effects in exposed workers, such as dermatitis, allergic and non-allergic respiratory effects and cancer. Dermatitis is a common health effect associated with exposure to wood dust. Wood, usually as sawdust or splinters, may affect the skin by mechanical action or by chemical irritation and sensitization. Respiratory effects in workers exposed to wood dust may also be caused by irritation or sensitization. A hypersensitivity reaction leading to asthma has been reported as a result of exposure to commonly used woods, including western red cedar, oak, mahogany, and redwood. The asthmatic reaction is believed to be species-specific. Respiratory symptoms and lung function decline can be caused by irritation of the bronchial tree, too. The main population of workers who suffer from respiratory problems are those who work in secondary wood products manufacturing, although these problems have been documented in sawmill workers. The National Institute for Occupational Safety and Health (NIOSH) considers both hardwood and softwood dust to be potentially carcinogenic for humans. The three types of cancer associated with wood dust exposure are nasal and sinus cavity cancer, lung cancer, and Hodgkin's disease. In addition, there is evidence that combined effect of wood dust and tobacco smoke increases the health risks in workers exposed to wood dust [7, 9-11].

In the present study we compared the prevalence of respiratory symptoms in the last 12 months and the values of the spirometric parameters, as well as its relation to duration of exposure between a group of never-smoking male workers exposed to natural hardwood dust and a group of unexposed male workers (office workers), matched by age and duration of employment at the actual workplace.

Methods

Study design and setting

A cross-sectional study was carried out in the Department of Cardiorespiratory Functional Diagnostics at the Institute for Occupational Health of R. Macedonia, Skopje - WHO Collaborating Center for Occupational Health and GA²LEN Collaborating Center in the period August – December 2014. Prevalence of respiratory symptoms in the last 12 months and mean values of spirometric parameters was compared between a group of never-smoking male workers exposed to hardwood dust and a group of never-smoking male office workers. All study subjects gave their informed consent before entering the study.

Subjects

We examined 37 male workers aged 33 to 62 years, employed in a company for parquet manufacture with duration of employment between 7 to 26 years. They worked in two large closed working areas in two working shifts lasting 8 hours and their working tasks included measuring, cutting and shaping different types of natural hardwood, such as oak, beech, ash and elm. The process control provided keeping of the exposure at permissible levels. During the working process the employees were protected by working outfits, gloves, glasses and masks. All examined parquet manufacturers were never-smokers, i.e. non-smokers who have never smoked at all, or have never been daily smokers and have smoked less than 100 cigarettes in their lifetime [12, 13].

In addition, an equal group of never-smoking male office workers matched to construction workers by age and duration of employment at the actual workplace was studied as a control.

In either group there were no subjects with chronic respiratory disease diagnosed by a physician (i.e. asthma, COPD, bronchiectasis, etc.), neither subject treated with bronchodilators and/or corticosteroids. In either group also there were no subjects in whom spirometry was contraindicated [14, 15].

Questionnaire

An interviewer-led questionnaire was completed by all study subjects. The questionnaire included questions on work history (e.g., chronological list of jobs; description of job activities at the actual workplace; type, extent and duration of exposure; and use of protective equipment), respiratory symptoms in the last 12 months and their relatedness to the workplace, chronic respiratory diseases diagnosed by a physician, family history of chronic bronchitis or asthma (taking into account the first-degree relatives), environmental exposure to tobacco smoke (ETS), accompanying disease, and medication use.

Respiratory symptoms in the last 12 months (cough, phlegm, dyspnea, wheezing, and chest tightness) were documented using the European Community for Coal and Steel questionnaire (ECCS-87), and the European Community Respiratory Health Survey (ECRHS) questionnaire [16, 17]. The work-relatedness of the respiratory symptoms was defined as more than usual cough, phlegm, dyspnea, wheezing, and chest tightness during daily work [18].

ETS or passive smoking or second-hand smoking was defined as an exposure to tobacco combustion products from smoking by others (at home, workplace, etc.), i.e. as a presence of at least one smoker in the household and/or at the workplace [19, 20].

Baseline spirometry

The baseline spirometry, including measures of forced vital capacity (FVC), forced expiratory volume in one second (FEV₁), FEV₁/FVC ratio, and maximal expiratory flow at 25%, 50%, 75%, and 25-75% of FVC (MEF₂₅, MEF₅₀, MEF₇₅, and MEF₂₅₋₇₅, respectively), was performed in all subjects using spirometer Ganshorn SanoScope LF8 (Ganshorn Medizin Electronic GmbH, Germany) recording the best result from three measurements of the values of FEV₁ which were within 5% of each other. The results of spirometry were expressed as percentages of the predicted values according to the actual recommendations of the European Respiratory Society (ERS) and the American Thoracic Society (ATS) [14, 15].

Statistical analysis

Continuous variables were expressed as mean values with standard deviation (SD), and the nominal variables as numbers and percentages. Analyses of the data involved testing the differences in prevalence, comparison of the means, as well as testing the association between respiratory symptoms in the last 12 months and mean values of the MEF parameters (less or more than 60% of the predicted value) and duration of the exposure at the actual workplace (less or more than 15 years). Chi-square test (or Fisher's exact test where appropriate) was used for testing difference in the prevalence. Comparison of spirometric measurements was performed by independent-samples T-test. A P-value less than 0.05 was considered as statistically significant. Statistical analysis was performed using the Statistical Package for the Social Sciences (SPSS) version 11.0 for Windows.

Results

Demographic characteristics of the study participants are given in Table 1.

Table 1: Demographics of the study subjects

Variable	Parquet manufacturers (n = 37)	Office workers (n = 37)
Age (years)	44.9 ± 7.8	45.2 ± 5.4
BMI (kg/m ²)	25.1 ± 3.1	25.8 ± 3.8
Duration of employment (actual workplace, years)	17.4 ± 4.3	18.1 ± 3.9
Duration of employment less than 15 years	21 (56.8%)	19 (51.3%)
Duration of employment more than 15 years	16 (43.2%)	18 (48.7%)
Family history of asthma/chronic bronchitis	5 (13.5%)	4 (10.8%)
Environmental ETS	18 (48.6%)	15 (40.5%)
Accompanying diseases		
Arterial hypertension	5 (13.5%)	5 (13.5%)
Diabetes mellitus type 2	2 (5.4%)	3 (8.1%)
Musculoskeletal disorders	6 (16.2%)	4 (10.8%)
Peptic ulcer	3 (8.1%)	4 (10.8%)

Numerical data are expressed as mean value with standard deviation; frequencies as number and percentage of study subjects with certain variable. BMI: body mass index; kg: kilogram; m: meter; ETS: exposure to tobacco smoke.

Prevalence of overall respiratory symptoms in the last 12 months was higher in parquet manufacturers than in office workers but the difference did not reach statistical significance.

Table 2: Prevalence of respiratory symptoms in the last 12 months

Respiratory symptoms in the last 12 months	Parquet manufacturers (n = 37)	Office workers (n = 37)	P-value*
Overall respiratory symptoms	16 (43.2%)	9 (24.3%)	0.092
Cough	11 (29.7%)	5 (13.5%)	0.044
Phlegm	6 (16.2%)	2 (5.4%)	0.021
Dyspnea	4 (10.8%)	3 (8.1%)	0.119
Wheezing	3 (8.1%)	2 (5.4%)	0.143
Chest tightness	5 (13.5%)	4 (10.8%)	0.138

Data are expressed as number and percentage of study subjects with certain variable. *Tested by Chi-square test (or Fisher's exact test where appropriate).

Prevalence of particular respiratory symptoms was also higher in parquet manufacturers than in office workers with statistically significant difference for cough and phlegm (Table 2, Figure 1 and Figure 2).




Figure 1: Prevalence of overall respiratory symptoms in the last 12 months

Majority of the respiratory symptoms in the last 12 months in parquet manufacturers were reported as work-related symptoms (13/16; 81.2%). The highest work-relatedness was reported for cough (8/11; 72.7%), phlegm (5/6; 83.3%) and dyspnea (3/4; 75.0%).




Figure 2: Prevalence of particular respiratory symptoms in the last 12 months

Work-relatedness of the respiratory symptoms was reported by three office workers with respiratory symptoms in the last 12 months (33.3%), i.e. by one office worker with cough, dyspnea and chest tightness (20.0%, 33.3% and 25.0%, respectively) (Figure 3).




Figure 3: Work-relatedness of the respiratory symptoms in the last 12 months

Prevalence of overall respiratory symptoms in the last 12 months was significantly lower in the parquet manufacturers with duration of exposure at the actual workplace longer than 15 years than in the parquet manufacturers with duration of exposure at the actual workplace shorter than 15 years. In addition, significant difference was found in the prevalence of cough and phlegm in the parquet manufacturers with duration of exposure longer than 15 years as compared to those with shorter duration of exposure at the actual workplace (Table 3). Such difference in the office workers was not registered neither for overall respiratory symptoms in the last 12 months, nor for any particular respiratory symptom in the last 12 months.

Table 3: Prevalence of respiratory symptoms in the last 12 months in the parquet manufacturers with duration of exposure at the actual workplace more and less than 15 years

Respiratory symptoms in the last 12 months	Duration of exposure > 15 years	Duration of exposure < 15 years	P-value*
Overall respiratory symptoms	12/16 (75.0%)	4/16 (25.0%)	0.037
Cough	8/11 (72.7%)	3/11 (27.3%)	0.040
Phlegm	5/6 (83.3%)	1/6 (16.7%)	0.014
Dyspnea	2/4 (50.0%)	2/4 (50.0%)	1.000
Wheezing	2/3 (66.6%)	1/3 (33.3%)	0.083
Chest tightness	2/5 (40.0%)	3/5 (60.0%)	0.108

Data are expressed as number and percentage of study subjects with certain variable. *Tested by Chi-square test (or Fisher's exact test where appropriate).

The mean values of baseline spirometric parameters with exception of the mean FVC value were significantly lower in the parquet manufacturers than in the office workers (Table 4).

Table 4: Mean baseline values of spirometric parameters

Spirometric parameter	Parquet manufacturers (n = 37)	Office workers (n = 37)	P-value*
FVC (%pred)	92.3 ± 14.7	94.2 ± 11.4	0.088
FEV ₁ (%pred)	82.6 ± 10.2	86.8 ± 9.4	0.034
FEV ₁ /FVC	0.75 ± 0.05	0.78 ± 0.03	0.037
MEF ₂₅ (%pred)	70.3 ± 8.9	76.9 ± 7.4	0.029
MEF ₅₀ (%pred)	64.6 ± 13.8	72.7 ± 10.5	0.011
MEF ₇₅ (%pred)	60.2 ± 16.7	69.7 ± 11.8	0.000
MEF ₂₅₋₇₅ (%pred)	65.9 ± 14.1	70.8 ± 12.9	0.006

Data are expressed as mean value with standard deviation. FVC: forced vital capacity; FEV₁: forced expiratory volume in one second; MEF₇₅, MEF₅₀, MEF₂₅, MEF₂₅₋₇₅: maximal expiratory flow at 25, 50%, 75% and 25-75% of FVC, respectively; % pred: % of predicted value. *Compared by Independent-samples T-test.

The mean values of all spirometric parameters were significantly lower in the parquet manufacturers with duration of exposure at the actual workplace longer than 15 years as compared to those with duration of exposure at the actual workplace shorter than 15 years (Table 5). Such difference in the office workers was registered only for MEF₇₅ and MEF₂₅₋₇₅ (71.8% vs. 67.4%; $P = 0.028$ and 72.6% vs. 68.1%; $P = 0.037$, respectively).

Table 5: Mean baseline values of spirometric parameters in the parquet manufacturers with duration of exposure at the actual workplace longer and shorter than 15 years

Spirometric parameter	Duration of exposure > 15 years	Duration of exposure < 15 years	P-value*
FVC (%pred)	90.4 ± 12.6	93.2 ± 10.9	0.038
FEV ₁ (%pred)	79.2 ± 9.8	84.3 ± 11.4	0.029
FEV ₁ /FVC	0.74 ± 0.02	0.76 ± 0.04	0.022
MEF ₂₅ (%pred)	68.5 ± 11.7	72.6 ± 10.8	0.012
MEF ₅₀ (%pred)	61.3 ± 10.2	67.1 ± 11.9	0.002
MEF ₇₅ (%pred)	56.1 ± 13.4	64.2 ± 12.8	0.000
MEF ₂₅₋₇₅ (%pred)	62.7 ± 14.3	68.9 ± 12.1	0.003

Data are expressed as mean value with standard deviation. FVC: forced vital capacity; FEV₁: forced expiratory volume in one second; MEF₇₅, MEF₅₀, MEF₂₅, MEF₂₅₋₇₅: maximal expiratory flow at 25, 50%, 75% and 25-75% of FVC, respectively; % pred: % of predicted value. *Compared by Independent-samples T-test.

Discussion

Respiratory effects of workplace exposure to wood dust are evaluated in a number of studies. These studies produced somewhat inconsistent results depending upon the type, level and duration of the workplace exposure to wood dust, the characteristics of the examined workers (age, smoking status, etc.) and the type of their work activities, as well as upon the type of the study.

In the present study we compared the prevalence of respiratory symptoms in the last 12 months and the mean values of spirometric parameters, as well as their relation to duration of the workplace exposure between the group of never-smoking male workers exposed to hardwood dust and the group of never-smoking male office workers matched by age and duration of employment at the actual workplace. Both groups had similar demographic characteristics regarding the body mass index value, family history of asthma/chronic bronchitis, environmental exposure to ETS and accompanying diseases.

We found higher prevalence of the respiratory symptoms in the last 12 months in the parquet manufacturers as compared to their prevalence in the office workers with statistically significant difference for cough and phlegm. Majority of the respiratory symptoms in the last 12 months in the parquet manufacturers were work-related and there was a close relationship between their prevalence and the duration of the workplace exposure to wood dust. Similar results were obtained in several studies which investigated the relationship between respiratory symptoms and workplace exposure to wood dust, such as the cross-sectional study performed by Okwari et al. including 221 workers exposed to wood dust in timber markets from Calabar, Nigeria [21], the cross-sectional study performed by Arbak et al. including 64 Turkish furniture-decoration students [22], the cross-sectional study performed by Schünssen et al. including 54 Danish furniture factory workers [23], and the cross-sectional study performed by Douwes et al. including 772 New Zealander pine sawmill workers [24]. Higher prevalence of respiratory symptoms in workers exposed to wood dust than in office workers was also reported by Ige & Onadeko [25] in a cross-sectional study including 500 sawmillers from Ibadan, Nigeria, but the prevalence of phlegm was twofold higher than its prevalence registered in the actual study (34.3% vs. 16.2%) that may be due to the specific wood types and specific work activities of these workers. In addition, in a cross-sectional study including workers employed in dusty occupations, Danilova et al. [26] reported similar prevalence of respiratory symptoms in furniture manufacturers in which the workplace exposure was somewhat different as compared to the workplace exposure of the wood workers examined in the actual study.

The mean values of the spirometric parameters in the parquet manufacturers were lower as compared to their mean values in the office workers. Statistically significant difference was registered for all measured spirometric parameters with exception of the mean values of FVC. Similar results were obtained in several studies which investigated the effect of different types of wood dust on the lung function of exposed workers. In a cross-sectional study including 50 males working in sawmills in different areas in Bhavnagar city, India, Kacha et al. registered significantly lower values of FVC, FEV₁, MEF₅₀, MEF₂₅ and PEF_R as compared to their predicted values [27]. In a cross-sectional study including 328 smoking and non-smoking woodworkers, Osman & Pala registered significantly lower mean values of FVC and FEV₁ in both smokers and non-smokers as compared to their mean values in the controls [28]. In addition, in a cross-sectional study including 46 non-smoking Pakistani woodworkers, Meo registered significantly lower mean values of FVC and FEV₁ as compared to their mean values in unexposed controls [29]. Dudhmal et al. in

their study, including 30 Indian non-smoking sawmill workers, found significantly lower mean values of FEV₁ and PEF_R as compared to their predicted values, whereas regarding the mean FVC value such difference was not found [30]. In the actual study we found significant relation between the reduction of the mean values of the measured spirometric parameters and the duration of exposure to wood dust. Similar results were registered in the cross-sectional studies performed by Kacha et al. and Meo [27, 29] mentioned above. In a cross-sectional study including 145 South African non-smoking woodworkers and 152 matched controls, Shamssain reported significantly lower FEV₁/FVC ratio in woodworkers than in the controls. Furthermore, the FEV₁/FVC ratio was significantly lower in woodworkers with duration of employment longer than 10 years as compared to those exposed shorter than 10 years [31]. On the contrary, in a longitudinal study including 31 Italian non-smoking forestry workers with assessment of the lung function at least five times in the average period of 12 years, Inocenti et al. did not register significant difference in the VC and FEV₁ loss between the woodworkers and the controls [32].

The present study had some limitations. First, relatively small number of subjects in the study groups could have certain implications on the data obtained and its interpretation. Second, environmental measurements were not performed, so we could not document the effect of the type and the level of workplace exposure in these workers. Third, as in the case of any cross-sectional study, the impact of healthy workers' effect (HWE) on the data obtained could not be excluded. The strength of the study is the investigation of respiratory effects of specific occupational exposure in the wood-working environment independently from the effects of tobacco smoke.

In conclusion, in a cross-sectional study including never-smoking male parquet manufacturers we found higher prevalence of respiratory symptoms in the last 12 months and lower values of spirometric parameters than in matched office workers. There was a close relationship both between the prevalence of the respiratory symptoms in the last 12 months and the reduction of the spirometric parameters and the duration of the workplace exposure to wood dust. Our results suggest adverse respiratory effects of workplace exposure to hardwood dust indicating the need of reduction of harmful workplace exposure via proper engineering control, as well as the need of regular medical examinations in order to identify affected workers and thereby institute adequate measures.

Authors Participations

DB participated in the study design, writing

the protocol, data collection, managing the analyses of the study, and writing all versions of the manuscript. SP participated in the study design, writing the protocol, managing the analyses of the study, as well as writing all versions of the manuscript. JM performed the statistical analysis and participated in the managing of the analyses of the study. DB and JM participated in the data collection and in the managing of the analyses of the study. All authors read and approved the final manuscript.

References

- Chan-Yeung M, Dimich-Ward H. Natural history of occupational lung diseases. *Eur Respir Mon.* 1999; 4: 46-63.
- Dimich-Ward H, Kennedy S, Chan-Yeung M. Occupational exposures and chronic airway limitation. *Can Respir J.* 1996; 3: 133-140.
- Lamprecht B, Schirrhofer L, Kaiser B, Studnicka M, Buist AS. Farming and the prevalence of non-reversible airways obstruction: results from a population-based study. *Am J Ind Med.* 2007; 50 (6): 421-426.
- Minov J, Karadzinska-Bislimovska J, Vasilevska K, Risteska-Kuc S, Stoleski S. Occupational asthma in subjects occupationally exposed to herbal and fruit tea dust. *Arh Hig Rada Toksikol.* 2007; 58: 211-221.
- Glindemeyer HW, Lefante JJ, Jones RN, Rando RJ, Weill H. Cotton dust and across-shift change FEV1 as predictors of annual change in FEV1. *Am J Respir Crit Care Med.* 1994; 149: 584-590.
- Forest Products Laboratory. Wood Handbook. Available at: www.woodweb.com (Assessed 22.06.2015).
- Bean TL, Butcher TW, Lawrence T. Wood dust exposure hazards. Available at: http://ohioline.osu.edu/aefact/0595_1.html (Assessed 22.06.2015).
- Greg R. A Career as a Carpenter. Available at: www.rosenpublishing.com (Assessed 22.06.2015).
- Noertjojo HK, Dimich-Ward H, Peelen S, Dittrick M, Kennedy S, Chan-Yeung M. Western red cedar dust exposure and lung function loss: a dose-response relationship. *Am J Respir Crit Care Med.* 1996; 154: 968-973.
- Pramanik P, Chaudhury A. Impact of occupational exposure to wood dust on pulmonary health of carpenters in small scale furniture industries in West Bengal. *DHR-IJBLS.* 2013; 4 (3): 204-211.
- Hayes RB, Gerin M, Raatgeven RJ, Bruyn A. Wood-related occupations, wood dust exposure, and sinonasal cancer. *Am J Epidemiol.* 1986; 124 (4): 569-577.
- World Health Organization. Guidelines for controlling and monitoring the tobacco epidemic. Geneva: WHO, 1998.
- Leffondre K, Abrahamowicz M, Siemiatycki J, Rachet B. Modelling smoking history: A comparison of different approaches. *Am J Epidemiol.* 2002; 156: 813-823.
- Spirometry Guide: 2010 Update. http://www.goldcopd.org/uploads/users/files/GOLD_Spirometry_2010.pdf.
- Miller MR, Hankinson J, Brusasco V, Burgos F, Casaburi R, Coates A, Crapo R, Enright P, van der Grinten CPM, Gustafsson P, Jensen R, Johnson DC, MacIntyre N, McKay R, Navajas D, Pedersen OF, Pellegrino R, Viegi G, Wanger J. Standardisation of spirometry. *Eur Respir J.* 2005; 26: 319-338.
- Minette A. Questionnaire of the European Community for Coal and Steel (ECSC) on respiratory symptoms. 1987 - updating of the 1962 and 1967 questionnaires for studying chronic bronchitis and emphysema. *Eur Respir J.* 1989; 2:165-177.
- European Community Respiratory Health Survey. Variations in the prevalence of respiratory symptoms, self-reported asthma attacks, and use of asthma medication in the European Respiratory Health Survey (ECRHS). *Eur Respir J.* 1996; 9: 687-695.
- Meijer E, Grobde DE, Heederik DJJ. Health surveillance for occupational chronic obstructive disease. *J Occup Environ Med.* 2001; 43: 444-450.
- U.S. Department of Health and Human Services. The health consequences of smoking: chronic obstructive pulmonary disease. A report of the Surgeon General. US Department of Health and Human Services, Public Health Service, Office of the Assistant for Health, Office of Smoking and Health. DHHS Publication No. 84-50 205, 1984.
- Janson C, Chinn S, Jarvis D, Zock JP, Toren K, Burney P, for the European Community Respiratory Health Survey. Effects of passive smoking on respiratory symptoms, bronchial responsiveness, lung function, and total serum IgE in the European Community Respiratory Health Survey: a cross-sectional study. *Lancet.* 2001; 358: 2103-2109.
- Okwari OO, Antai AB, Owu DU, Peters EJ, Osim EE. Lung function status of workers exposed to wood dust in timber markets in Calabar, Nigeria. *Afr J Med Med Sci.* 2005; 34(2): 141-145.
- Arbak P, Bilgin C, Balbay O, Yesidal N, Annakkaya AN, Ugler F. Respiratory symptoms and peak expiratory flow rates among furniture-decoration students. *Ann Agric Environ Med.* 2004; 11: 13-17.
- Schlünssen V, Schaumburg J, Taudorf E, Mikkelsen AB, Sigsgaard T. Respiratory symptoms and lung function among Danish woodworkers. *J Occup Environ Med.* 2002; 44(1): 82-98.
- Douwes J, McLean D, Slater T, Pearce N. Asthma and other respiratory symptoms in New Zealand pine processing sawmill workers. *Am J Ind Med.* 2001; 39 (6): 608-615.
- Ige OM, Onadeko OB. Respiratory symptoms and ventilator function of the sawmillers in Ibadan, Nigeria. *Afr J Med Med Sci.* 2000; 29(2): 101-104.
- Danilova M, Stoleski S, Mijakoski D. Respiratory symptoms and ventilatory function in never-smoking males working in dusty occupations. *OA Maced J Med Sci.* 2014; 2(4): 645-649.
- Kacha Y, Nayak Y, Varu M, Mehta H, Shah CJ. Effects of wood dust on respiratory functions in saw mill workers. *International journal of basic and applied physiology.* 2014; 3(1): 122-128.
- Osman E, Pala K. Occupational exposure to wood dust and health effects on the respiratory system in a minor industrial estate in Bursa, Turkey. *Int J Occup Med Environ Health.* 2009; 22(1): 43-50.
- Meo SA. Lung function in Pakistani wood workers. *Int J Environ Health Res.* 2006; 16(3): 193-203.
- Dudhmal VB, Afroz S, Jadhav SS, Karadkhedkar SS. Pulmonary function tests in saw mill factory workers. *Indian J Physiol Pharmacol.* 2006; 50(3): 313-315.
- Shamssain MH. Pulmonary function and symptoms in workers exposed to wood dust. *Thorax.* 1992; 47(2): 84-87.
- Innocenti A, Ciapini C, Natale D, Nerozzi F, Pippi P, Selmi M. Longitudinal changes of pulmonary functions in workers with high wood dust exposure levels. *Med Lav.* 2006; 97(1): 30-35.