

doi:10.1186/s12876-015-0201-5

which should be cited to refer to this work.

## GES-type and OXA-23 carbapenemase-producing *Acinetobacter baumannii* in Turkey

Arzu Nazli Zeka<sup>1,2</sup>, Laurent Poirel<sup>1,3\*</sup>, Oguz Resat Sipahi<sup>2</sup>, Rémy A. Bonnin<sup>1</sup>, Bilgin Arda<sup>2</sup>, Memetali Özinel<sup>4</sup>, Sercan Ulusoy<sup>2</sup>, Canan Bor<sup>5</sup> and Patrice Nordmann<sup>1,3</sup>

<sup>1</sup>INSERM U914 'Emerging Resistance to Antibiotics', Le Kremlin-Bicêtre, France; <sup>2</sup>Department of Infectious Diseases and Clinical Microbiology, Ege University Medical Faculty, İzmir, Turkey; <sup>3</sup>Medical and Molecular Microbiology Unit, Department of Medicine, Faculty of Science, University of Fribourg, Fribourg, Switzerland; <sup>4</sup>Department of Medical Microbiology, Ege University Medical Faculty, İzmir, Turkey; <sup>5</sup>Intensive Care and Reanimation Unit, Ege University Medical Faculty, İzmir, Turkey

\*Corresponding author. Medical and Molecular Microbiology Unit, Department of Medicine, Faculty of Science, University of Fribourg, Rue Albert Gockel 3, Fribourg, Switzerland. Tel: +41-26-300-9582; Fax: +41-26-300-9582; E-mail: laurent.poirel@unifr.ch

**Keywords:** GES-11, *A. baumannii*,  $\beta$ -lactamases

Sir,  
*Acinetobacter baumannii* is one of the most common organisms causing healthcare-associated infections in hospitals, especially within intensive care units (ICUs). Carbapenem resistance in *A. baumannii* is most often mediated by acquired class D carbapenem-hydrolysing  $\beta$ -lactamases and in particular OXA-23.<sup>1</sup> Recently, the emergence of Ambler class A GES-type carbapenemases in *A. baumannii* has been demonstrated, being identified in France, Belgium and Kuwait.<sup>2-4</sup>

The aim of our study was to analyse the molecular epidemiology and clinical features of carbapenem-resistant *A. baumannii* (CRAB) clinical isolates recovered in a Turkish hospital. Isolates were recovered from patients hospitalized in two ICUs (respiratory and anaesthesiology) at the 2000 bed Ege University Medical School Hospital located in İzmir, western Turkey, from 1 March to 1 September 2012 (a 6 month period). The total number of patients in the two ICUs over the study period was 318. The total number of patient days (length of stay  $\times$  patient number) was 5935.

The CRAB isolates were recovered from clinical specimens of patients diagnosed with ventilator-associated pneumonia (VAP) and/or bacteraemia. Patients who were colonized with *A. baumannii*, but without any sign of infection, were excluded from the study.

A total of 60 patients (24 females and 36 males; mean age  $64 \pm 2$  years) were included in the study. The 60 CRAB isolates were identified by incubation at  $44^\circ\text{C}$  and by using the Vitek 2 identification system (bioMérieux, La Balme-les-Grottes, France). They

were then further confirmed by specific PCR targeting the *bla*<sub>OXA-51</sub>-like genes intrinsic to *A. baumannii*.<sup>2</sup> The mean APACHE II score of the patients was  $19 \pm 7.2$ . Forty-five patients (75%) had VAP and 15 (25%) had bacteraemia. None of these patients had received an appropriate empirical antimicrobial therapy. Fifteen (10 bacteraemia and 5 VAP) patients died before having being treated with an appropriate antimicrobial therapy within the first 48 h (mortality rate 100%). The overall crude mortality rate for the 45 patients for whom the antimicrobial treatment was appropriate was 55.6% ( $P < 0.001$ ). The antibiotic regimens were mostly combinations, being either colistin/tigecycline ( $n = 20$ ), colistin/netilmicin ( $n = 12$ ), colistin/meropenem ( $n = 3$ ), tigecycline/meropenem ( $n = 6$ ) or colistin monotherapy ( $n = 4$ ) (Table 1).

Antibiotic susceptibility was determined by disc diffusion and MICs of tigecycline, colistin, imipenem and meropenem were determined by using Etest strips (AB bioMérieux, La Balme-les-Grottes, France) and interpreted according to the updated CLSI guidelines when applicable.<sup>5</sup> All of the 60 isolates were resistant to all  $\beta$ -lactams, including broad-spectrum cephalosporins and carbapenems, and to fluoroquinolones. Resistance rates to amikacin, gentamicin and netilmicin were 73%, 65% and 36%, respectively. In addition, 33% of the isolates were resistant to all aminoglycosides tested. Four isolates showed reduced susceptibility to tigecycline (MICs of 1 mg/L) and five isolates were highly resistant (MICs of 3 mg/L). All of the isolates remained susceptible to colistin.

Identification of carbapenemase and class A extended-spectrum  $\beta$ -lactamase (ESBL) genes was performed by PCR and sequencing,<sup>2-4</sup> showing that the 60 isolates were all positive for the *bla*<sub>OXA-23</sub> carbapenemase gene. In addition, five isolates were positive for the *bla*<sub>GES-11</sub> gene, encoding an ESBL possessing a weak carbapenemase activity.<sup>6</sup> PCR mapping identified the *aacA4* gene cassette encoding the AAC(6')-Ib aminoglycoside acetyltransferase together with the *dfrA7* gene cassette encoding resistance to trimethoprim downstream of *bla*<sub>GES-11</sub>.

In order to determine the genetic location of carbapenemase- or ESBL-encoding genes, conjugation assays were performed using *A. baumannii* BM4547 (rifampicin resistant) as the recipient strain, as described previously.<sup>2</sup> Mating-out assays allowed *A. baumannii* BM4547 transconjugants harbouring the *bla*<sub>GES-11</sub> gene to be obtained. Those *A. baumannii* transconjugants expressed both OXA-23 and GES-11 and the corresponding *bla*<sub>GES-11</sub> and *bla*<sub>OXA-23</sub> genes were identified on a single 100 kb plasmid. The transconjugants showed resistance to all  $\beta$ -lactams, including carbapenems and aminoglycosides. Plasmid typing performed by *A. baumannii* PCR-based replicon typing<sup>7</sup> identified the 100 kb plasmid as belonging to group 6, carrying the two carbapenemase genes in all transconjugants.

Genotypic analysis of the 60 clinical isolates was performed by PFGE and allowed them to be grouped into five distinct pulsotypes (data not shown). This grouping was further confirmed by using the Diversilab technique (bioMérieux), which indicated that the five GES-11-positive isolates belonged to three different clonal lineages, with three isolates belonging to a single clone (data not shown). Multilocus sequence typing performed as described previously<sup>8</sup> on the five GES-11-positive *A. baumannii* isolates revealed

**Table 1.** Clinical characteristics of patients infected by the GES-producing *A. baumannii* isolates

Age (years)	Gender	Comorbidity	Infection	APACHE II score	Length of stay in ICU <sup>a</sup>	Empirical treatment	Treatment after culture	Outcome
48	female	connective tissue disorder	VAP	21	9	meropenem	tigecycline + colistin	death
67	male	solid organ tumour + abdominal surgery	VAP	32	14	meropenem	netilmicin + colistin	death
83	male	multiple trauma	VAP + bacteraemia	11	5	meropenem	—	death
79	male	solid organ tumour	bacteraemia	10	5	levofloxacin	tigecycline + meropenem	survival
49	female	solid organ tumour + abdominal surgery	bacteraemia	15	14	ceftriaxone	tigecycline + netilmicin	survival

<sup>a</sup>Number of days before isolation of *A. baumannii*.

that they all belonged to ST2, being part of the worldwide-distributed clone II group.

The current emergence of CRAB is a significant source of concern worldwide. Strains producing the carbapenemases OXA-58 and OXA-23 have been previously identified in Turkey. Here, only OXA-23 producers were identified, despite the fact that five clones were concomitantly identified in this hospital. The clonal diversity identified here revealed a polyclonal outbreak situation in the ICUs during that period of time. As previously reported, the overall mortality of the patients infected with those CRABs was very high, especially when they did not receive adequate empirical therapy. According to the resistance profile of the studied isolates, colistin was the most effective antimicrobial.

While our work was in progress, another study reported the occurrence of GES-11 and GES-12 enzymes among *A. baumannii* isolates in Turkey.<sup>9</sup> This highlights the need for implementation of adequate screening techniques to identify patients colonized by those multidrug-resistant strains.

## Funding

This work was mostly funded by the INSERM, France, and by grants from the European Community (R-GNOSIS, FP7/HEALTH-F3-2011-282512, and MAGIC-BULLET, FP7/HEALTH-F3-2001-278232). A. N. Z. was supported by an overseas research grant from EKMUD (Turkish Infectious Diseases and Clinical Microbiology Specialist Society).

## Transparency declarations

None to declare.

## References

- 1 Poirel L, Nordmann P. Carbapenem resistance in *Acinetobacter baumannii*: mechanisms and epidemiology. *Clin Microbiol Infect* 2006; **12**: 826–36.
- 2 Bonnin RA, Nordmann P, Potron A *et al.* Carbapenem-hydrolyzing GES-type extended-spectrum  $\beta$ -lactamase in *Acinetobacter baumannii*. *Antimicrob Agents Chemother* 2011; **55**: 349–54.
- 3 Bonnin RA, Rotimi VO, Al Hubail M *et al.* Wide dissemination of GES-type carbapenemases in *Acinetobacter baumannii* isolates in Kuwait. *Antimicrob Agents Chemother* 2013; **57**: 183–8.
- 4 Bogaerts P, Naas T, El Garch F *et al.* GES extended-spectrum  $\beta$ -lactamases in *Acinetobacter baumannii* isolates in Belgium. *Antimicrob Agents Chemother* 2010; **54**: 4872–8.
- 5 Clinical and Laboratory Standards Institute. *Performance Standards for Antimicrobial Susceptibility Testing: Twenty-third Informational Supplement M100-S23*. CLSI, Wayne, PA, USA, 2013.
- 6 Moubareck C, Brémont S, Conroy MC *et al.* GES-11, a novel integron-associated GES variant in *Acinetobacter baumannii*. *Antimicrob Agents Chemother* 2009; **53**: 3579–81.
- 7 Bertini A, Poirel L, Mugnier PD *et al.* Characterization and PCR-based replicon typing of resistance plasmids in *Acinetobacter baumannii*. *Antimicrob Agents Chemother* 2010; **54**: 4168–77.
- 8 Diancourt L, Passet V, Nemeč A *et al.* The population structure of *Acinetobacter baumannii*: expanding multiresistant clones from an ancestral susceptible genetic pool. *PLoS One* 2010; **5**: e10034.
- 9 Cicek AC, Saral A, Iraz M *et al.* OXA- and GES-type  $\beta$ -lactamases predominate in extensively drug-resistant *Acinetobacter baumannii* isolates from a Turkish University Hospital. *Clin Microbiol Infect* 2013; doi:10.1111/1469-0691.12338.