

RESEARCH ARTICLE

Open Access

Laryngeal Mask Airway for neonatal resuscitation in a developing country: evaluation of an educational intervention. Neonatal LMA: an educational intervention in DRC

Vincenzo Zanardo^{1*}, Alphonse Simbi¹, Massimo Micaglio², Francesco Cavallin¹, Leon Tshilolo¹, Daniele Trevisanuto¹

Abstract

Background: Studies carried out in developing countries have indicated that training courses in newborn resuscitation are efficacious in teaching local birth attendants how to properly utilize simple resuscitation devices. The aim of this study was to assess the knowledge and expertise gained by physicians and midwives who participated in a Neonatal Resuscitation Course and workshop organized in a Third World Country on the use of Laryngeal Mask Airway (LMA).

Methods: A 28-item questionnaire, derived from the standard test contained in the American Heart Association and the American Academy of Pediatrics Neonatal Resuscitation Manual, was administered to 21 physicians and 7 midwives before and after a course, which included a practical, hands-on workshop focusing on LMA positioning and bag-ventilation in a neonatal manikin.

Results: The knowledge gained by the physicians was superior to that demonstrated by the midwives. The physicians, in fact, demonstrated a significant improvement with respect to their pre-course knowledge. Both the physicians and the midwives showed a good level of expertise in manipulating the manipulating the manikin during the practical trial session. The midwives and physicians almost unanimously manifested a high degree of approval of neonatal resuscitation by LMA, as they defined it a sustainable and cost-effective method requiring minimal expertise.

Conclusions: Further studies are warranted to test the advantages and limits of the neonatal LMA training courses in developing countries.

Background

Intrapartum hypoxia and birth asphyxia are widely regarded as major causes of morbidity and mortality in developing countries [1,2]. All those involved in delivery room care must, consequently, possess the knowledge and expertise to perform neonatal resuscitation [3].

Maintaining a patent airway and providing effective positive pressure ventilation (PPV), which is currently achieved in the delivery room by means of a face mask

(FM) or a tracheal tube (TT), is standard treatment and forms the cornerstone of emergency neonatal care in First World countries [2], but making these interventions feasible in settings where resources are limited is particularly challenging [4,5]. Since its introduction into clinical practice, the LMA has gained increasing popularity for resuscitation of adult as well as pediatric patients, and more recently in neonatal resuscitation. Some organizations have, in fact, made it a part of their guidelines and LMA has been included in the AAP and AHA Guidelines of Neonatal Resuscitation Program (NRP) since 2000 [3,6]. It has been seen that LMA provides a low pressure airtight seal against the glottis

* Correspondence: zanardo@pediatria.unipd.it

¹Department of Pediatrics, Padua University, (3, Via Giustiniani) Padua (35128), Italy

Full list of author information is available at the end of the article

[7-10] and studies on the efficacy of ventilation by medical and paramedical personnel in neonatal training models have shown that it is characterized by ease of insertion and rapid, adequate airway patency [11]. Although this device cannot be considered a substitute for the TT, it could, at least theoretically, play an ancillary role in developing countries [12] where it offers practical, cost-effective, sustainable advantages over the face mask.

Some studies carried out in Third World countries have indicated that local birth attendants who have been trained in newborn resuscitation are capable of learning and properly utilizing simple resuscitation techniques (e.g. Mouth-to-mask breathing and room air) [4], reducing asphyxia-related deaths [13-16]. It is uncertain, however, what aspects of these training programs were responsible for that outcome.

The aim of this study is to assess the knowledge gained by local birth attendants (physicians and midwives) who participated in a NRP course and workshop on LMA organized in the Democratic Republic of Congo (DRC).

Methods

A 3-day NRP course was held by the Continuing Education for Africa (CEFA), in Kinshasa, the DRC, in September 2006. The course consisted of a number of didactic sessions proportionally divided into the seven steps of the NRP: (I) principles of resuscitation; (II) initial steps in resuscitation; (III) bag mask ventilation; (IV) chest compressions; (V) tracheal intubation; (VI) medications; and (VII) special considerations [3], followed by practical hands-on training session, including LMA positioning and bag-ventilation in a neonatal manikin (Neonate Airway Trainer; Laerdal, Norway).

Twenty-eight local birth attendants, (21 physicians and 7 midwives) from the Congo Brazzaville (1), Benin (1), Cameroun (1), and the remaining from other areas of the DRC, participated in the course. All of the participants took - both before and after the course - a 28 question test, an adaptation of a standard one contained in the AHA/AAP Neonatal Resuscitation Manual [3] and underwent a practical trial evaluating their proficiency in manipulating LMA and a neonatal manikin. The participants had 60 minutes to complete the test, which included multiple choice, fill-in-the blank, and true/false questions. The test, which was written in French, was strictly supervised.

The questions concerned: (I) features of LMA; (II) advantages of LMA over the face mask; (III) advantages of LMA over the endotracheal tube; (IV) disadvantages of LMA; and (V) potential applications in neonatal resuscitation. Additional file 1

The theoretical test was followed by a practical trial during which the trainees were asked to attempt LMA insertion in a manikin (designed for skills training in neonatal resuscitation) in less than 15 seconds (successful insertion was defined as effective thorax expansion verified by the instructor). The participants were also asked to anonymously express their opinion about their degree of satisfaction (high/low) with the course and the sustainability and cost-effectiveness (yes/no) of LMA in their respective countries.

The study was approved by the Ethical and Scientific Committee of the CEFA Agency.

Statistical analysis

"Improvement", defined as the difference (number or percentage) between the pre and post test scores, in the two groups (physicians and midwives) was analyzed separately by the McNemar test and Fisher's exact test, as appropriate. A post hoc multiple comparison analysis using the Bonferroni's adjustment was performed when statistically significant ($p < 0.05$) differences were found in the groups. A p value < 0.05 was considered significant.

Results

Theoretical knowledge about and practical skills in neonatal resuscitation using LMA on a neonatal manikin gained by the participants (physicians and midwives) at a neonatal resuscitation course are outlined in Tables 1 and 2.

Compared with the initial score, the overall knowledge gained by the physicians and midwives who participated in the NRP course and workshop on neonatal resuscitation by LMA was increased, but with different percentages regarding the single test questions. In particular, physicians significantly improved in 12/28 post-test items analyzed by McNemar's test and in 7/12 by Bonferroni's adjustment. Table 1

The "improvement", defined by the number of correct answers in post-test coupled with a wrong answer in pre-test, was statistically significant for the physicians and midwives with reference to the questions concerning their theoretical knowledge about LMA. The "improvement", defined by the percent of correct answers in post-test coupled with a wrong answer in pre-test, with regard to the questions dealing with the advantages of LMA over the facial mask ($p < 0.02$) and its potential applications ($p < 0.005$) was statistically significant for the physicians alone. Table 2

The degree of approval by the physicians and midwives of neonatal resuscitation by LMA and their opinions regarding its sustainability and cost-effectiveness are outlined in Table 3.

Table 1 Knowledge gained by participants during a NRP training course on neonatal resuscitation focusing on LMA

Neonatal Laryngeal Mask Airway Features	Pre-test correct answers n.(%)		Post-test correct answers n.(%)	
	Physicians: n. 21	Midwives: n.7	Physicians: n.21	Midwives: n.7
Range	5	3	7	3
Size	6	1	10	3
Lubrication	3	2	11 [^]	4
Tip flattening	5	3	11 [^]	3
Advancing	5	2	10	4
Location	4	1	12 [^]	3
Cuff inflation	5	0	11 [^]	3
Connection for PPV	13	5	11	5
Correct positioning	1	0	14*	7
	47/189 (24.8)	17/63 (26.9)	97/189 (51.3)	35/63 (55.5)
Advantages over the face mask				
Less skill required	4	0	11 [^]	6
Easy placement	3	1	4	2
Better airtight seal	1	1	12*	3
Improved SO ₂	3	1	11	6
Function unaffected by anatomical factors	1	1	6	1
Less hand fatigue	7	0	10	7
	19/126 (15.0)	4/42 (9.5)	54/126 (42.8)	25/42 (9.52)
Advantages over the tracheal tube				
Easier and quicker placement by trained medical and non-medical personnel	5	2	10	4
Avoids laryngoscopy	2	5	14*	7
Avoids tracheal oedema	4	0	14*	6
Efficacy in upper airway malformations when intubation and mask ventilation fail	9	4	14	6
Avoids use of neuromuscular blocking agents	4	3	15*	7
	24/105 (22.8)	14/35 (40.0)	67/105 (63.8)	30/35 (85.7)
Disadvantages				
Gastric insufflation and aspiration	0	0	8	4
Inadequate alveolar ventilation	13	3	14	6
Impossibility of suctioning the airway	4	3	8	6
Impossibility of administering drug endotracheally	12	3	14	7
	29/84 (34.5)	9/28 (32.1)	44/84 (52.3)	17/28 (60.7)
Potential applications in neonatal resuscitation				
When face mask and tracheal tube resuscitation fail	10	6	15	7
In neonatal training models allows a patent airway in a shorter time than endotracheal tube	4	0	14*	6
Incidence of failure low with LMA	10	5	15	7
Ruinously	5	2	15*	7
	29/84 (34.5)	13/28 (46.4)	59/84 (70.2)	27/28 (96.4)

Pre- and post-test correct answers.

[^]McNemar test p value (p < 0.05), corrected by Bonferroni's adjustment (p < 0.05)*

Table 2 "Improvement", defined as the difference between the pre- and post-test scores

Neonatal LMA		Improvement n.	McNemar Test [§]	Improvement *%	Fisher test*
Features	Physicians	50	p < 0.0001	35.2	p < 0.72
	Midwives	18	p < 0.0001	40.9	
Advantages over the face mask	Physicians	35	p < 0.0001	32.7	p < 0.02
	Midwives	21	p < 0.0001	55.3	
Advantages over the tracheal tube	Physicians	43	p < 0.0001	53.1	p < 0.08
	Midwives	16	p < 0.0001	76.2	
Disadvantages	Physicians	15	p < 0.0001	27.3	p < 0.26
	Midwives	8	p = 0.08	42.1	
Potential applications	Physicians	30	p < 0.0001	54.5	p < 0.006
	Midwives	14	p = 0.001	93.3	

Facial mask, FM; tracheal tube, TT

[§] Improvement (n) = a pre-test wrong answer coupled with a post-test correct answer by McNemar's test. *Improvement (%) = improvement number over pre-test wrong answer number ratio by Fischer's test. p value (p < 0.05).

The practical part of the test showed that the physicians and midwives were skillful in their attempts at LMA insertion. One physician and one midwife failed, respectively, 3 and 2 times to insert the device correctly. All the midwives and physicians, with the exception of

one, expressed a high degree of approval with regard to neonatal resuscitation by LMA and defined it a sustainable and cost-effective procedure. Table 3

Table 3 Practical trial in inserting LMA in a neonatal manikin

LMA::	Physicians: n. 21	Midwives: n. 7
Successful placement:		
- number of attempts:		
1	20	6
2	0	1
3	1	0
Insertion time:		
- positioning < 15 seconds		
	20	6
- positioning > 15 seconds		
	1	1
Degree of approval:		
- high		
	20	7
- low		
	1	0
Sustainability:		
- yes		
	20	7
- no		
	1	0
Cost-effectiveness:		
- yes		
	20	7
- no		
	1	0

Discussion

Neonatal mortality, amounting to an estimated 4 million deaths worldwide each year takes place, in 98% of cases, in developing countries. As approximately 19% of these deaths are due to birth asphyxia [1], identifying solutions to achieve the Millennium Development Goal of halving child mortality by 2015 by means of a wide-scale implementation of cost-effective interventions has become urgent [17].

In this study, we proved the capability of learning and properly using in a manikin model LMA by physicians and midwives, trained in a developing country to a NRP course and to a workshop on LMA. The knowledge gained by the physicians related to the LMA was superior than that achieved by the midwives. Manikin-based comparative practical skills on LMA insertion showed instead, a similar high efficacy between trained physicians and midwives, in terms of number of successful attempts and of time for placement. Unanimously, midwives and physicians, except one, also manifested an high degree of approval of the neonatal resuscitation by LMA, giving a positive evaluation of his/her sustainability and cost-effectiveness in a low income country.

Some studies have demonstrated that NRP courses are effective in teaching neonatal resuscitation in developing countries [18] and have indicated approaches and

techniques that have proved efficacious in preliminary trials in saving newborn lives [14,19]. Birth attendants in India [20] and China [21] involved in the NRP have been able to reduce asphyxia-related deaths. There is also evidence that mouth-to-mask and bag-and-mask resuscitation are comparable techniques with regards to neonatal mortality and morbidity [4,22]. The evidence that room air [23,24] and mouth-to-mask could be satisfactory used for neonatal resuscitation [25] suggests that firstly, it is reasonable to promote the basic elements of resuscitation of newborn resuscitation, as routine part of newborn care. Meanwhile, where feasible, all birth attendants should be trained to provide positive ventilation with other conventional methods currently used in neonatal resuscitation, bag-and-mask or bag-and-LMA ventilation.

The LMA may theoretically offer many practical (i.e. easier transport and sterilization), cost-effectiveness, and sustainable advantages (i.e. low cost coupled with the fact that it can be reused) over the face mask. As mentioned above, LMA allows a low pressure airtight seal against the glottis [7-9] and studies on the efficacy of ventilation by medical and paramedical personnel in neonatal training models have shown that the LMA combines ease of insertion and adequate, rapid airway patency [11].

The AHA recommends bag-and-mask ventilation, a challenging procedure for those inexperienced in neonatal resuscitation when a newborn requires PPV, while tracheal intubation may be impossible due to lack of skill or the presence of severe congenital abnormalities. Some case reports have, moreover, shown the successful use of the LMA in resuscitation of newborns with congenital airway abnormality under inadequate ventilation and difficult intubation settings [26,27]. Tests on neonatal intubation training models have shown that midwives and interns can obtain a clear airway more rapidly with LMA than TT and with fewer failures with LMA than with TT [28].

While our study demonstrates that LMA can be easily taught to local healthcare workers in developing countries, it does have some limitations. The theoretical knowledge gained and the manual skills involved in manipulating LMA and a manikin concern only a scant handful of birth attendants and were never verified in real asphyxiated neonates in the delivery room or in other birth settings. The cost-effectiveness and sustainability of resuscitation by LMA in countries where resources are extremely limited require further scientific scrutiny, just as it would be useful to analyze the retention of the knowledge and skills gained by these and other birth attendants attending specialized courses.

Conclusions

In view of the minimum amount of time and resources necessary to train the participants in NRP courses and its many advantages with respect to FM and TT, LMA

should be considered for further evaluation and use in developing countries.

Additional material

Additional file 1: Neonatal Resuscitation Program (NRP). Questions on neonatal Laryngeal Mask Airway. Questions concerning: (I) features of LMA; (II) advantages of LMA over the face mask; (III) advantages of LMA over the endotracheal tube; (IV) disadvantages of LMA; and (V) potential applications in neonatal resuscitation.

Acknowledgements

None

Author details

¹Department of Pediatrics, Padua University, (3, Via Giustiniani) Padua (35128), Italy. ²Department of Pharmacology and Anesthesiology, Padua University, (3, Via Giustiniani) Padua (35128), Italy.

Authors' contributions

VZ had the original idea for strategy of analysis in collaboration with DT and AS. MM retrieved the data. LT reviewed the literature. FC did the statistical analysis. VZ wrote the paper. All authors read and approved the final manuscript.

Competing interests

The authors declare that they have no competing interests.

Received: 11 August 2009 Accepted: 31 August 2010

Published: 31 August 2010

References

1. Bhutta ZA, Darmstadt GL, Hasan BS, Haws RA: **Community-based interventions for improving perinatal and neonatal health outcomes in developing countries: a review of the evidence.** *Pediatrics* 2005, **115**:519-617.
2. The International Liaison Committee on Resuscitation (ILCOR) consensus on science with treatment recommendations for pediatric and neonatal patients: neonatal resuscitation. *Pediatrics* 2006, **117**:e978-e988.
3. Bloom RS, Cropley C: **Textbook of Neonatal Resuscitation.** Elk Grove Village, IL. American Heart Association, American Academy of Pediatrics, 4 2000.
4. Massawe A, Kilewo C, Irani S, Verna RJ, Chakrapam AB, Ribbe T, et al: **Assessment of mouth-to-mask ventilation in resuscitation of asphyxial newborn babies. A pilot study.** *Trop Med Int Health* 1996, **1**:865-873.
5. Trevisanuto D, Ibrahim SA, Doglioni N, Salvadori S, Ferrarese P, Zanardo V: **Neonatal resuscitation courses for pediatric residents: comparison between Khartoum (Sudan) and Padova (Italy).** *Paediatr Anaesth* 2007, **17**:28-31.
6. Grein AJ, Weiner GM: **Laryngeal mask airway versus bag-mask ventilation or endotracheal intubation for neonatal resuscitation [protocol].** *Cochrane Library Document* 2002.
7. Brain AJ, Vergese C, Addy EV: **The intubating laryngeal mask. I. Development of a new device for intubation of the trachea.** *Br J Anaesth* 1997, **79**:699-703.
8. Brain AJ, Vergese C, Addy EV, Kapila A, Brimacombe J: **The intubating laryngeal mask II: a preliminary clinical report of a new means of intubating the trachea.** *Br J Anaesth* 1997, **79**:704-709.
9. Harnett M, Kinirorns B, Heffernan A, Motherway C, Casey W: **Airway complications in infants: comparison of laryngeal mask airway and the facemask-oral airway.** *Can J Anaesth* 2000, **47**:315-318.
10. Mora EU, Weiner GM: **Alternative Ventilation Strategies: Laryngeal Masks.** *Clin Perinatol* 2006, **33**:99-110.
11. Micaglio M, Doglioni N, Parotto M, Zanardo V, Ori C, Trevisanuto D: **Training for neonatal resuscitation with the laryngeal mask airway: a comparison of the LMA-ProSeal and the LMA-Classical in an airway management manikin.** *Paediatr Anaesth* 2006, **16**:1028-31.

12. Office of Population Research: **World fertility surveys**. Princeton, NJ: Office of Population Research, Princeton University 1973 [<http://opr.princeton.edu/archive/wfs/>], Accessed December 28, 2004.
13. Deorari AK, Paul VK, Singh M, Vidyasagar D: **Impact of education and training on neonatal resuscitation practices in 14 teaching hospitals in India**. *Ann Trop Paediatr* 2001, **21**:29-33.
14. Ergenekon E, Koc E, Atalay Y, Soysal S: **Neonatal resuscitation course experience in Turkey**. *Resuscitation* 2000, **45**:225-227.
15. Trevisanuto D, Ferrarese P, Cavicchioli P, Fasson A, Zanardo V, Zacchello F: **Knowledge gained by pediatric residents after neonatal resuscitation program courses**. *Pediatr Anesth* 2005, **15**:944-947.
16. Martin PD, Cyna AM, Hunter WAH, Henry J, Ramayya GP: **Training nursing staff in airway management for resuscitation: a clinical comparison of the face mask and laryngeal mask**. *Anaesthesia* 1993, **18**:33-37.
17. Bhutta ZA, Gupta I, deSilva H, Manandhar D, Awasthi S, Hossain SM, Salam MA: **Maternal and child health: is South Asia ready for change?** *BMJ* 2004, **328**:816-819.
18. Marsh D, Darmstadt G, Moore J, Daly P, Oot D, Tinker A: **Advancing newborn health and survival in developing countries: a conceptual framework**. *J Perinatol* 2002, **22**:572-576.
19. Darmstadt G, Lawn J, Costello A: **Advancing the state of the world's newborns**. *Bull World Health Organ* 2003, **81**:224-225.
20. Deorari A, Paul V, Singh M, Vidyasagar D: **The national movement of neonatal resuscitation in India**. *J Trop Pediatr* 2000, **46**:315-317.
21. Zhu X, Fang H, Zeng S, Li Y, Lin H, Shi S: **The impact of the neonatal resuscitation program guidelines (NRPG) on the neonatal mortality in a hospital in Zhuhai, China**. *Singapore Med J* 1997, **38**:485-487.
22. Kamenir S: **Neonatal resuscitation and newborn outcomes in rural Kenya**. *J Trop Pediatr* 1997, **43**:170-173.
23. Saugstad O, Rootwelt T, Aalen O: **Resuscitation of asphyxiated newborn infants with room air or oxygen: an international controlled trial: the Resair 2 study**. *Pediatrics* 1998, **102**(1).
24. Ramji S, Ahuja S, Thirupuram S, Rootwelt T, Rooth G, Saugstad O: **Resuscitation of asphyxiated newborn infants with room air or 100% oxygen**. *Pediatr Res* 1993, **34**:809-812.
25. Kamenir S: **Neonatal resuscitation and newborn outcomes in rural Kenya**. *J Trop Pediatr* 1997, **43**:170-173.
26. Mawer RJ: **Equipment for pediatric resuscitation**. *Anaesthesia* 1995, **50**(1):87-88.
27. Baraka A: **Larynx mask airway for resuscitation of a newborn with Pierre-Robin syndrome**. *Anesthesiology* 1995, **80**:1248-1253.
28. Lavies NG: **Use of the laryngeal mask airway in neonatal resuscitation**. *Anaesthesia* 1993, **48**:352.

Pre-publication history

The pre-publication history for this paper can be accessed here:
<http://www.biomedcentral.com/1472-6963/10/254/prepub>

doi:10.1186/1472-6963-10-254

Cite this article as: Zanardo et al.: Laryngeal Mask Airway for neonatal resuscitation in a developing country: evaluation of an educational intervention. Neonatal LMA: an educational intervention in DRC. *BMC Health Services Research* 2010 **10**:254.

**Submit your next manuscript to BioMed Central
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

