

RESEARCH ARTICLE

Open Access

Eigenvalues of complementary Lidstone boundary value problems

Ravi P Agarwal^{1,2*} and Patricia JY Wong³

* Correspondence:

Agarwal@tamuk.edu
¹Department of Mathematics, Texas
A&M University - Kingsville,
Kingsville, TX 78363, USA
Full list of author information is
available at the end of the article**Abstract**

We consider the following complementary Lidstone boundary value problem

$$\begin{aligned} (-1)^m y^{(2m+1)}(t) &= \lambda F(t, y(t), y'(t)), \quad t \in (0, 1) \\ y(0) = 0, \quad y^{(2k-1)}(0) &= y^{(2k-1)}(1) = 0, \quad 1 \leq k \leq m \end{aligned}$$

where $\lambda > 0$. The values of λ are characterized so that the boundary value problem has a positive solution. Moreover, we derive explicit intervals of λ such that for any λ in the interval, the existence of a positive solution of the boundary value problem is guaranteed. Some examples are also included to illustrate the results obtained. Note that the nonlinear term F depends on y' and this derivative dependence is seldom investigated in the literature.

AMS Subject Classification: 34B15.**Keywords:** eigenvalues, positive solutions, complementary Lidstone boundary value problems**1 Introduction**In this article, we shall consider the *complementary Lidstone* boundary value problem

$$\begin{aligned} (-1)^m y^{(2m+1)}(t) &= \lambda F(t, y(t), y'(t)), \quad t \in (0, 1) \\ y(0) = 0, \quad y^{(2k-1)}(0) &= y^{(2k-1)}(1) = 0, \quad 1 \leq k \leq m \end{aligned} \quad (1.1)$$

where $m \geq 1$, $\lambda > 0$, and F is continuous at least in the interior of the domain of interest. Note that the nonlinear term F involves a derivative of the dependent variable—this is seldom studied in the literature and most research articles on boundary value problems consider nonlinear terms that involve y only.

We are interested in the existence of a positive solution of (1.1). By a *positive solution* y of (1.1), we mean a nontrivial $y \in C^{(2m+1)}(0, 1)$ satisfying (1.1) and $y(t) \geq 0$ for $t \in (0, 1)$. If, for a particular λ the boundary value problem (1.1) has a positive solution y , then λ is called an *eigenvalue* and y is a corresponding *eigenfunction* of (1.1). We shall denote the set of eigenvalues of (1.1) by E , i.e.,

$$E = \{\lambda > 0 \mid (1.1) \text{ has a positive solution}\}.$$

The focus of this article is *eigenvalue problem*, as such we shall characterize the values of λ so that the boundary value problem (1.1) has a positive solution. To be specific, we shall establish criteria for E to contain an interval, and for E to be an interval

(which may either be bounded or unbounded). In addition explicit subintervals of E are derived.

The *complementary Lidstone* interpolation and boundary value problems are very recently introduced in [1], and studied by Agarwal et. al. [2,3] where they consider an odd order $((2m+ 1)$ th order) differential equation together with boundary data at the odd order derivatives

$$\gamma(0) = a_0, \quad \gamma^{(2k-1)}(0) = a_k, \quad \gamma^{(2k-1)}(1) = b_k, \quad 1 \leq k \leq m. \tag{1.2}$$

The boundary conditions (1.2) are known as *complementary Lidstone* boundary conditions, they naturally complement the *Lidstone* boundary conditions [4-7] which involve even order derivatives. To be precise, the *Lidstone* boundary value problem comprises an even order $(2m$ th order) differential equation and the *Lidstone* boundary conditions

$$\gamma^{(2k)}(0) = a_k, \quad \gamma^{(2k)}(1) = b_k, \quad 0 \leq k \leq m - 1. \tag{1.3}$$

There is a vast literature on Lidstone interpolation and boundary value problems. The Lidstone interpolation has a long history from 1929 when Lidstone [8] introduced a generalization of Taylor’s series that approximates a given function in the neighborhood of two points instead of one. Further characterization can be found in the study of [9-16]. More research on Lidstone interpolation as well as Lidstone spline is seen in [1,17-23]. On the other hand, the Lidstone boundary value problems and several of its particular cases have been the subject matter of numerous investigations, see [4,18,24-37] and the references cited therein. It is noted that in most of these studies the nonlinear terms considered do *not* involve derivatives of the dependent variable, only a handful of articles [30,31,34,35] tackle nonlinear terms that involve even order derivatives. In the present study, our study of the *complementary Lidstone* boundary value problem (1.1) where F depends on a *derivative* certainly extends and complements the rich literature on boundary value problems and in particular on Lidstone boundary value problems.

The plan of the article is as follows. In Section 2, we shall state a fixed point theorem due to Krasnosel’skii [38], and develop some inequalities for certain Green’s function which are needed later. The characterization of the set E is presented in Section 3. Finally, in Section 4, we establish explicit subintervals of E .

2 Preliminaries

Theorem 2.1. [38] Let B be a Banach space, and let $C(\subset B)$ be a cone. Assume Ω_1, Ω_2 are open subsets of B with $0 \in \Omega_1, \bar{\Omega}_1 \subset \Omega_2$, and let $S : C \cap (\bar{\Omega}_2 \setminus \Omega_1) \rightarrow C$ be a completely continuous operator such that, either

- (a) $\|Sy\| \leq \|y\|, y \in C \cap \partial\Omega_1$, and $\|Sy\| \geq \|y\|, y \in C \cap \partial\Omega_2$, or
- (b) $\|Sy\| \geq \|y\|, y \in C \cap \partial\Omega_1$, and $\|Sy\| \leq \|y\|, y \in C \cap \partial\Omega_2$.

Then, S has a fixed point in $C \cap (\bar{\Omega}_2 \setminus \Omega_1)$.

To tackle the *complementary Lidstone* boundary value problem (1.1), let us review certain attributes of the *Lidstone* boundary value problem. Let $g_m(t, s)$ be the Green’s function of the Lidstone boundary value problem

$$\begin{aligned} x^{(2m)}(t) &= 0, \quad t \in (0, 1) \\ x^{(2k)}(0) &= x^{(2k)}(1) = 0, \quad 0 \leq k \leq m - 1. \end{aligned} \tag{2.1}$$

The Green's function $g_m(t, s)$ can be expressed as [4,5]

$$g_m(t, s) = \int_0^1 g(t, u)g_{m-1}(u, s)du \tag{2.2}$$

where

$$g_1(t, s) = g(t, s) = \begin{cases} t(s - 1), & 0 \leq t \leq s \leq 1 \\ s(t - 1), & 0 \leq s \leq t \leq 1. \end{cases} \tag{2.3}$$

Further, it is known that

$$|g_m(t, s)| = (-1)^m g_m(t, s) \quad \text{and} \quad g_m(t, s) = g_m(s, t), \quad (t, s) \in (0, 1) \times (0, 1). \tag{2.4}$$

We also have the inequality

$$\frac{2}{\pi^2} \sin \pi t \leq t(1 - t) \leq \frac{1}{\pi} \sin \pi t, \quad t \in [0, 1]. \tag{2.5}$$

The following two lemmas give the upper and lower bounds of $|g_m(t, s)|$, they play an important role in subsequent development.

Lemma 2.1. For $(t, s) \in [0, 1] \times [0, 1]$, we have

$$|g_m(t, s)| \leq \frac{1}{\pi^{2m-1}} \sin \pi s. \tag{2.6}$$

Proof. For $(t, s) \in [0, 1] \times [0, 1]$, it is clear from (2.3) that

$$|g(t, s)| \leq s(1 - s). \tag{2.7}$$

Using (2.7), (2.4), and (2.5) in (2.2) yields for $(t, s) \in [0, 1] \times [0, 1]$,

$$\begin{aligned} |g_m(t, s)| &= \int_0^1 |g(t, u)| \cdot |g_{m-1}(u, s)| du \leq \int_0^1 |g_{m-1}(u, s)| u(1 - u) du \\ &\leq \frac{1}{\pi} \int_0^1 |g_{m-1}(s, u)| \sin \pi u du. \end{aligned} \tag{2.8}$$

By induction, we can show that

$$\int_0^1 |g_m(t, s)| \sin \pi s ds = \frac{1}{\pi^{2m}} \sin \pi t, \quad t \in [0, 1]. \tag{2.9}$$

Now (2.6) is immediate by applying (2.9) to (2.8).

Lemma 2.2. Let $\delta \in \left(0, \frac{1}{2}\right)$ be given. For $(t, s) \in [\delta, 1-\delta] \times [0, 1]$, we have

$$|g_m(t, s)| \geq \frac{2\delta}{\pi^{2m}} \sin \pi s. \tag{2.10}$$

Proof. For $(t, s) \in [\delta, 1-\delta] \times [0, 1]$, from (2.3) we find

$$|g(t, s)| \geq \begin{cases} \delta(1-s), & 1 \leq s \\ [1 - (1-\delta)]s, & s \leq t \end{cases} \geq \delta s(1-s). \tag{2.11}$$

Then, using (2.11), (2.4), and (2.5) in (2.2), we get for $(t, s) \in [\delta, 1-\delta] \times [0, 1]$,

$$\begin{aligned} |g_m(t, s)| &= \int_0^1 |g(t, u)| \cdot |g_{m-1}(u, s)| \, du \geq \delta \int_0^1 |g_{m-1}(u, s)| u(1-u) \, du \\ &\geq \frac{2\delta}{\pi^2} \int_0^1 |g_{m-1}(s, u)| \sin \pi u \, du, \end{aligned}$$

which, in view of (2.9), gives (2.10) immediately.

Remark 2.1. The bounds in Lemmas 2.1 and 2.2 are *sharper* than those given in the literature [4,5,35,37].

3 Eigenvalues of (1.1)

To tackle (1.1) we first consider the initial value problem

$$\begin{aligned} \gamma'(t) &= x(t), \quad t \in (0, 1) \\ \gamma(0) &= 0 \end{aligned} \tag{3.1}$$

whose solution is simply

$$\gamma(t) = \int_0^t x(s) \, ds. \tag{3.2}$$

Taking into account (3.1) and (3.2), the *complementary Lidstone* boundary value problem (1.1) reduces to the *Lidstone* boundary value problem

$$\begin{aligned} (-1)^m x^{(2m)}(t) &= \lambda F \left(t, \int_0^t x(s) \, ds, x(t) \right), \quad t \in (0, 1) \\ x^{(2k-2)}(0) &= x^{(2k-2)}(1) = 0, \quad 1 \leq k \leq m. \end{aligned} \tag{3.3}$$

If (3.3) has a positive solution x^* , then by virtue of (3.2), $\gamma^*(t) = \int_0^t x^*(s) \, ds$ is a positive solution of (1.1). Hence, the existence of a positive solution of the *complementary Lidstone* boundary value problem (1.1) follows from the existence of a positive solution of the *Lidstone* boundary value problem (3.3). It is clear that an eigenvalue of (3.3) is also an eigenvalue of (1.1), thus

$$E = \{\lambda > 0 | (1.1) \text{ has a positive solution}\} = \{\lambda > 0 | (3.3) \text{ has a positive solution}\}.$$

With the lemmas developed in Section 2 and a technique to handle the nonlinear term F , we shall study the eigenvalue problem (1.1) via (3.3).

For easy reference, we list below the conditions that are used later. In these conditions, f , α , and β are continuous functions with $f : (0, \infty) \times (0, \infty) \rightarrow (0, \infty)$ and $\alpha, \beta : (0, 1) \rightarrow [0, \infty)$.

(A1) f is nondecreasing in each of its arguments, i.e., for $u, u_1, u_2, v, v_1, v_2 \in (0, \infty)$ with $u_1 \leq u_2$ and $v_1 \leq v_2$, we have

$$f(u_1, v) \leq f(u_2, v) \quad \text{and} \quad f(u, v_1) \leq f(u, v_2);$$

(A2) for $t \in (0, 1)$ and $u, v \in (0, \infty)$,

$$\alpha(t)f(u, v) \leq F(t, u, v) \leq \beta(t)f(u, v);$$

(A3) $\alpha(t)$ is not identically zero on any nondegenerate subinterval of $(0, 1)$ and there exists $a_0 \in (0, 1]$ such that $\alpha(t) \geq a_0\beta(t)$ for all $t \in (0, 1)$;

$$(A4) \quad 0 < \int_0^1 \beta(t) \sin \pi t dt < \infty;$$

(A5) for $t \in (0, 1)$ and $u, u_1, u_2, v, v_1, v_2 \in (0, \infty)$ with $u_1 \leq u_2$ and $v_1 \leq v_2$, we have

$$F(t, u_1, v) \leq F(t, u_2, v) \quad \text{and} \quad F(t, u, v_1) \leq F(t, u, v_2).$$

We shall consider the Banach space $B = C[0, 1]$ equipped with the norm

$$\|x\| = \sup_{t \in [0, 1]} |x(t)|, \quad x \in B.$$

For a given $\delta \in \left(0, \frac{1}{2}\right)$, let the cone C_δ be defined by

$$C_\delta = \left\{ x \in B \mid x(t) \geq 0, \quad t \in [0, 1]; \quad \min_{t \in [\delta, 1-\delta]} x(t) \geq \gamma \|x\| \right\}$$

where $\gamma = \frac{2\delta}{\pi} a_0$ (a_0 is defined in (A3)). Further, let

$$C_\delta(M) = \{x \in C_\delta \mid \|x\| \leq M\}.$$

Let the operator $S : C_\delta \rightarrow B$ be defined by

$$\begin{aligned} Sx(t) &= \lambda \int_0^1 (-1)^m g_m(t, s) F \left(s, \int_0^s x(\tau) d\tau, x(s) \right) ds \\ &= \lambda \int_0^1 |g_m(t, s)| F \left(s, \int_0^s x(\tau) d\tau, x(s) \right) ds, \quad t \in [0, 1]. \end{aligned} \tag{3.4}$$

To obtain a positive solution of (3.3), we shall seek a fixed point of the operator S in the cone C_δ .

Further, we define the operators $U, V : C_\delta \rightarrow B$ by

$$Ux(t) = \lambda \int_0^1 |g_m(t, s)| \alpha(s) f \left(\int_0^s x(\tau) d\tau, x(s) \right) ds$$

and

$$Vx(t) = \lambda \int_0^1 |g_m(t, s)| \beta(s) f \left(\int_0^s x(\tau) d\tau, x(s) \right) ds.$$

If (A2) holds, then

$$Ux(t) \leq Sx(t) \leq Vx(t), \quad t \in [0, 1]. \tag{3.5}$$

Lemma 3.1. Let (A1)-(A4) hold. Then, the operator S is compact on the cone C_δ .

Proof. Let us consider the case when $\alpha(t)$ is unbounded in a deleted right neighborhood of 0 and also in a deleted left neighborhood of 1. Clearly, $\beta(t)$ is also unbounded near 0 and 1.

For $n \in \{1, 2, 3, \dots\}$, let $\alpha_n, \beta_n : [0, 1] \rightarrow [0, \infty)$ be defined by

$$\alpha_n(t) = \begin{cases} \alpha\left(\frac{1}{n+1}\right), & 0 \leq t \leq \frac{1}{n+1} \\ \alpha(t), & \frac{1}{n+1} \leq t \leq \frac{n}{n+1} \\ \alpha\left(\frac{n}{n+1}\right), & \frac{n}{n+1} \leq t \leq 1 \end{cases}$$

and

$$\beta_n(t) = \begin{cases} \beta\left(\frac{1}{n+1}\right), & 0 \leq t \leq \frac{1}{n+1} \\ \beta(t), & \frac{1}{n+1} \leq t \leq \frac{n}{n+1} \\ \beta\left(\frac{n}{n+1}\right), & \frac{n}{n+1} \leq t \leq 1. \end{cases}$$

Also, we define the operators $U_n, V_n : C_\delta \rightarrow B$ by

$$U_n x(t) = \lambda \int_0^1 |g_m(t, s)| \alpha_n(s) f\left(\int_0^s x(\tau) d\tau, x(s)\right) ds$$

and

$$V_n x(t) = \lambda \int_0^1 |g_m(t, s)| \beta_n(s) f\left(\int_0^s x(\tau) d\tau, x(s)\right) ds.$$

It is standard that for each n , both U_n and V_n are compact operators on C_δ . Let $M > 0$ and $x \in C_\delta(M)$. For $t \in [0, 1]$, we get

$$\begin{aligned} |V_n x(t) - Vx(t)| &\leq \int_0^1 |g_m(t, s)| |\beta_n(s) - \beta(s)| f\left(\int_0^s x(\tau) d\tau, x(s)\right) ds \\ &= \frac{1}{n+1} \int_0^1 |g_m(t, s)| |\beta_n(s) - \beta(s)| f\left(\int_0^s x(\tau) d\tau, x(s)\right) ds \\ &\quad + \frac{1}{n} \int_0^1 |g_m(t, s)| |\beta_n(s) - \beta(s)| f\left(\int_0^s x(\tau) d\tau, x(s)\right) ds. \end{aligned}$$

By the monotonicity of f (see (A1)), we have

$$f\left(\int_0^s x(\tau) d\tau, x(s)\right) \leq f\left(\int_0^1 M d\tau, M\right) = f(M, M) \tag{3.6}$$

Coupling with Lemma 2.1, it follows that

$$|V_n x(t) - Vx(t)| \leq f(M, M) \left[\int_0^{\frac{1}{n+1}} \frac{1}{\pi^{2m-1}} \left| \beta\left(\frac{1}{n+1}\right) - \beta(s) \right| \sin \pi s ds + \int_{\frac{n}{n+1}}^1 \frac{1}{\pi^{2m-1}} \left| \beta\left(\frac{n}{n+1}\right) - \beta(s) \right| \sin \pi s ds \right].$$

The integrability of $\beta(t) \sin \pi t$ (see (A4)) ensures that V_n converges uniformly to V on $C_\delta(M)$. Hence, V is compact on C_δ . By a similar argument, we see that U_n converges uniformly to U on $C_\delta(M)$ and therefore U is also compact on C_δ . It follows immediately from inequality (3.5) that the operator S is compact on C_δ .

Remark 3.1. From the proof of Lemma 3.1, we see that if the functions α and β are continuous on the close interval $[0, 1]$, then the conditions (A1) and (A4) are *not* needed in Lemma 3.1.

The first result shows that E contains an interval.

Theorem 3.1. Let (A1)-(A4) hold. Then, there exists $c > 0$ such that the interval $(0, c] \subseteq E$.

Proof. Let $M > 0$ be given. Define

$$c = \frac{M}{f(M, M)} \left[\int_0^1 \frac{1}{\pi^{2m-1}} \beta(s) \sin \pi s ds \right]^{-1}. \tag{3.7}$$

Let $\lambda \in (0, c]$. We shall prove that $S(C_\delta(M)) \subseteq C_\delta(M)$. Let $x \in C_\delta(M)$. First, we shall show that $Sx \in C_\delta$. It is clear from (3.5) that

$$Sx(t) \geq \lambda \int_0^1 |g_m(t, s)| \alpha(s) f\left(\int_0^s x(\tau) d\tau, x(s)\right) ds \geq 0, \quad t \in [0, 1]. \tag{3.8}$$

Further, from (3.5) and Lemma 2.1 we get

$$\begin{aligned} Sx(t) &\leq \lambda \int_0^1 |g_m(t, s)| \beta(s) f\left(\int_0^s x(\tau) d\tau, x(s)\right) ds \\ &\leq \lambda \int_0^1 \frac{1}{\pi^{2m-1}} \beta(s) f\left(\int_0^s x(\tau) d\tau, x(s)\right) \sin \pi s ds, \quad t \in [0, 1] \end{aligned}$$

which leads to

$$\|Sx\| \leq \lambda \int_0^1 \frac{1}{\pi^{2m-1}} \beta(s) f\left(\int_0^s x(\tau) d\tau, x(s)\right) \sin \pi s ds. \tag{3.9}$$

Now, applying (3.5), Lemma 2.2, (A3) and (3.9) successively, we find for $t \in [\delta, 1-\delta]$,

$$\begin{aligned} Sx(t) &\geq \lambda \int_0^1 |g_m(t,s)| \alpha(s) f \left(\int_0^s x(\tau) d\tau, x(s) \right) ds \\ &\geq \lambda \int_0^1 \frac{2\delta}{\pi^{2m}} \alpha(s) f \left(\int_0^s x(\tau) d\tau, x(s) \right) \sin \pi s ds \\ &\geq \lambda \int_0^1 \frac{2\delta}{\pi^{2m}} a_0 \beta(s) f \left(\int_0^s x(\tau) d\tau, x(s) \right) \sin \pi s ds \\ &\geq \frac{2\delta}{\pi} a_0 \|Sx\| = \gamma \|Sx\|. \end{aligned}$$

Therefore,

$$\min_{t \in [\delta, 1-\delta]} Sx(t) \geq \gamma \|Sx\|. \tag{3.10}$$

Inequalities (3.8) and (3.10) imply that $Sx \in C_\delta$.

Next, we shall verify that $\|Sx\| \leq M$. For this, an application of (3.5), Lemma 2.1, (3.6) and (3.7) provides

$$\begin{aligned} Sx(t) &\leq c \int_0^1 |g_m(t,s)| \beta(s) f \left(\int_0^s x(\tau) d\tau, x(s) \right) ds \\ &\leq cf(M, M) \int_0^1 \frac{1}{\pi^{2m-1}} \beta(s) \sin \pi s ds = M, \quad t \in [0, 1] \end{aligned}$$

or equivalently

$$\|Sx\| \leq M.$$

Hence, $S(C_\delta(M)) \subseteq C_\delta(M)$. Also, the standard arguments yield that S is completely continuous. By Schauder fixed point theorem, S has a fixed point in $C_\delta(M)$. Clearly, this fixed point is a positive solution of (3.3) and therefore λ is an eigenvalue of (3.3). Since $\lambda \in (0, c]$ is arbitrary, it follows immediately that the interval $(0, c] \subseteq E$.

Remark 3.2. From the proof of Theorem 3.1, we see that (A2) and (A3) lead to $S : C_\delta \rightarrow C_\delta$.

Theorem 3.2. Let (A1)-(A5) hold. Suppose that $\lambda^* \in E$, for any $\lambda \in (0, \lambda^*)$, we have $\lambda \in E$, i.e., $(0, \lambda^*] \subseteq E$.

Proof. Let x^* be the eigenfunction corresponding to the eigenvalue λ^* . Thus, we have

$$x^*(t) = Sx^*(t) = \lambda^* \int_0^1 |g_m(t,s)| F \left(s, \int_0^s x^*(\tau) d\tau, x^*(s) \right) ds, \quad t \in [0, 1]. \tag{3.11}$$

Define

$$K^* = \{x \in B \mid 0 \leq x(t) \leq x^*(t), \quad t \in [0, 1]\}.$$

Let $\lambda \in (0, \lambda^*)$ and $x \in K^*$. Using (A5), we get

$$\begin{aligned} 0 \leq Sx(t) &= \lambda \int_0^1 |g_m(t, s)| F \left(s, \int_0^s x(\tau) d\tau, x(s) \right) ds \\ &\leq \lambda^* \int_0^1 |g_m(t, s)| F \left(s, \int_0^s x^*(\tau) d\tau, x^*(s) \right) ds \\ &= Sx^*(t), \quad t \in [0, 1] \end{aligned}$$

where the last equality follows from (3.11). This immediately implies that the operator S maps K^* into K^* . Moreover, the operator S is continuous and completely continuous. Schauder's fixed point theorem guarantees that S has a fixed point in K^* , which is a positive solution of (3.3). Hence, λ is an eigenvalue, i.e., $\lambda \in E$.

The following result shows that E is an interval.

Corollary 3.1. Let (A1)-(A5) hold. If $E \neq \emptyset$, then E is an interval.

Proof. Suppose E is not an interval. Then, there exist $\lambda_0, \lambda'_0 \in E (\lambda_0 < \lambda'_0)$ and $\tau \in (\lambda_0, \lambda'_0)$ with $\tau \notin E$. However, this is not possible as Theorem 3.2 guarantees that $\tau \in E$. Hence, E is an interval.

The following two results give the upper and lower bounds of an eigenvalue in terms of some parameters of the corresponding eigenfunction.

Theorem 3.3. Let (A1) and (A2) hold. Assume that m is odd. Let λ be an eigenvalue of (3.3) and $x \in C_\delta$ be a corresponding eigenfunction. If $x^{(i)}(0) = b_i, i = 1, 3, \dots, 2m - 1$, where $b_{2m-1} > 0$, then λ satisfies

$$M_1 \leq \lambda \leq M_2 \tag{3.12}$$

where

$$\begin{aligned} M_1 &= \max_{0 \leq k \leq m-1} \left[\sum_{i=k}^{m-1} \frac{b_{2i+1}}{(2(i-k)+1)!} \right] \left[f(D, D) \int_0^1 \frac{(1-s)^{2(m-k)-1}}{(2(m-k)-1)!} \beta(s) ds \right]^{-1}, \\ M_2 &= \min_{0 \leq k \leq m-1} \left[\sum_{i=k}^{m-1} \frac{b_{2i+1}}{(2(i-k)+1)!} \right] \left[f(0, 0) \int_0^1 \frac{(1-s)^{2(m-k)-1}}{(2(m-k)-1)!} \alpha(s) ds \right]^{-1} \end{aligned}$$

and

$$D = \max_{t \in [0, 1]} \sum_{i=0}^{m-1} b_{2i+1} \frac{t^{2i+1}}{(2i+1)!}.$$

Proof. For $n \in \{1, 2, 3, \dots\}$, we define $f_n = f^* \omega_n$, where ω_n is a standard mollifier [25] such that f_n is Lipschitz and converges uniformly to f .

For a fixed n , let λ_n be an eigenvalue and $x_n(t)$, with $x_n^{(i)}(0) = b_i, i = 1, 3, \dots, 2m - 1$ be a corresponding eigenfunction of the following boundary value problem

$$(-1)^m x_n^{(2m)}(t) = \lambda_n F_n \left(t, \int_0^t x_n(s) ds, x_n(t) \right), \quad t \in [0, 1] \tag{3.13}$$

$$x_n^{(2i)}(0) = x_n^{(2i)}(1) = 0, \quad 0 \leq i \leq m - 1 \tag{3.14}$$

where F_n converges uniformly to F , and for $u, v \in (0, \infty)$,

$$\alpha_n(t)f_n(u, v) \leq F_n(t, u, v) \leq \beta_n(t)f_n(u, v), \quad t \in (0, 1) \tag{3.15}$$

(see the proof of Lemma 3.1 for the definitions of $\alpha_n(t)$ and $\beta_n(t)$).

It is clear that $x_n(t)$ is the unique solution of the initial value problem (3.13),

$$\begin{aligned} x_n^{(i)}(0) &= 0, \quad i = 0, 2, \dots, 2m - 2 \\ x_n^{(i)}(0) &= b_i, \quad i = 1, 3, \dots, 2m - 1. \end{aligned} \tag{3.16}$$

First, we shall establish an upper bound for x_n . Since

$$(-1)^m x_n^{(2m)}(t) = \lambda_n F_n \left(t, \int_0^t x_n(s) ds, x_n(t) \right) \geq \lambda_n \alpha_n(t) f_n \left(\int_0^t x_n(s) ds, x_n(t) \right) \geq 0,$$

we have $x_n^{(2m-1)}(t)$ is nonincreasing and hence

$$x_n^{(2m-1)}(t) \leq x_n^{(2m-1)}(0) = b_{2m-1}, \quad t \in [0, 1]. \tag{3.17}$$

In view of the initial conditions (3.16) and also (3.17), we find

$$x_n^{(2m-2)}(t) = \int_0^t x_n^{(2m-1)}(s) ds \leq b_{2m-1} t, \quad t \in [0, 1]. \tag{3.18}$$

Next, an application of (3.18) gives

$$x_n^{(2m-3)}(t) = b_{2m-3} + \int_0^t x_n^{(2m-2)}(s) ds \leq b_{2m-3} + b_{2m-1} \frac{t^2}{2!}, \quad t \in [0, 1].$$

By repeating the process, we get

$$x_n(t) \leq b_1 t + b_3 \frac{t^3}{3!} + \dots + b_{2m-1} \frac{t^{2m-1}}{(2m-1)!} \leq D, \quad t \in [0, 1]. \tag{3.19}$$

By the monotonicity of f_n , we have

$$f_n \left(\int_0^t x_n(s) ds, x_n(t) \right) \leq f_n \left(\int_0^1 D ds, D \right) = f_n(D, D)$$

and

$$f_n \left(\int_0^t x_n(s) ds, x_n(t) \right) \geq f_n \left(\int_0^t 0 ds, 0 \right) = f_n(0, 0).$$

Coupling with (3.13) and (3.15), it follows that

$$\lambda_n \alpha_n(t) f_n(0, 0) \leq (-1)^m x_n^{(2m)}(t) \leq \lambda_n \beta_n(t) f_n(D, D), \quad t \in [0, 1]. \tag{3.20}$$

Once again, using the initial conditions (3.16), repeated integration of (3.20) from 0 to t provides

$$\phi_{1,k}(t) \leq x_n^{(2k)}(t) \leq \phi_{2,k}(t), \quad t \in [0, 1], \quad 0 \leq k \leq m - 1 \tag{3.21}$$

where

$$\phi_{1,k}(t) = \sum_{i=k}^{m-1} b_{2i+1} \frac{t^{2(i-k)+1}}{(2(i-k)+1)!} - \lambda_n f_n(D, D) \int_0^t \frac{(t-s)^{2(m-k)-1}}{(2(m-k)-1)!} \beta_n(s) ds$$

and

$$\phi_{2,k}(t) = \sum_{i=k}^{m-1} b_{2i+1} \frac{t^{2(i-k)+1}}{(2(i-k)+1)!} - \lambda_n f_n(0, 0) \int_0^t \frac{(t-s)^{2(m-k)-1}}{(2(m-k)-1)!} \alpha_n(s) ds.$$

In order to satisfy the boundary conditions $x_n^{(2k)}(1) = 0, \quad 0 \leq k \leq m - 1$, from inequality (3.21) it is necessary that

$$\phi_{1,k}(1) \leq 0 \quad \text{and} \quad \phi_{2,k}(1) \geq 0, \quad 0 \leq k \leq m - 1.$$

This readily implies

$$M_{1,n} \leq \lambda_n \leq M_{2,n} \tag{3.22}$$

where

$$M_{1,n} = \max_{0 \leq k \leq m-1} \left[\sum_{i=k}^{m-1} \frac{b_{2i+1}}{(2(i-k)+1)!} \right] \left[f_n(D, D) \int_0^1 \frac{(1-s)^{2(m-k)-1}}{(2(m-k)-1)!} \beta_n(s) ds \right]^{-1}$$

and

$$M_{2,n} = \min_{0 \leq k \leq m-1} \left[\sum_{i=k}^{m-1} \frac{b_{2i+1}}{(2(i-k)+1)!} \right] \left[f_n(0, 0) \int_0^1 \frac{(1-s)^{2(m-k)-1}}{(2(m-k)-1)!} \alpha_n(s) ds \right]^{-1}.$$

From (3.20) it is observed (by using the initial conditions (3.16) and repeated integration) that $\{x_n^{(i)}\}_{n=1}^\infty, \quad 0 \leq i \leq 2m - 1$ is a uniformly bounded sequence on $[0, 1]$. Thus, there exists a subsequence, which can be relabeled as $\{x_n\}_{n=1}^\infty$, that converges uniformly (in fact, in $C^{(2m-1)}$ -norm) to some x on $[0, 1]$. We note that each $x_n(t)$ can be expressed as

$$x_n(t) = \lambda_n \int_0^1 |g_m(t, s)| F_n \left(s, \int_0^s x_n(\tau) d\tau, x_n(s) \right) ds, \quad t \in [0, 1]. \tag{3.23}$$

Since $\{\lambda_n\}_{n=1}^\infty$ is a bounded sequence (from (3.22)), there is a subsequence, which can be relabeled as $\{\lambda_n\}_{n=1}^\infty$, that converges to some λ . Then, letting $n \rightarrow \infty$ in (3.23) yields

$$x(t) = \lambda \int_0^1 |g_m(t, s)| F \left(s, \int_0^s x(\tau) d\tau, x(s) \right) ds, \quad t \in [0, 1].$$

This means that $x(t)$ is an eigenfunction of (3.3) corresponding to the eigenvalue λ . Further, $x^{(i)}(0) = b_i, \quad i = 1, 3, \dots, 2m - 1$ and inequality (3.12) follows from (3.22) immediately.

Theorem 3.4. Let (A1)-(A4) hold. Let λ be an eigenvalue of (3.3) and $x \in C_\delta$ be a corresponding eigenfunction. Further, let $\|x\| = p$. Then,

$$\lambda \geq \frac{p}{f(p,p)} \left[\frac{1}{\pi^{2m-1}} \int_0^1 \beta(s) \sin \pi s ds \right]^{-1} \tag{3.24}$$

and

$$\lambda \leq \frac{p}{f(\gamma p(\frac{1}{2} - \delta), \gamma p)} \left[\int_{\frac{1}{2}}^{1-\delta} |g_m(t_1, s)| \alpha(s) ds \right]^{-1} \tag{3.25}$$

where t_1 is any number in $(0, 1)$ such that $x(t_1) \neq 0$.

Proof. Let $t_0 \in [0, 1]$ be such that

$$p = \|x\| = x(t_0).$$

Then, using (3.5), Lemma 2.1 and the monotonicity of f , we find

$$\begin{aligned} p = x(t_0) = Sx(t_0) &\leq \lambda \int_0^1 |g_m(t_0, s)| \beta(s) f \left(\int_0^s x(\tau) d\tau, x(s) \right) ds \\ &\leq \lambda \int_0^1 \frac{1}{\pi^{2m-1}} \beta(s) f \left(\int_0^1 p d\tau, p \right) \sin \pi s ds \\ &= \lambda \frac{f(p,p)}{\pi^{2m-1}} \int_0^1 \beta(s) \sin \pi s ds \end{aligned}$$

which gives (3.24) readily.

Next, we employ (3.5), the monotonicity of f and the fact that $\min_{t \in [\delta, 1-\delta]} x(t) \geq \gamma p$ to get

$$\begin{aligned} p &\geq x(t_1) \geq \lambda \int_0^1 |g_m(t_1, s)| \alpha(s) f \left(\int_0^s x(\tau) d\tau, x(s) \right) ds \\ &\geq \lambda \int_{\frac{1}{2}}^{1-\delta} |g_m(t_1, s)| \alpha(s) f \left(\int_0^s x(\tau) d\tau, x(s) \right) ds \\ &\geq \lambda \int_{\frac{1}{2}}^{1-\delta} |g_m(t_1, s)| \alpha(s) f \left(\int_0^{\frac{1}{2}} x(\tau) d\tau, x(s) \right) ds \\ &\geq \lambda \int_{\frac{1}{2}}^{1-\delta} |g_m(t_1, s)| \alpha(s) f \left(\int_\delta^{\frac{1}{2}} x(\tau) d\tau, x(s) \right) ds \\ &\geq \lambda \int_{\frac{1}{2}}^{1-\delta} |g_m(t_1, s)| \alpha(s) f \left(\int_\delta^{\frac{1}{2}} \gamma p d\tau, \gamma p \right) ds \\ &= \lambda f \left(\gamma p \left(\frac{1}{2} - \delta \right), \gamma p \right) \int_{\frac{1}{2}}^{1-\delta} |g_m(t_1, s)| \alpha(s) ds \end{aligned}$$

from which (3.25) is immediate.

The following result gives the criteria for E to be a bounded/unbounded interval.

Theorem 3.5. Define

$$\begin{aligned}
 W_B &= \left\{ f \mid \frac{u}{f(u, u)} \text{ is bounded for } u \in (0, \infty) \right\}, \\
 W_0 &= \left\{ f \mid \lim_{u \rightarrow \infty} \frac{u}{f(u, u)} = 0 \right\}, \\
 W_\infty &= \left\{ f \mid \lim_{u \rightarrow \infty} \frac{u}{f(u, u)} = \infty \right\}.
 \end{aligned}$$

- (a) Let (A1)-(A5) hold. If $f \in W_B$, then $E = (0, c)$ or $(0, c]$ for some $c \in (0, \infty)$.
- (b) Let (A1)-(A5) hold. If $f \in W_0$, then $E = (0, c]$ for some $c \in (0, \infty)$.
- (c) Let (A1)-(A4) hold. If $f \in W_\infty$, then $E = (0, \infty)$.

Proof. (a) This is immediate from (3.25) and Corollary 3.1.

- (b) Since $W_0 \subseteq W_B$, it follows from Case (a) that $E = (0, c)$ or $(0, c]$ for some $c \in (0, \infty)$.

In particular,

$$c = \sup E.$$

Let $\{\lambda_n\}_{n=1}^\infty$ be a monotonically increasing sequence in E which converges to c , and let $\{x_n\}_{n=1}^\infty$ be a corresponding sequence of eigenfunctions in the context of (3.3). Further, let $p_n = \|x_n\|$. Then, (3.25) together with $f \in W_0$ implies that no subsequence of $\{p_n\}_{n=1}^\infty$ can diverge to infinity. Thus, there exists $R > 0$ such that $p_n \leq R$ for all n . So $\{x_n\}_{n=1}^\infty$ is uniformly bounded. This implies that there is a subsequence of $\{x_n\}_{n=1}^\infty$, relabeled as the original sequence, which converges uniformly to some x , where $x(t) \geq 0$ for $t \in [0, 1]$. Clearly, we have $Sx_n = x_n$, i.e.,

$$x_n(t) = \lambda_n \int_0^1 |g_m(t, s)| F \left(s, \int_0^s x_n(\tau) d\tau, x_n(s) \right) ds, \quad t \in [0, 1]. \tag{3.26}$$

Since x_n converges to x and λ_n converges to c , letting $n \rightarrow \infty$ in (3.26) yields

$$x(t) = c \int_0^1 |g_m(t, s)| F \left(s, \int_0^s x(\tau) d\tau, x(s) \right) ds, \quad t \in [0, 1].$$

Hence, c is an eigenvalue with corresponding eigenfunction x , i.e., $c = \sup E \in E$. This completes the proof for Case (b).

- (c) Let $\lambda > 0$ be fixed. Choose $\varepsilon > 0$ so that

$$\frac{\lambda}{\pi^{2m-1}} \int_0^1 \beta(s) \sin \pi s ds \leq \frac{1}{\varepsilon}. \tag{3.27}$$

By definition, if $f \in W_\infty$, then there exists $M = M(\varepsilon) > 0$ such that

$$f(u, u) < \varepsilon u, \quad u \geq M. \tag{3.28}$$

We shall prove that $S(C_\delta(M)) \subseteq C_\delta(M)$. Let $x \in C_\delta(M)$. As in the proof of Theorem 3.1, we have (3.8) and (3.10) and so $Sx \in C_\delta$. Thus, it remains to show that $\|Sx\| \leq M$. Using (3.5), Lemma 2.1, (3.6), (3.28), and (3.27), we find for $t \in [0, 1]$,

$$\begin{aligned} Sx(t) &\leq \lambda \int_0^1 |g_m(t, s)| \beta(s) f \left(\int_0^s x(\tau) d\tau, x(s) \right) ds \\ &\leq \lambda f(M, M) \int_0^1 \frac{1}{\pi^{2m-1}} \beta(s) \sin \pi s ds \\ &\leq \lambda \varepsilon M \int_0^1 \frac{1}{\pi^{2m-1}} \beta(s) \sin \pi s ds \leq M. \end{aligned}$$

It follows that $\|Sx\| \leq M$ and hence $S(C_\delta(M)) \subseteq C_\delta(M)$. Also, S is continuous and completely continuous. Schauder's fixed point theorem guarantees that S has a fixed point in $C_\delta(M)$. Clearly, this fixed point is a positive solution of (3.3) and therefore λ is an eigenvalue of (3.3). Since $\lambda > 0$ is arbitrary, we have proved that $E = (0, \infty)$.

Example 3.1. Consider the complementary Lidstone boundary value problem

$$\begin{aligned} \gamma^{(5)} &= \lambda \left(\frac{t^5}{10} + \frac{t^4}{4} - t^3 + \frac{t^2}{4} + \frac{t}{2} + 2 \right)^{-q} \left[\frac{\gamma + \gamma'}{2} + 2 \right]^q, \quad t \in (0, 1) \\ \gamma(0) &= \gamma'(0) = \gamma'''(0) = \gamma'(1) = \gamma'''(1) = 0 \end{aligned} \tag{3.29}$$

where $\lambda > 0$ and $q \geq 0$.

Here, $m = 2$ and

$$F(t, \gamma, \gamma') = \left(\frac{t^5}{10} + \frac{t^4}{4} - t^3 + \frac{t^2}{4} + \frac{t}{2} + 2 \right)^{-q} \left(\frac{\gamma + \gamma'}{2} + 2 \right)^q.$$

Clearly, $F(t, u, v)$ is nondecreasing in u and v , thus (A5) is satisfied.

Choose

$$\alpha(t) = \beta(t) = \left(\frac{t^5}{10} + \frac{t^4}{4} - t^3 + \frac{t^2}{4} + \frac{t}{2} + 2 \right)^{-q}$$

and

$$f(u, v) = \left(\frac{u + v}{2} + 2 \right)^q.$$

We see that (A1)-(A4) are satisfied.

Case 1. $0 \leq q < 1$. Clearly $f \in W_\infty$. It follows from Theorem 3.5(c) that the set $E = (0, \infty)$. As an example, when $\lambda = 24$, the boundary value problem (3.29) has a positive solution given by $\gamma(t) = \frac{t^5}{5} - \frac{t^4}{2} + \frac{t^2}{2}$.

Case 2. $q = 1$. Here $f \in W_B$. By Theorem 3.5(a) the set E is an open or a half-closed interval. Further, from Case 1 and Theorem 3.2 we note that E contains the interval $(0, 24]$.

Case 3. $q > 1$. Clearly $f \in W_0$. By Theorem 3.5(b) the set E is a half-closed interval. Again, as in Case 2 we note that $(0, 24] \subseteq E$.

4 Eigenvalue intervals

In this section, we shall establish explicit subintervals of E . Here, the functions α and β in (A2)-(A4) are assumed to be continuous on the closed interval $[0, 1]$. Hence, noting Remark 3.1, we shall not require conditions (A1) and (A4) to show the compactness of the operator S . For the function f in (A2), we define

$$\begin{aligned} \bar{f}_0 &= \limsup_{u \rightarrow 0, v \rightarrow 0} \frac{f(u, v)}{v}, & f_{-0} &= \liminf_{u \rightarrow 0, v \rightarrow 0} \frac{f(u, v)}{v}, \\ \bar{f}_\infty &= \limsup_{u \rightarrow \infty, v \rightarrow \infty} \frac{f(u, v)}{v}, & f_{-\infty} &= \liminf_{u \rightarrow \infty, v \rightarrow \infty} \frac{f(u, v)}{v}. \end{aligned}$$

Let $\delta \in (0, \frac{1}{2})$ be given. Define $t^*, \hat{t} \in [0, 1]$ by

$$\begin{aligned} \int_{\frac{1}{2}}^{1-\delta} |g_m(t^*, s)| \alpha(s) ds &= \sup_{t \in [0, 1]} \int_{\frac{1}{2}}^{1-\delta} |g_m(t, s)| \alpha(s) ds, \\ \int_{\delta}^{1-\delta} |g_m(\hat{t}, s)| \alpha(s) ds &= \sup_{t \in [0, 1]} \int_{\delta}^{1-\delta} |g_m(t, s)| \alpha(s) ds, \end{aligned} \tag{4.1}$$

Theorem 4.1. Let (A2)-(A4) hold. Then, $\lambda \in E$ if λ satisfies

$$\frac{1}{f_{-\infty}} \left[\gamma \int_{\frac{1}{2}}^{1-\delta} |g_m(t^*, s)| \alpha(s) ds \right]^{-1} < \lambda < \frac{1}{f_0} \left[\frac{1}{\pi^{2m-1}} \int_0^1 \beta(s) \sin \pi s ds \right]^{-1}. \tag{4.2}$$

Proof. We shall use Theorem 2.1. Let λ satisfy (4.2) and let $\varepsilon > 0$ be such that

$$\frac{1}{f_{-\infty} - \varepsilon} \left[\gamma \int_{\frac{1}{2}}^{1-\delta} |g_m(t^*, s)| \alpha(s) ds \right]^{-1} \leq \lambda \leq \frac{1}{f_0 + \varepsilon} \left[\frac{1}{\pi^{2m-1}} \int_0^1 \beta(s) \sin \pi s ds \right]^{-1} \tag{4.3}$$

First, we pick $p > 0$ so that

$$f(u, v) \leq (\bar{f}_0 + \varepsilon)v, \quad 0 < u \leq p, \quad 0 < v \leq p. \tag{4.4}$$

Let $x \in C_\delta$ be such that $\|x\| = p$. Note that for $s \in [0, 1]$,

$$\int_0^s x(\tau) d\tau \leq \int_0^1 p d\tau = p.$$

Then, using (3.5), Lemma 2.1, (4.4) and (4.3) successively, we find for $t \in [0, 1]$,

$$\begin{aligned} Sx(t) &\leq \lambda \int_0^1 \frac{1}{\pi^{2m-1}} \beta(s) f \left(\int_0^s x(\tau) d\tau, x(s) \right) \sin \pi s ds \\ &\leq \lambda \int_0^1 \frac{1}{\pi^{2m-1}} \beta(s) (\bar{f}_0 + \varepsilon) x(s) \sin \pi s ds \\ &\leq \lambda \int_0^1 \frac{1}{\pi^{2m-1}} \beta(s) (\bar{f}_0 + \varepsilon) \|x\| \sin \pi s ds \leq \|x\|. \end{aligned}$$

Hence,

$$\|Sx\| \leq \|x\|. \tag{4.5}$$

If we set $\Omega_1 = \{x \in B \mid \|x\| < p\}$, then (4.5) holds for $x \in C_\delta \cap \partial \Omega_1$.

Next, let $q > 0$ be such that

$$f(u, v) \geq (f_{-\infty} - \varepsilon)v, \quad u \geq q, \quad v \geq q. \tag{4.6}$$

Let $x \in C_\delta$ be such that

$$\|x\| = \max \left\{ p + 1, \frac{q}{\gamma}, \frac{q}{\gamma} \left(\frac{1}{2} - \delta \right)^{-1} \right\} = \max \left\{ p + 1, \frac{q}{\gamma} \left(\frac{1}{2} - \delta \right)^{-1} \right\} \equiv q_0.$$

It is clear that

$$\begin{cases} x(s) \geq \gamma \|x\| \geq q, & s \in \left[\frac{1}{2}, 1 - \delta \right] \\ \int_{\frac{1}{2}}^{\frac{1}{2}} x(\tau) d\tau \geq \int_{\frac{1}{2}}^{\frac{1}{2}} \gamma \|x\| d\tau = \left(\frac{1}{2} - \delta \right) \gamma \|x\| \geq q. \end{cases} \tag{4.7}$$

Then, an application of (3.5), (4.7), and (4.6) gives for $t \in [0, 1]$,

$$\begin{aligned} Sx(t) &\geq \lambda \int_{\frac{1}{2}}^{1-\delta} |g_m(t, s)| \alpha(s) f \left(\int_0^s x(\tau) d\tau, x(s) \right) ds \\ &\geq \lambda \int_{\frac{1}{2}}^{1-\delta} |g_m(t, s)| \alpha(s) f \left(\int_{\frac{1}{2}}^{\frac{1}{2}} x(\tau) d\tau, x(s) \right) ds \\ &\geq \lambda \int_{\frac{1}{2}}^{1-\delta} |g_m(t, s)| \alpha(s) (f_{-\infty} - \varepsilon) x(s) ds \\ &\geq \lambda \int_{\frac{1}{2}}^{1-\delta} |g_m(t, s)| \alpha(s) (f_{-\infty} - \varepsilon) \gamma \|x\| ds. \end{aligned}$$

Taking supremum both sides and using (4.3) then provides (see (4.1) for the definition of t^*)

$$\|Sx\| \geq \lambda \int_{\frac{1}{2}}^{1-\delta} |g_m(t^*, s)| \alpha(s) ds \cdot (f_{-\infty} - \varepsilon) \gamma \|x\| \geq \|x\|.$$

Therefore, if we set $\Omega_2 = \{x \in B \mid \|x\| < q_0\}$, then for $x \in C_\delta \cap \partial \Omega_2$ we have

$$\|Sx\| \geq \|x\|. \tag{4.8}$$

Now that we have obtained (4.5) and (4.8), it follows from Remark 3.2 and Theorem 2.1 that S has a fixed point $x \in C_\delta \cap (\bar{\Omega}_2 \setminus \Omega_1)$ such that $p \leq \|x\| \leq q_0$. Obviously, this x is a positive solution of (3.3) and hence $\lambda \in E$. \square

Theorem 4.2. Let (A2)-(A4) hold. Then, $\lambda \in E$ if λ satisfies

$$\frac{1}{f_{-0}} \left[\gamma \int_{\delta}^{1-\delta} |g_m(\hat{t}, s)| \alpha(s) ds \right]^{-1} < \lambda < \frac{1}{f_{\infty}} \left[\frac{1}{\pi^{2m-1}} \int_0^1 \beta(s) \sin \pi s ds \right]^{-1}. \tag{4.9}$$

Proof. We shall apply Theorem 2.1 again. Let λ satisfy (4.9) and let $\varepsilon > 0$ be such that

$$\frac{1}{f_{-0} - \varepsilon} \left[\gamma \int_{\delta}^{1-\delta} |g_m(\hat{t}, s)| \alpha(s) ds \right]^{-1} \leq \lambda \leq \frac{1}{f_{\infty} + \varepsilon} \left[\frac{1}{\pi^{2m-1}} \int_0^1 \beta(s) \sin \pi s ds \right]^{-1} \tag{4.10}$$

First, we choose $r > 0$ so that

$$f(u, v) \geq (f_{-0} - \varepsilon)v, \quad 0 < u \leq r, \quad 0 < v \leq r. \tag{4.11}$$

Let $x \in C_\delta$ be such that $\|x\| = r$. Then, on using (3.5), (4.11), and (4.10) successively, we have for $t \in [0, 1]$,

$$\begin{aligned} Sx(t) &\geq \lambda \int_0^1 |g_m(t, s)| \alpha(s) f \left(\int_0^s x(\tau) d\tau, x(s) \right) ds \\ &\geq \lambda \int_0^1 |g_m(t, s)| \alpha(s) (f_{-0} - \varepsilon) x(s) ds \\ &\geq \lambda \int_{\delta}^{1-\delta} |g_m(t, s)| \alpha(s) (f_{-0} - \varepsilon) \gamma \|x\| ds. \end{aligned}$$

Taking supremum both sides and using (4.10) then yields (see (4.1) for the definition of \hat{t})

$$\|Sx\| \geq \lambda \int_{\delta}^{1-\delta} |g_m(\hat{t}, s)| \alpha(s) ds \cdot (f_{-0} - \varepsilon) \gamma \|x\| \geq \|x\|.$$

Hence, if we set $\Omega_1 = \{y \in B \mid \|y\| < r\}$, then (4.8) holds for $x \in C_\delta \cap \partial \Omega_1$.

Next, pick $w > 0$ such that

$$f(u, v) \leq (f_{\infty} + \varepsilon)v, \quad u \geq w, \quad v \geq w. \tag{4.12}$$

We shall consider two cases - when f is bounded and when f is unbounded.

Case 1. Suppose that f is bounded. Then, there exists some $M > 0$ such that

$$f(u, v) \leq M, \quad u, v \in (0, \infty). \tag{4.13}$$

Let $x \in C_\delta$ be such that

$$\|x\| = \max \left\{ r + 1, \frac{\lambda M}{\pi^{2m-1}} \int_0^1 \beta(s) \sin \pi s ds \right\} \equiv w_0.$$

From (3.5), Lemma 2.1 and (4.13), it is clear for $t \in [0, 1]$ that

$$\begin{aligned} Sx(t) &\leq \lambda \int_0^1 \frac{1}{\pi^{2m-1}} \beta(s) f \left(\int_0^s x(\tau) d\tau, x(s) \right) \sin \pi s ds \\ &\leq \lambda \int_0^1 \frac{1}{\pi^{2m-1}} \beta(s) M \sin \pi s ds \leq w_0 = \|x\|. \end{aligned}$$

Hence, (4.5) holds.

Case 2. Suppose that f is unbounded. Then, there exists $w_0 > \max \{r + 1, w\}$ such that

$$f(u, v) \leq f(w_0, w_0), \quad 0 < u \leq w_0, \quad 0 < v \leq w_0. \tag{4.14}$$

Let $x \in C_\delta$ be such that $\|x\| = w_0$. Then, applying (3.5), Lemma 2.1, (4.14), (4.12), and (4.10) successively gives for $t \in [0, 1]$,

$$\begin{aligned} Sx(t) &\leq \lambda \int_0^1 \frac{1}{\pi^{2m-1}} \beta(s) f \left(\int_0^s x(\tau) d\tau, x(s) \right) \sin \pi s ds \\ &\leq \lambda \int_0^1 \frac{1}{\pi^{2m-1}} \beta(s) f(w_0, w_0) \sin \pi s ds \\ &\leq \lambda \int_0^1 \frac{1}{\pi^{2m-1}} \beta(s) (\bar{f}_\infty + \varepsilon) w_0 \sin \pi s ds \\ &= \lambda \int_0^1 \frac{1}{\pi^{2m-1}} \beta(s) (\bar{f}_\infty + \varepsilon) \|x\| \sin \pi s ds \leq \|x\|. \end{aligned}$$

Thus, (4.5) follows immediately.

In both Cases 1 and 2, if we set $\Omega_2 = \{x \in B \mid \|x\| < w_0\}$, then (4.5) holds for $x \in C_\delta \cap \partial\Omega_2$.

Now that we have obtained (4.8) and (4.5), it follows from Remark 3.2 and Theorem 2.1 that S has a fixed point $x \in C_\delta \cap (\bar{\Omega}_2 \setminus \Omega_1)$ such that $r \leq \|x\| \leq w_0$. It is clear that this x is a positive solution of (3.3) and hence $\lambda \in E$.

Remark 4.1. In (4.2) and (4.9), although t^* and \hat{t} can be computed from (4.1), we can circumvent the computation by giving further bounds. Indeed, applying Lemma 2.2 we find

$$\int_{\frac{1}{2}}^{1-\delta} |g_m(t^*, s)| \alpha(s) ds \geq \sup_{t \in [\delta, 1-\delta]} \int_{\frac{1}{2}}^{1-\delta} |g_m(t, s)| \alpha(s) ds \geq \frac{2\delta}{\pi^{2m}} \int_{\frac{1}{2}}^{1-\delta} \alpha(s) \sin \pi s ds$$

and

$$\int_{\delta}^{1-\delta} |g_m(\hat{t}, s)| \alpha(s) ds \geq \sup_{t \in [\delta, 1-\delta]} \int_{\delta}^{1-\delta} |g_m(t, s)| \alpha(s) ds \geq \frac{2\delta}{\pi^{2m}} \int_{\delta}^{1-\delta} \alpha(s) \sin \pi s ds.$$

The following corollary is immediate from Theorems 4.1, 4.2 and Remark 4.1.

Corollary 4.1. Let (A2)-(A4) hold. Then,

$$\begin{aligned} E &\supseteq \left(\frac{1}{f_{-\infty}} \left[\gamma \int_{\frac{1}{2}}^{1-\delta} |g_m(t^*, s)| \alpha(s) ds \right]^{-1}, \frac{1}{\bar{f}_0} \left[\frac{1}{\pi^{2m-1}} \int_0^1 \beta(s) \sin \pi s ds \right]^{-1} \right) \\ &\supseteq \left(\frac{1}{f_{-\infty}} \left[\frac{2\gamma\delta}{\pi^{2m}} \int_{\frac{1}{2}}^{1-\delta} \alpha(s) \sin \pi s ds \right]^{-1}, \frac{1}{\bar{f}_0} \left[\frac{1}{\pi^{2m-1}} \int_0^1 \beta(s) \sin \pi s ds \right]^{-1} \right) \end{aligned}$$

and

$$\begin{aligned} E &\supseteq \left(\frac{1}{f_{-0}} \left[\gamma \int_{\delta}^{1-\delta} |g_m(\hat{t}, s)| \alpha(s) ds \right]^{-1}, \frac{1}{\bar{f}_{\infty}} \left[\frac{1}{\pi^{2m-1}} \int_0^1 \beta(s) \sin \pi s ds \right]^{-1} \right) \\ &\supseteq \left(\frac{1}{f_{-0}} \left[\frac{2\gamma\delta}{\pi^{2m}} \int_{\delta}^{1-\delta} \alpha(s) \sin \pi s ds \right]^{-1}, \frac{1}{\bar{f}_{\infty}} \left[\frac{1}{\pi^{2m-1}} \int_0^1 \beta(s) \sin \pi s ds \right]^{-1} \right). \end{aligned}$$

Remark 4.2. If f is superlinear (i.e., $\bar{f}_0 = 0$ and $f_{-\infty} = \infty$) or sublinear (i.e., $f_{-0} = \infty$ and $\bar{f}_{\infty} = 0$), then we conclude from Corollary 4.1 that $E = (0, \infty)$, i.e., the boundary value problem (3.3) (or (1.1)) has a positive solution for any $\lambda > 0$.

Example 4.1. Consider the complementary Lidstone boundary value problem

$$\begin{aligned} \gamma^{(5)} &= \lambda \left[a \left(t^5 - \frac{5t^4}{2} + \frac{5t^2}{2} \right) + b(5t^4 - 10t^3 + 5t) + c \right]^{-r} (a\gamma + b\gamma' + c)^r, \quad t \in (0, 1) \\ \gamma(0) &= \gamma'(0) = \gamma'''(0) = \gamma'(1) = \gamma'''(1) = 0 \end{aligned} \tag{4.15}$$

where $\lambda, a, b, c > 0$ and $r \leq 1$.

Here, $m = 2$. It is clear that (A2)-(A4) are satisfied with

$$\alpha(t) = \beta(t) = \left[a \left(t^5 - \frac{5t^4}{2} + \frac{5t^2}{2} \right) + b(5t^4 - 10t^3 + 5t) + c \right]^{-r} \quad \text{and} \quad f(u, v) = (au + bv + c)^r.$$

Case 1. $r < 1$. It is clear that f is sublinear. Therefore, by Remark 4.2 the boundary value problem (4.15) has a positive solution for any $\lambda > 0$. In fact, we note that when $\lambda = 120$, (4.15) has a positive solution given by $x(t) = t^5 - \frac{5t^4}{2} + \frac{5t^2}{2}$.

Case 2. $r = 1$, $a = b = 0.5$, $c = 10$. Here, $f_{-0} = \infty$ and $\bar{f}_{\infty} = 1$. It follows from Corollary 4.1 that

$$E \supseteq \left(0, \frac{\pi^3}{\bar{f}_{\infty}} \left[\int_0^1 \beta(s) \sin \pi s ds \right]^{-1} \right) = (0, 528.99).$$

Once again we note that when $\lambda = 120 \in (0, 528.99)$, the corresponding eigenfunction is given by $x(t) = t^5 - \frac{5t^4}{2} + \frac{5t^2}{2}$.

Author details

¹Department of Mathematics, Texas A&M University - Kingsville, Kingsville, TX 78363, USA ²Department of Mathematics, Faculty of Science, King Abdulaziz University, Jeddah 21589, Saudi Arabia ³School of Electrical and Electronic Engineering, Nanyang Technological University, 50 Nanyang Avenue, Singapore 639798, Singapore

Authors' contributions

All authors read and approved the final manuscript.

Competing interests

The authors declare that they have no competing interests.

Received: 9 December 2011 Accepted: 24 April 2012 Published: 24 April 2012

References

1. Costabile, FA, Dell'Accio, F, Luceri, R: Explicit polynomial expansions of regular real functions by means of even order Bernoulli polynomials and boundary values. *J Comput Appl Math.* **175**, 77–99 (2005). doi:10.1016/j.cam.2004.06.006
2. Agarwal, RP, Pinelas, S, Wong, PJY: Complementary Lidstone interpolation and boundary value problems. *J Inequal Appl.* **2009**, 1–30 (2009)
3. Agarwal, RP, Wong, PJY: Piecewise complementary Lidstone interpolation and error inequalities. *J Comput Appl Math.* **234**, 2543–2561 (2010). doi:10.1016/j.cam.2010.03.029
4. Agarwal, RP, Wong, PJY: Lidstone polynomials and boundary value problems. *Comput Math Appl.* **17**, 1397–1421 (1989). doi:10.1016/0898-1221(89)90023-0
5. Agarwal, RP, Wong, PJY: *Error Inequalities in Polynomial Interpolation and their Applications.* Kluwer, Dordrecht (1993)
6. Davis, PJ: *Interpolation and Approximation.* Blaisdell Publishing Co., Boston (1961)
7. Varma, AK, Howell, G: Best error bounds for derivatives in two point Birkhoff interpolation problem. *J Approx Theory.* **38**, 258–268 (1983). doi:10.1016/0021-9045(83)90132-6
8. Lidstone, GJ: Notes on the extension of Aitken's theorem (for polynomial interpolation) to the Everett types. *Proc Edinburgh Math Soc.* **2**, 16–19 (1929)
9. Boas, RP: A note on functions of exponential type. *Bull Am Math Soc.* **47**, 750–754 (1941). doi:10.1090/S0002-9904-1941-07552-X
10. Boas, RP: Representation of functions by Lidstone series. *Duke Math J.* **10**, 239–245 (1943). doi:10.1215/S0012-7094-43-01021-X
11. Poritsky, H: On certain polynomial and other approximations to analytic functions. *Trans Am Math Soc.* **34**, 274–331 (1932). doi:10.1090/S0002-9947-1932-1501639-4
12. Schoenberg, IJ: On certain two-point expansions of integral functions of exponential type. *Bull Am Math Soc.* **42**, 284–288 (1936). doi:10.1090/S0002-9904-1936-06293-2
13. Whittaker, JM: On Lidstone's series and two-point expansions of analytic functions. *Proc Lond Math Soc.* **36**, 451–469. (1933-34)
14. Whittaker, JM: *Interpolatory Function Theory.* Cambridge University Press, Cambridge (1935)
15. Widder, DV: Functions whose even derivatives have a prescribed sign. *Proc Nat Acad Sci.* **26**, 657–659 (1940). doi:10.1073/pnas.26.11.657
16. Widder, DV: Completely convex functions and Lidstone series. *Trans Am Math Soc.* **51**, 387–398 (1942)
17. Agarwal, RP, Wong, PJY: Explicit error bounds for the derivatives of piecewise-Lidstone interpolation. *J Comp Appl Math.* **58**, 67–81 (1995). doi:10.1016/0377-0427(93)E0262-K
18. Agarwal, RP, Wong, PJY: Error bounds for the derivatives of Lidstone interpolation and applications, in: In: Govil NK (ed.) *Approximation Theory, in memory of A.K. Varma.* pp. 1–41. Marcal Dekker, New York (1998)
19. Costabile, FA, Dell'Accio, F: Lidstone approximation on the triangle. *Appl Numer Math.* **52**, 339–361 (2005). doi:10.1016/j.apnum.2004.08.003
20. Costabile, FA, Serpe, A: An algebraic approach to Lidstone polynomials. *Appl Math Lett.* **20**, 387–390 (2007). doi:10.1016/j.aml.2006.02.034
21. Wong, PJY: On Lidstone splines and some of their applications. *Neural Parallel Sci Comput.* **1**, 472–475 (1995)
22. Wong, PJY, Agarwal, RP: Sharp error bounds for the derivatives of Lidstone-spline interpolation. *Comput Math Appl.* **28**(9), 23–53 (1994). doi:10.1016/0898-1221(94)00176-6
23. Wong, PJY, Agarwal, RP: Sharp error bounds for the derivatives of Lidstone-spline interpolation II. *Comput Math Appl.* **31**(3), 61–90 (1996). doi:10.1016/0898-1221(95)00206-5

24. Agarwal, RP, Akrivis, G: Boundary value problems occurring in plate deflection theory. *J Comput Appl Math.* **8**, 145–154 (1982). doi:10.1016/0771-050X(82)90035-3
25. Agarwal, RP, O'Regan, D, Wong, PJY: *Positive Solutions of Differential, Difference and Integral Equations.* Kluwer, Dordrecht (1999)
26. Agarwal, RP, Wong, PJY: Quasilinearization and approximate quasilinearization for Lidstone boundary value problems. *Int J Comput Math.* **42**, 99–116 (1992). doi:10.1080/00207169208804054
27. Baldwin, P: Asymptotic estimates of the eigenvalues of a sixth-order boundary-value problem obtained by using global phase-integral method. *Phil Trans Roy Soc London Ser A.* **322**, 281–305 (1987). doi:10.1098/rsta.1987.0051
28. Baldwin, P: A localized instability in a Bénard layer. *Appl Anal.* **24**, 117–156 (1987). doi:10.1080/00036818708839658
29. Boutayeb, A, Twizell, EH: Finite-difference methods for twelfth-order boundary value problems. *J Comput Appl Math.* **35**, 133–138 (1991). doi:10.1016/0377-0427(91)90202-U
30. Davis, JM, Eloe, PW, Henderson, J: Triple positive solutions and dependence on higher order derivatives. *J Math Anal Appl.* **237**, 710–720 (1999). doi:10.1006/jmaa.1999.6500
31. Davis, JM, Henderson, J, Wong, PJY: General Lidstone problems: multiplicity and symmetry of solutions. *J Math Anal Appl.* **251**, 527–548 (2000). doi:10.1006/jmaa.2000.7028
32. Eloe, PW, Henderson, J, Thompson, HB: Extremal points for impulsive Lidstone boundary value problems. *Math Comput Model.* **32**, 687–698 (2000). doi:10.1016/S0895-7177(00)00165-5
33. Forster, P: Existenzaussagen und Fehlerabschätzungen bei gewissen nichtlinearen Randwertaufgaben mit gewöhnlichen Differentialgleichungen. *Numerische Mathematik.* **10**, 410–422 (1967). doi:10.1007/BF02162874
34. Guo, Y, Ge, W: Twin positive symmetric solutions for Lidstone boundary value problems. *Taiwanese J Math.* **8**, 271–283 (2004)
35. Ma, Y: Existence of positive solutions of Lidstone boundary value problems. *J Math Anal Appl.* **314**, 97–108 (2006). doi:10.1016/j.jmaa.2005.03.059
36. Twizell, EH, Boutayeb, A: Numerical methods for the solution of special and general sixth-order boundary value problems, with applications to Bénard layer eigenvalue problems. *Proc Roy Soc London Ser A.* **431**, 433–450 (1990). doi:10.1098/rspa.1990.0142
37. Yao, Q: On the positive solutions of Lidstone boundary value problems. *Appl Math Comput.* **137**, 477–485 (2003). doi:10.1016/S0096-3003(02)00152-2
38. Krasnosel'skii, MA: *Positive Solutions of Operator Equations.* Noordhoff, Groningen (1964)

doi:10.1186/1687-2770-2012-49

Cite this article as: Agarwal and Wong: Eigenvalues of complementary Lidstone boundary value problems. *Boundary Value Problems* 2012 **2012**:49.

Submit your manuscript to a SpringerOpen[®] journal and benefit from:

- ▶ Convenient online submission
- ▶ Rigorous peer review
- ▶ Immediate publication on acceptance
- ▶ Open access: articles freely available online
- ▶ High visibility within the field
- ▶ Retaining the copyright to your article

Submit your next manuscript at ▶ springeropen.com
