



# Isolation and characterization of microsatellite markers for *Sturnira parvidens* and cross-species amplification in *Sturnira* species

Edgar G. Gutiérrez<sup>1,\*</sup>, Giovanni Hernández Canchola<sup>2,\*</sup>, Livia S. León Paniagua<sup>2,\*</sup>, Norberto Martínez Méndez<sup>1,\*</sup> and Jorge Ortega<sup>1</sup>

<sup>1</sup> Department of Zoología, Instituto Politécnico Nacional/ENCB, CDMX, CDMX, México

<sup>2</sup> Department of Biología Evolutiva, Facultad de Ciencias, UNAM, CDMX, CDMX, México

\* These authors contributed equally to this work.

## ABSTRACT

**Background.** *Sturnira* is one of the most species-rich genera in the Neotropics, and it is found from Mexico and the Lesser Antilles to Argentina. This genus forms a well-supported monophyletic clade with at least twenty-one recognized species, as well as several others under taxonomic review. *Sturnira parvidens* is a widespread frugivorous bat of the deciduous forests of the Neotropics, is highly abundant, and is a major component in fruit dispersal to regenerate ecosystems.

**Methods.** We used a technique based on Illumina paired-end sequencing of a library highly enriched for microsatellite repeats to develop loci for *S. parvidens*. We analyzed millions of resulting reads with specialized software to extract those reads that contained di-, tri-, tetra-, penta-, and hexanucleotide microsatellites.

**Results.** We selected and tested 14 polymorphic (di, tri, and tetra) microsatellites. All markers were genotyped on 26 different individuals from distinct locations of the distributional area of *S. parvidens*. We observed medium—high genetic variation across most loci, but only 12 were functionally polymorphic. Levels of expected heterozygosity across all markers were high to medium (mean  $H_E = 0.79$ , mean  $H_O = 0.72$ ). We examined ascertainment bias in twelve bats of the genus, obtaining null/monomorphic/polymorphic amplifications.

**Discussion.** The Illumina paired-end sequencing system is capable of identifying massive numbers of microsatellite loci, while expending little time, reducing costs, and providing a large amount of data. The described polymorphic loci for *S. parvidens* in particular, and for the genus in general, could be suitable for further genetic analysis, including taxonomic inconsistencies, parentage/relatedness analysis, and population genetics assessments.

**Subjects** Ecology, Genetics, Molecular Biology, Zoology

**Keywords** Microsatellites, *Sturnira parvidens*, Pal\_finder, Illumina

## INTRODUCTION

The yellow-shouldered Mesoamerican bat (*Sturnira parvidens*) is primarily associated with lower elevations (0 to 2,000 m), and is found mainly in tropical/subtropical habitats and ecotones (*Villalobos & Valerio, 2002*). *S. parvidens* is found from the northern Mexican

Submitted 15 March 2017

Accepted 28 April 2017

Published 24 May 2017

Corresponding author

Jorge Ortega, [artibeus2@aol.com](mailto:artibeus2@aol.com)

Academic editor

Erica Goss

Additional Information and  
Declarations can be found on  
page 8

DOI 10.7717/peerj.3367

 Copyright

2017 Gutiérrez et al.

Distributed under  
Creative Commons CC-BY 4.0

**OPEN ACCESS**

Pacific Slope and the northern Mexican Gulf Slope southward to Northern Costa Rica, and including the Yucatan Peninsula (G Hernández-Canchola & L León-Paniagua, 2017, unpublished data). *S. parvidens* has been caught in the understory and subcanopy of tropical and subtropical forests, in xeric scrubs, and in secondary and temperate forests. They are commonly found roosting in the foliage of forests of advanced successional stages, but their home ranges include mature and secondary forest (Evelyn & Stiles, 2003). They mainly consume fruit from plants representing early stages of plant succession, like pioneer trees (*Cecropia peltata*), pioneer herbs (*Solanum americanum*, *S. torvun*, *S. ochraceo-ferrugineum*, *Capsicum annuum*), or pioneer shrubs (*Piper hispidum*, *P. lapathifolium*; *Olea-Wagner et al.*, 2007). This frugivorous species is an important seed disperser, carrying out an important ecosystemic role in the restoration of secondary tropical forests. It is considered abundant but, as fragmentation intensifies, the species is particularly vulnerable to local extinction (Evelyn & Stiles, 2003).

Pleistocene climatic oscillations and the complex orogeny of its distributional area shaped the phylogeography of this bat, generating two lowland lineages. The two genetic lineages, one in the Western Slope region of Mexico, and the other in the Eastern Slope region of Mexico and Central America, diverged into haplogroups around c. 0.423 Ma, and demographic expansion was detected later, after the splitting event (G Hernández-Canchola & L León-Paniagua, 2017, unpublished data). *Sturnira* is the most speciose genus of frugivorous bats. Due to its ability to colonize new areas, it adapted to produce complex groups showing different genetic lineages (Velazco & Patterson, 2013; Velazco & Patterson, 2014; G Hernández-Canchola & L León-Paniagua, 2017, unpublished data). The genus *Sturnira* involves a highly diversified and complex group of species. This speciose group of bats inhabits the entire Neotropic realm and includes three mountain basal species: *S. aratathomasi*, *S. bidens*, and *S. nana*. Also, it has been described as a clade formed by species that usually inhabit highland mountain forests: *S. bogotensis*, *S. burtonlimi*, *S. erythromos*, *S. hondurensis*, *S. koopmanhilli*, *S. ludovici*, *S. magna*, *S. mordax*, *S. oporaphilum*, *S. perla*, *S. tildae* and *S. adrianae* (Velazco & Patterson, 2013; Molinari et al., 2017). Lastly, it includes a group of species that inhabit lowland tropical forests: *S. angeli*, *S. bakeri*, *S. lilium*, *S. luisi*, *S. new species 3*, *S. paulsoni*, and *S. parvidens* (Velazco & Patterson, 2013).

No microsatellite molecular markers are known for *Sturnira parvidens*; our goal was to isolate and characterize polymorphic microsatellite loci for the species by using Next-Generation Sequencing. The development of these markers can be useful for understanding the genetic structure of subpopulations in its distributional range. They can be used to identify the impact of humans on the fragmentation of the populations and assess the divergent lineages formed by genetic drift. They can also be used to evaluate movements of individuals in the mosaic-fragmented landscapes, and discern the genetic component in the social structure of the population by assessing relatedness and paternity. We showed cross-species amplification in twelve species of the *Sturnira* genus, under the hypothesis of having a positive ascertainment bias due to the phylogenetic relatedness among species (Crawford et al., 1998; Li & Kimmel, 2013). Suitable cross-species amplification will facilitate studies in *Sturnira* related bat populations of Middle and South America.


## MATERIALS AND METHODS

We obtained tissue samples from 26 distinct individuals of *S. parvidens* from different localities in its distributional range in Mexico. Specimens were provided by *Colección de Mamíferos del Museo de Zoología “Alfonso L. Herrera”, Facultad de Ciencias-Universidad Nacional Autónoma de México*. Tissue samples were stored individually in 95% ethanol until analysis. We followed the guidelines set forth by the American Society of Mammalogists for the use of wildlife (*Gannon, Sikes & Animal Care and Use Committee of the American Society of Mammalogists, 2007*). Fieldwork was conducted with the permission of SEMARNAT (Secretaría del Medio Ambiente y Recursos Naturales de México—permit FAUT-0307). Six samples were sent to the Savannah River Ecology Laboratory, for an enrichment library process. The facility follows their own protocol and provides a database of the resulting microsatellites. Meanwhile the rest of the specimens were used to standardize protocols and assess polymorphism in microsatellites.

DNA was extracted following the instructions of the Qiagen protocol (Blood and Tissue Kit, Cat No. 69504; Qiagen, Hilden, Germany) for shotgun sequences, and we used the Universal Salt Protocol to extract DNA from the remaining specimens (*Aljanabi & Martinez, 1997*). An Illumina paired-end shotgun library was prepared by shearing 11 g of tissue DNA using a Covaris S220 and following the standard protocol of the Illumina TruSeq DNA Library Kit. Five million of the resulting reads were analyzed with the program PAL\_FINDER\_v0.02.03 (*Castoe et al., 2012*), in order to extract those reads that contained di-, tri-, tetra-, penta-, and hexanucleotide microsatellites.

Once positive reads were identified in PAL\_FINDER, they were batched to a local installation of the program MSATCOMMANDER v 0.8.2 for primer design (*Faircloth, 2008*). We recovered 6,790 unique loci (48 hexa, 97 penta, 1,260 tetra, 1,097 tri and 4,288 dinucleotide—[Fig. 1](#)), but only 14 were chosen for PCR trials that were performed in a MultiGene™ Gradient Thermal Cycler (Labnet, Edison, NJ, USA). We directly labelled forward primers (FAM) for each of the chosen loci. PCR reactions were performed in a 10  $\mu$ l volume containing 30 ng of DNA, 0.2 mM of dNTPs, 10 mM of each primer, 1 Taq buffer (Buffer PCR 10 $\times$ ), 0.3  $\mu$ L MgCl<sub>2</sub> (25 mM), and 1.0 U of FlexiTaq polymerase. PCR cycling conditions were as follows: initial denaturation at 95 °C for 3 min; followed by 30 cycles of 95 °C for 3 min, gradient temperature (ranging from 56 to 60 °C) for 30 s, and 72 °C for 2 min; extension of 68 °C for 8 min; and final ending of 4 °C. Exact annealing temperatures for each primer are given in [Table 1](#). We visualized the PCR products by electrophoresis on 1.5% agarose gels. Markers were tested for amplification success, polymorphism and specificity in 26 individuals of *S. parvidens*.

The results of the microsatellite profiles were examined using GeneMarker® v. 2.4.2 (SoftGenetics, State College, PA, USA) and peaks were scored by hand. We obtained the number of homozygotes and heterozygotes by scoring data. We estimated the proportion of polymorphic loci and the average number of alleles per locus by using the GDA software (*Lewis & Zaykin, 2001*). We assessed the observed ( $H_O$ ) and the expected heterozygosity ( $H_E$ ), linkage disequilibrium, and Hardy–Weinberg proportions by using Genepop 4.2 (*Rousset, 2008*), and corroborated with Arlequin 3.5 (*Excoffier, Laval & Schneider,*



**Figure 1** Potentially amplifiable loci (PAL's) with positive microsatellites found in the enriched library. Perfect, imperfect and compound loci separated out for dinucleotide to hexanucleotide microsatellite forms.

**Table 1** Primer sequences and characteristics of the 14 microsatellite loci isolated for *Sturnira parvidens*. Annealing temperature not obtained (X).

Locus	Primer (Forward) (5–3')	Primer (Reverse) (5–3')	Motif	Annealing T (°C)
Spar01	6 FAM-TGCCCTGAAGAACCTTGTGAGC	CCCATACTTCTCCCTCACAGC	AAAG(92)	58
Spar02	6 FAM-AGAAAGAAAGGGAGGGCGG	TTCTTTATGCCCTTTGCTCTAGG	AAAG(104)	60
Spar05	6 FAM-TGCCTGCCTAGTCTGTCCACC	AAGCAGTTCCCATCACATGC	ATC(33)	56
Spar06	6-FAM-CCTGGGATGAAGTTTCTGACG	GAATAATGGGAATACCAGAATAAGACG	TTC(30)	×
Spar07	6 FAM-CTCCCACGGACAATCAACG	CCCAGATTGCTGCCTCTCC	TGC(30)	56
Spar08	6 FAM-GGAGTCTCCTTCATTAAGTGCC	GGATGTGTGTGAAGATTGTGC	ATT(30)	56
Spar09	6 FAM-AAGTCCATTTCAAGGCTGGG	CCCATCATAACCCTCCTTTGC	AC(44)	60
Spar010	6 FAM-TCTGGCCTGAGGTATTTGGG	ACTGTAGCCACTTCCCTGCC	AC(44)	60
Spar011	6 FAM-AAGCCACTGCCTTGTGCCC	GACTCTCTGGACATTGGCCC	TC(44)	60
Spar012	6 FAM-GGGAGTGAATGAGAAAGATAAAGTCC	CTGTCATTGCATGGGTTGG	AC(44)	60
Spar013	6 FAM-AAAGATTCTGGAGATCATACCC	TGAATGTATCCTAGGGCGAGC	AC(42)	60
Spar014	6-FAM-TTTCTCTCACTGTCTAACTCTGCC	AGTCCCTGGCAGGTGTGTCC	TC(32)	×
Spar030	6 FAM-AATGGCACCATATTATTCTACATAGG	CCGTCTAGGCTCAGTTTCC	ATT(36)	60
Spar040	6 FAM-GACTGAGACAATTGCTTGAGATAGC	GAGTTTCAGGAGTATTTCACTGC	ATC(33)	60

2003). We used MICROCHECKER to screen null alleles in each locus (Van Oosterhout et al., 2004). We measured polymorphic information content (PIC) with Cervus 3.0.7 (Kalinowski, Taper & Marshall, 2007).

We probed cross-species amplification in tissues of twelve species of the genus: *S. hondurensis*, *S. burtonlimi*, *S. oporophilum*, *S. mordax*, *S. tildae*, *S. erythromos*,

**Table 2** Diagnostic characteristics of selected microsatellites. Number of alleles, size range, polymorphic information (PI), observed heterozygosity ( $H_O$ ), expected heterozygosity ( $H_E$ ), Hardy–Weinberg equilibrium (HWE), and null alleles.

Locus	GenBank accession number	No. alleles	Size range (bp)	PI	$H_O$	$H_E$	HWE	Null alleles
Spar01	KY645946	7	132–236	0.7098	0.941	0.761	0.08	×
Spar02	KY645947	6	130–222	0.6455	0.765	0.692	0.08	×
Spar05	KY645948	6	124–226	0.6069	0.412	0.699	0.05	✓
Spar07	KY645949	10	121–226	0.8028	0.824	0.865	0.18	✓
Spar08	KY645950	11	130–382	0.8052	0.800	0.860	0.13	×
Spar09	KY645951	13	134–230	0.8864	0.875	0.933	0.11	×
Spar10	KY645952	12	132–236	0.8698	0.882	0.919	0.08	×
Spar11	KY645953	8	124–222	0.8125	0.588	0.863	0.12	×
Spar12	KY645954	8	128–214	0.7068	0.750	0.772	0.08	×
Spar13	KY645955	10	124–220	0.8577	0.500	0.867	0.05	✓
Spar030	KY645957	6	133–169	0.7088	0.741	0.735	0.08	×
Spar040	KY645958	6	124–190	0.6721	0.662	0.669	0.08	×

*S. bogotensis*, *S. magna*, *S. new species 3*, *S. luisi*, *S. lilium*, and *S. bakeri* (Supplemental Information 1). All polymorphic loci were tested in the mentioned species by using similar PCR conditions. We followed the ascertainment bias hypothesis of broad amplification in similar phylogenetic species (Schlötterer, 2000).

## RESULTS

We obtained a total of 6,790 potentially amplifiable loci (PALs), containing perfect, imperfect, and compound microsatellites (Fig. 1). Dinucleotide microsatellites were the most abundant (4,288), followed by tetra (1,260); hexa microsatellites were the least abundant in our readings (48). PCR reactions showed that of the 14 loci tested, two were non-specific or monomorphic, and only 12 loci were polymorphic such that we were able to get proper amplification (Table 1). Annealing temperature ranged from 56 to 60 °C.

We found moderate levels of allelic richness, with an average of 8.8 alleles per locus in the representative selection from the wide area of the distribution of *Sturnira parvidens*. Polymorphic information content (PIC) presented values above 0.5 showing a significant content of alleles per locus. Allele frequencies showed a remarkable diversity of alleles per locus, driving a superior number of valuable loci to be used in different genetic analyses (Supplemental Information 2). No evidence of linkage disequilibrium was found on the analyzed loci. We did not observe any loci out of Hardy–Weinberg equilibrium. Levels of expected heterozygosity ( $H_E$ ) ranged from medium to high for all markers (mean  $H_E = 0.79$ , and mean  $H_O = 0.72$ ). In the majority, there was no evidence of null alleles, but three loci (Spar05, Spar07, Spar13) showed significant frequencies of null alleles (above 15%–Table 2).

Cross-species amplification showed differences for the twelve related species (Table 3). *S. new species 3* presented the largest number of amplified microsatellites (8), followed by *S. bakeri* (7). *S. mordax* had the lowest number of amplified loci (4).

**Table 3** Cross-species amplifications of the designed primers for *S. parvidens*. We followed same PCR conditions in the twelve related species. (×) no positive amplification, (✓p) positive polymorphic amplification, (✓m) positive monomorphic amplification, (✓\*) polymorphism not proven because PCR conditions were not standardized.

Locus	<i>S. hondurensis</i> (n = 3)	<i>S. burtonlimi</i> (n = 3)	<i>S. oporaphilum</i> (n = 1)	<i>S. mordax</i> (n = 2)
Spar01	×	✓p	×	×
Spar02	✓p	×	✓*	×
Spar05	✓p	✓p	✓*	✓*
Spar07	×	×	×	×
Spar08	✓*	✓p	✓*	✓p
Spar09	×	✓p	✓*	✓*
Spar10	×	✓*	✓*	×
Spar11	✓*	✓p	✓*	✓p
Spar12	✓m	×	✓*	×
Spar13	×	×	✓*	×

<i>S. tilda</i> (n = 1)	<i>S. erythromos</i> (n = 1)	<i>S. magna</i> (n = 1)	<i>S. bogotensis</i> (n = 1)	<i>S. newspecies_3</i> (n = 3)	<i>S. luisi</i> (n = 3)	<i>S. lilium</i> (n = 3)	<i>S. bakeri</i> (n = 2)
×	×	×	×	✓p	×	✓*	✓*
×	×	✓*	×	✓*	×	×	✓*
✓*	×	✓*	×	✓*	✓*	✓*	×
×	×	×	×	✓p	×	×	✓p
✓*	✓*	×	✓*	✓p	✓*	✓*	✓p
✓*	✓*	×	✓*	✓p	✓*	✓*	✓p
✓*	✓*	✓*	✓*	✓p	✓*	✓p	✓p
✓*	✓*	✓*	✓*	✓*	✓p	✓*	✓*
✓*	✓*	✓*	✓*	×	×	×	×
✓*	✓*	✓*	✓*	×	×	×	×

## DISCUSSION

Next Generation Sequencing allowed the project to obtain a large number of microsatellite loci for *Sturnira parvidens*. This method has been probed for several bat species, and it is becoming a standard method for acquiring specific molecular markers (McCulloch & Stevens, 2011). Given the natural applicability of microsatellites to solve ecological questions, these molecular markers have emerged as a multipurpose indicator for ecological applications (Zane, Bargelloni & Patarnello, 2002; Selkoe & Toonen, 2006). Its applicability spreads to different academic fields such as population genetics, behavioral ecology, genomics, phylogenies, etc.

Our microsatellites conformed to the normal standard measures (Balloux & Lugon-Moulin, 2002). These indicators provide a straightforward approach for describing genetic variation due to the high level of existing alleles. Low allelic richness can affect accuracy in estimating population genetic parameters, leading to significant errors in assessing genetic diversity of target populations (Bashalkhanov, Pandey & Rajora, 2009). Here, we present a novel set of microsatellite loci with the potential to estimate genetic diversity in a non-model species. Standard measures for our microsatellites may have important



implications in the evolutionary biology of the target species, because they can be used to develop conservation strategies for Neotropical bats. Highly informative microsatellites have been used to assess genetic diversity in a broad range of bat populations and to propose measures for conservation (i.e., [Rossiter et al., 2000](#); [Romero-Nava, León-Paniagua & Ortega, 2014](#); [Korstian, Hale & Williams, 2015](#)).

Amplified microsatellites for *S. parvidens* presented levels of polymorphism and heterozygosity similar to those found in other bat species (i.e., *Artibeus jamaicensis*—[Ortega et al., 2002](#); *Rhinolophus ferrumequinum*—[Dawson et al., 2004](#); *Desmodus rotundus*—[Piaggio, Johnston & Perkins, 2008](#); *Corynorhinus* spp.—[Lee, Howell & Van Den Bussche, 2011](#); *Myotis* spp.—[Jan et al., 2012](#); *Carollia castanea*—[Cleary, Waits & Hohenlohe, 2016](#)).

Microsatellite markers are widely used to infer levels of genetic diversity in natural populations. Molecular markers are not always developed for the target species and the use of microsatellite loci from related species can be accurate. Ascertainment bias limited the microsatellite-based amplification due to the particular selection of polymorphic markers in the target species, plus the reduced sensitivity of the markers due to the phylogenetic constrictions of the particular evolutionary traits of each sister species ([Crawford et al., 1998](#); [Schlötterer, 2000](#); [Li & Kimmel, 2013](#)). The bias leads to a lower average allele length due to the phylogenetic restriction provided by the unique evolutionary history of each species ([Li & Kimmel, 2013](#)). We tested the potential use of our markers in related species, finding multilocus heterozygosities inside the *Sturnira* genus. This positive effect suggests the use of the developed markers to extrapolate genetic diversity in future studies for this highly speciose genus, in which the past demographic shared histories barely affect the cross-species amplification consolidation.

## CONCLUSIONS

We used Illumina Paired-Sequences to efficiently develop microsatellite loci for *Sturnira parvidens*. We formed a genomic library to obtain 12 specific and polymorphic microsatellites for this bat. Microsatellites showed high allelic richness per locus, showing their effectiveness for further studies (i.e., population genetics, behavioral ecology, etc.). Cross-species amplification was effective for the 12 related species, but with no positive amplifications in several cases.

## ACKNOWLEDGEMENTS

We are grateful for the supporting fieldwork provided by students of the Facultad de Ciencias, UNAM (MZFC-M). We would like to thank the Field Museum of Natural History, Chicago (FMNH); Louisiana State University, Louisiana State University, Museum of Zoology, Baton Rouge (LSUMZ); Museo de Zoología de la Universidad de Costa Rica, San José, Costa Rica (MZUCR); Museum of Texas Tech University, Lubbock (TTU) for providing tissues from their collections. Particular special thanks go to Bruce D. Patterson and Natalia Cortés-Delgado from FMNH; David Villalobos and Bernal Rodríguez from MZUCR; Frederick H. Sheldon and Donna L. Dittmann from LSUMZ; and Caleb D. Phillips from TTU for providing samples.

## ADDITIONAL INFORMATION AND DECLARATIONS

### Funding

This work was granted by Proyecto Ciencia Básica CONACyT (239482) and Proyecto de Investigación SIP IPN (20160744). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

### Grant Disclosures

The following grant information was disclosed by the authors:

Proyecto Ciencia Básica CONACyT: 239482.

Proyecto de Investigación SIP IPN: 20160744.

### Competing Interests

The authors declare there are no competing interests.

### Author Contributions

- Edgar G. Gutiérrez and Giovani Hernández Canchola conceived and designed the experiments, analyzed the data, wrote the paper, prepared figures and/or tables, reviewed drafts of the paper.
- Livia S. León Paniagua and Jorge Ortega conceived and designed the experiments, contributed reagents/materials/analysis tools, wrote the paper, reviewed drafts of the paper.
- Norberto Martínez Méndez conceived and designed the experiments, analyzed the data, wrote the paper, reviewed drafts of the paper.

### Field Study Permissions

The following information was supplied relating to field study approvals (i.e., approving body and any reference numbers):

Secretaría de Medio Ambiente y Recursos Naturales (SEMARNAT) provided permit for tissue collection FAUT-0307.

### DNA Deposition

The following information was supplied regarding the deposition of DNA sequences:

Gen Bank: Spar01 [KY645946](#), Spar02 [KY645947](#), Spar05 [KY645948](#), Spar07 [KY645949](#), Spar08 [KY645950](#), Spar09 [KY645951](#), Spar010 [KY645952](#), Spar011 [KY645953](#), Spar012 [KY645954](#), Spar013 [KY645955](#), Spar030 [KY645957](#), Spar040 [KY645958](#).

### Data Availability

The following information was supplied regarding data availability:

The raw data has been supplied as a [Supplementary File](#).

### Supplemental Information

Supplemental information for this article can be found online at <http://dx.doi.org/10.7717/peerj.3367#supplemental-information>.



## REFERENCES

- Aljanabi S, Martinez I. 1997.** Universal and rapid salt extraction of high quality genomic DNA for PCR-based techniques. *Nucleic Acid Research* **25**:4692–4693 DOI [10.1093/nar/25.22.4692](https://doi.org/10.1093/nar/25.22.4692).
- Balloux F, Lugon-Moulin N. 2002.** The estimation of population differentiation with microsatellite markers. *Molecular Ecology* **11**:155–165 DOI [10.1046/j.0962-1083.2001.01436.x](https://doi.org/10.1046/j.0962-1083.2001.01436.x).
- Bashalkhanov S, Pandey M, Rajora OP. 2009.** A simple method for estimating genetic diversity in large populations from finite sample sizes. *BMC Genetics* **10**:84 DOI [10.1186/1471-2156-10-84](https://doi.org/10.1186/1471-2156-10-84).
- Castoe TA, Poole AW, De Koning APJ, Jones KL, Tomback DF, Oyler-McCance SJ, Fike JA, Lance SL, Streicher JW, Smith EN, Pollack DD. 2012.** Rapid microsatellite identification from Illumina paired-end genomic sequencing in two birds and a snake. *PLOS ONE* **7**:e30953 DOI [10.1371/journal.pone.0030953](https://doi.org/10.1371/journal.pone.0030953).
- Cleary KA, Waits LP, Hohenlohe PA. 2016.** Development and characterization of fourteen novel microsatellite markers for the chestnut short-tailed fruit bat (*Carollia castanea*), and cross-amplification to related species. *PeerJ* **4**:e2465 DOI [10.7717/peerj.2465](https://doi.org/10.7717/peerj.2465).
- Crawford AM, Kappes SM, Paterson KA, De Gortari MJ, Dodds KG, Freking BA, Stone RT, Beattie CW. 1998.** Microsatellite evolution: testing the ascertainment bias hypothesis. *Journal of Molecular Evolution* **46**:256–260 DOI [10.1007/PL00006301](https://doi.org/10.1007/PL00006301).
- Dawson DA, Rossiter SJ, Jones G, Faulkes CG. 2004.** Microsatellite loci for the greater horseshoe bat, *Rhinolophus ferrumequinum* (Rhinolophidae, Chiroptera) and their cross-utility in 17 other bat species. *Molecular Ecology Notes* **4**:96–100 DOI [10.1046/j.1471-8286.2003.00580.x](https://doi.org/10.1046/j.1471-8286.2003.00580.x).
- Evelyn MJ, Stiles DA. 2003.** Roosting requirements of two frugivorous bats (*Sturnira lilium* and *Artibeus intermedius*) in fragmented Neotropical forest. *Biotropica* **35**:405–418 DOI [10.1111/j.1744-7429.2003.tb00594.x](https://doi.org/10.1111/j.1744-7429.2003.tb00594.x).
- Excoffier L, Laval G, Schneider S. 2003.** Arlequin ver. 3.0: an integrated software package for population genetics data analysis. *Evolutionary Bioinformatics Online* **1**:47–50.
- Faircloth BC. 2008.** MSTACOMMANDER: detection of microsatellite repeat arrays and automated, locus-specific primer design. *Molecular Ecology Resources* **8**:92–94 DOI [10.1111/j.1471-8286.2007.01884.x](https://doi.org/10.1111/j.1471-8286.2007.01884.x).
- Gannon WL, Sikes RS, Animal Care and Use Committee of the American Society of Mammalogists. 2007.** Guidelines of the American Society of Mammalogists for the use of wild mammals in research. *Journal of Mammalogy* **88**:809–823 DOI [10.1644/06-MAMM-F-185R1.1](https://doi.org/10.1644/06-MAMM-F-185R1.1).
- Jan C, Dawson DD, Altringham JD, Burke T, Butlin RK. 2012.** Development of conserved microsatellite markers of high cross-species utility in bat species (Vespertilionidae, Chiroptera, Mammalia). *Molecular Ecology Resources* **12**:532–548 DOI [10.1111/j.1755-0998.2012.03114.x](https://doi.org/10.1111/j.1755-0998.2012.03114.x).

- Kalinowski ST, Taper ML, Marshall TC. 2007.** Revising how the computer program CERVUS accommodates genotyping error increases success in paternity assignment. *Molecular Ecology* **16**:1099–1106 DOI [10.1111/j.1365-294X.2007.03089.x](https://doi.org/10.1111/j.1365-294X.2007.03089.x).
- Korstian JM, Hale AM, Williams DA. 2015.** Genetic diversity, historic population size, and population structure in 2 North American tree bats. *Journal of Mammalogy* **96**:972–980 DOI [10.1093/jmammal/gyv101](https://doi.org/10.1093/jmammal/gyv101).
- Lee DN, Howell JM, Van Den Bussche RA. 2011.** Development and characterization of 15 polymorphic tetranucleotide microsatellite loci for Townsend's big-eared bat (*Corynorhinus townsendii*) and cross amplification in Rafinesque's big-eared bat (*Corynorhinus rafinesquii*). *Conservation Genetic Resources* **4**:429–433 DOI [10.1007/s12686-011-9566-5](https://doi.org/10.1007/s12686-011-9566-5).
- Lewis P, Zaykin D. 2001.** Genetic data analysis: computer program for the analysis of allelic data. Version 10 (d16c). Available at <http://lewis.eeb.uconn.edu/lewishome/software.html> (accessed on October 2016).
- Li B, Kimmel M. 2013.** Factors influencing ascertainment bias of microsatellite allele sizes: impact on estimates of mutation rates. *Genetics* **195**:563–572 DOI [10.1534/genetics.113.154161](https://doi.org/10.1534/genetics.113.154161).
- McCulloch ES, Stevens RS. 2011.** Rapid development and screening of microsatellite loci for *Artibeus lituratus* and their utility for six related species within Phyllostomidae. *Molecular Ecology* **11**:903–913 DOI [10.1111/j.1755-0998.2011.03027.x](https://doi.org/10.1111/j.1755-0998.2011.03027.x).
- Molinari J, Bustos XE, Burneo SF, Camacho MA, Moreno SA, Fermín G. 2017.** A new polytypic species of yellow-shouldered bats, genus *Sturnira* (Mammalia: Chiroptera: Phyllostomidae), from the Andean and coastal mountain systems of Venezuela and Colombia. *Zootaxa* **4243**(1):75–96 DOI [10.11646/zootaxa.4243.1.3](https://doi.org/10.11646/zootaxa.4243.1.3).
- Olea-Wagner A, Lorenzo C, Naranjo E, Ortiz D, León-Paniagua L. 2007.** Diversidad de frutos que consumen tres especies de murciélagos (Chiroptera: Phyllostomidae) en la selva Lacandona, Chiapas, México. *Revista Mexicana de Biodiversidad* **78**:191–200.
- Ortega J, Maldonado JE, Arita HT, Wilkinson G, Fleischer R. 2002.** Characterization of microsatellite loci in the Jamaican fruit-eating bat *Artibeus jamaicensis* and cross-species amplification. *Molecular Ecology Notes* **2**:462–464 DOI [10.1046/j.1471-8286.2002.00278.x](https://doi.org/10.1046/j.1471-8286.2002.00278.x).
- Piaggio AJ, Johnston JJ, Perkins SL. 2008.** Permanent genetic resources: development of polymorphic microsatellite loci for the common vampire bat, *Desmodus rotundus* (Chiroptera: Phyllostomidae). *Molecular Ecology Resources* **8**:440–442 DOI [10.1111/j.1471-8286.2007.01986.x](https://doi.org/10.1111/j.1471-8286.2007.01986.x).
- Romero-Nava C, León-Paniagua LS, Ortega J. 2014.** Microsatellites loci reveal heterozygosity and population structure in vampire bats (*Desmodus rotundus*) (Chiroptera: Phyllostomidae) of Mexico. *Revista de Biología Tropical* **52**:659–669.
- Rossiter SJ, Jones G, Ransome RD, Barrat EM. 2000.** Genetic variation and population structure in the endangered greater horseshoe bat *Rhinolophus ferrumequinum*. *Molecular Ecology* **9**:1131–1135 DOI [10.1046/j.1365-294x.2000.00982.x](https://doi.org/10.1046/j.1365-294x.2000.00982.x).

- Rousset F. 2008.** Genepop'007: a complete reimplementation of the Genepop software for Windows and Linux. *Molecular Ecology Resources* **8**:103–106 DOI [10.1111/j.1471-8286.2007.01931.x](https://doi.org/10.1111/j.1471-8286.2007.01931.x).
- Selkoe KA, Toonen RJ. 2006.** Microsatellites for ecologists: a practical guide to using and evaluating microsatellite markers. *Ecology Letters* **9**:615–629 DOI [10.1111/j.1461-0248.2006.00889.x](https://doi.org/10.1111/j.1461-0248.2006.00889.x).
- Schlötterer C. 2000.** Evolutionary dynamics of microsatellite DNA. *Chromosoma* **109**:365–371 DOI [10.1007/s004120000089](https://doi.org/10.1007/s004120000089).
- Van Oosterhout C, Hutchinson WF, Wills DPM, Shipley P. 2004.** MICRO-CHECKER: software for identifying and correcting genotyping errors in microsatellite data. *Molecular Ecology Notes* **4**:535–538 DOI [10.1111/j.1471-8286.2004.00684.x](https://doi.org/10.1111/j.1471-8286.2004.00684.x).
- Velazco PM, Patterson BD. 2013.** Diversification of the yellow-shouldered bats, genus *Sturnira* (Chiroptera, Phyllostomidae), in the New World tropics. *Molecular Phylogenetics and Evolution* **68**:683–698 DOI [10.1016/j.ympev.2013.04.016](https://doi.org/10.1016/j.ympev.2013.04.016).
- Velazco PM, Patterson BD. 2014.** Two new species of yellow-shouldered bats, genus *Sturnira* Gray, 1842 (Chiroptera: Phyllostomidae) from Costa Rica, Panama and western Ecuador. *ZooKeys* **402**:43–66 DOI [10.3897/zookeys.402.7228](https://doi.org/10.3897/zookeys.402.7228).
- Villalobos F, Valerio AA. 2002.** The phylogenetic relationships of the bat genus *Sturnira* Gray, 1842 (Chiroptera: Phyllostomidae). *Mammalian Biology* **67**:268–275.
- Zane L, Bargelloni L, Patarnello T. 2002.** Strategies for microsatellite isolation: a review. *Molecular Ecology* **11**:1–6.