



Spatial distribution of environmental DNA in a nearshore marine habitat

James L. O'Donnell¹, Ryan P. Kelly¹, Andrew Olaf Shelton²,
Jameal F. Samhoury³, Natalie C. Lowell^{1,4} and Gregory D. Williams⁵

¹ School of Marine and Environmental Affairs, University of Washington, Seattle, WA, United States of America

² Earth Resource Technology, Inc., Under Contract to the Northwest Fisheries Science Center, National Marine Fisheries Service, National Oceanic and Atmospheric Administration, Seattle, WA, United States of America

³ Conservation Biology Division, Northwest Fisheries Science Center, National Marine Fisheries Service, National Oceanic and Atmospheric Administration, Seattle, WA, United States of America

⁴ School of Aquatic and Fishery Sciences, University of Washington, Seattle, WA, United States of America

⁵ Pacific States Marine Fisheries Commission, Under Contract to the Northwest Fisheries Science Center, National Marine Fisheries Service, National Oceanic and Atmospheric Administration, Seattle, WA, United States of America

ABSTRACT

In the face of increasing threats to biodiversity, the advancement of methods for surveying biological communities is a major priority for ecologists. Recent advances in molecular biological technologies have made it possible to detect and sequence DNA from environmental samples (environmental DNA or eDNA); however, eDNA techniques have not yet seen widespread adoption as a routine method for biological surveillance primarily due to gaps in our understanding of the dynamics of eDNA in space and time. In order to identify the effective spatial scale of this approach in a dynamic marine environment, we collected marine surface water samples from transects ranging from the intertidal zone to four kilometers from shore. Using PCR primers that target a diverse assemblage of metazoans, we amplified a region of mitochondrial 16S rDNA from the samples and sequenced the products on an Illumina platform in order to detect communities and quantify their spatial patterns using a variety of statistical tools. We find evidence for multiple, discrete eDNA communities in this habitat, and show that these communities decrease in similarity as they become further apart. Offshore communities tend to be richer but less even than those inshore, though diversity was not spatially autocorrelated. Taxon-specific relative abundance coincided with our expectations of spatial distribution in taxa lacking a microscopic, pelagic life-history stage, though most of the taxa detected do not meet these criteria. Finally, we use carefully replicated laboratory procedures to show that laboratory treatments were remarkably similar in most cases, while allowing us to detect a faulty replicate, emphasizing the importance of replication to metabarcoding studies. While there is much work to be done before eDNA techniques can be confidently deployed as a standard method for ecological monitoring, this study serves as a first analysis of diversity at the fine spatial scales relevant to marine ecologists and confirms the promise of eDNA in dynamic environments.

Submitted 28 November 2016

Accepted 29 January 2017

Published 28 February 2017

Corresponding author

James L. O'Donnell,
jimmyod@uw.edu

Academic editor

Magnus Johnson

Additional Information and
Declarations can be found on
page 18

DOI 10.7717/peerj.3044

Distributed under
Creative Commons Public
Domain Dedication

OPEN ACCESS

Subjects Biodiversity, Ecology, Marine Biology, Molecular Biology

Keywords Marine, Metabarcoding, Metagenomics, Estuarine, Molecular ecology, Environmental monitoring

INTRODUCTION

The patterns and causes of variability in ecological communities across space are both seminal and contentious areas of study in ecology (Hubbell, 2001; Anderson et al., 2011). One consistently observed pattern of community spatial heterogeneity is that communities close to one another tend to be more similar than those that are farther apart (Nekola & White, 1999). This decrease in community similarity with increasing spatial separation is called distance decay and has been reported from communities of tropical trees (Condit, 2002; Chust et al., 2006), ectomycorrhizal fungi (Bahram et al., 2013), salt marsh plants (Guo et al., 2015), and microorganisms (Martiny et al., 2011; Chust et al., 2013; Wetzel et al., 2012; Bell, 2010). Typically, this relationship is assessed by regressing a measure of community similarity against a measure of spatial separation for a set of sites at which a set of species' abundances (or presences) is calculated. Yet no existing biodiversity survey method completely censuses all of the organisms in a given area. The lack of a single 'silver bullet' method of sampling contributes to inconclusiveness to the study of spatial patterning in ecology (Levin, 1992), and leaves open the possibility of new and more comprehensive methods.

From a boat or aircraft, scientists can count whales by sight, but not the krill on which they feed. For example, towed fishing nets can efficiently sample organisms larger than the mesh and slower than the boat, but overlook viruses and have undesirable effects on charismatic air-breathing species. However, DNA-based surveys show great promise as an efficient technique for detecting a previously unthinkable breadth of organisms from a single sample.

Microbiologists have used nucleic acid sequencing to quantify the composition and function of microbial communities in a wide variety of habitats (Handelsman et al., 1998; Tyson et al., 2004; Venter et al., 2004; Iverson et al., 2012). To do so, microorganisms are collected in a sample of environmental medium (e.g., water), their DNA or RNA is isolated and sequenced, and the identity and abundance of sequences is considered to reflect the community of organisms contained in the sample, which indirectly estimates the quantity of organisms in an area.

Macroorganisms shed DNA-containing cells into the environment (environmental DNA or eDNA) that can be sampled in the same way (Ficetola et al., 2008; Thomsen et al., 2012). Potentially, eDNA methods allow a broad swath of macroorganisms to be surveyed from basic environmental samples. However, the accuracy and reliability of indirect estimates of macroorganismal abundance has been debated because the entire organisms are not contained within the sample (Coward et al., 2015). Concern surrounding eDNA methods is rooted in uncertainty about the attributes of eDNA in the environment relative to actual organisms (Shelton et al., 2016; Evans et al., 2016). Basic questions such as how long DNA can persist in that environment and how far DNA can travel remain largely unknown

(but see *Klymus et al. (2015)*; *Turner, Uy & Everhart (2015)*; *Strickler, Fremier & Goldberg (2015)*; *Deiner & Altermatt (2014)*) and impede inference about local organismal presence from an environmental sample. As a result, estimating the spatial and temporal resolution of eDNA studies in the field is a key step in making these methods practical.

The relationship between local organismal abundance and eDNA is further complicated in habitats where the environmental medium itself may transport eDNA away from its source. We know that genetic material can move away from its source precisely because organisms can be detected indirectly without being present in the sample (*Kelly et al., 2016b*). One might reasonably expect eDNA to travel farther in a highly dynamic fluid such as the open ocean or flowing river than it would through the sediment at the bottom of a stagnant pond (*Deiner & Altermatt, 2014*; *Shogren et al., 2016*). Yet even studies of extremely dynamic habitats such as coastlines with high wave energy have found remarkable evidence that eDNA transport is limited enough that DNA methods can detect differences among communities separated by less than 100 m (*Port et al., 2016*).

While rigorous laboratory studies have investigated the effects of some environmental factors on eDNA persistence (*Klymus et al., 2015*; *Barnes et al., 2014*; *Sassoubre et al., 2016*) and the transport of eDNA in specific contexts (*Deiner & Altermatt, 2014*), we suggest that field studies comparing the spatial distribution of communities of eDNA with expectations based on prior knowledge of organisms' distributions are also critical to developing a working understanding of eDNA in the real world. Research to date has documented the non-random spatial distribution of meiofaunal (*Fonseca et al., 2014*; *Guardiola et al., 2016*), microbial (*Lallias et al., 2015*), and extracellular (*Guardiola et al., 2015*) eDNA of marine and estuarine sediments, and of microscopic plankton in open ocean waters (*De Vargas et al., 2015*). These studies conducted targeted sampling at intermediate (thousands of meters) to global (thousands of kilometers) scales. Here, we use a grid-based environmental sampling strategy to assess spatial variability of eDNA in a coastal marine environment at a fine scale (tens to thousands of meters), using molecular methods that focus on microbial metazoans.

We apply methods derived from community ecology to understand spatial patterns and patchiness of eDNA. The underlying mechanism thought to drive the slope of the distance decay relationship in ecological communities is the rate of movement of individuals among sites, which may be driven by underlying processes such as habitat suitability. Because eDNA is shed and transported away from its source, the increased movement of eDNA particles should homogenize community similarity, and thus erode the distance decay relationship of eDNA communities.

Puget Sound is a deep, narrow fjord in Washington, USA, where a narrow band of shallow bottom hugs the shoreline and abruptly gives way to a central depth of up to 300 m. This form allows the juxtaposition of communities associated with distinctly different habitats: shallow, intertidal benthos, and euphotic pelagic (*Burns, 1985*). At the upper reaches of the intertidal, the shoreline substrate varies from soft, fine sediment to cobble and boulder rubble. Soft intertidal sediments are inhabited by burrowing bivalves (Bivalvia), segmented worms (Annelida), and acorn worms (Enteropneusta), and in some lower intertidal and high subtidal ranges by eelgrass (*Zostera marina*) (*Kozloff, 1973*; *Dethier, 2010*). Eelgrass

meadows harbor epifaunal and infaunal biota, and attract transient species which use the meadows for shelter and to feed on resident organisms. Hard intertidal surfaces support a well-documented biota including barnacles (Sessilia) and other crustaceans, mussels (Bivalvia:Mytilidae), anemones (Actiniaria), sea stars (Asteroidea), urchins (Echinoidea), Bryozoans (Ectoprocta), crustaceans (Decapoda), and a variety of algae ([Dethier, 2010](#)). Hard bottoms of the lower intertidal and high subtidal are home to macroalgae such as Laminariales and Desmarestiales which provide habitat for a distinct community of fish and invertebrates. The upper pelagic is home to a diverse assemblage of microscopic plankton including diatoms, copepods, and larvae ([Strickland, 1983](#)), as well as transitory fish and marine mammals.

We took advantage of this setting to explore the spatial variation and distribution of marine eDNA communities. Using PCR-based methods and massively parallel sequencing, we surveyed mitochondrial 16S sequences from a suite of marine animals in water samples collected over a grid of sites extending from the shoreline out to 4 kilometer offshore in Puget Sound, Washington, USA. We leverage this sampling design to perform an explicitly spatial analysis of eDNA-derived community similarity. We investigate two primary objectives. First we examine the spatial patterning of eDNA and determine the degree to which eDNA community similarity can be predicted by physical proximity. We expect that physical proximity will be a strong predictor of community similarity, and that community differences can be detected over small distances. Second, we examine the distribution of diversity from eDNA data, and compare it to our expectations based on distributions of macrobial communities. We expect that distinct eDNA communities exist in this setting, and that their spatial distribution coincides with that of adult macrobial organisms. Because of the vastly different communities of benthic macrobial metazoans as a function of distance from shore, we expect that more than one eDNA community is present across our 4 kilometer sampling grid, and that communities change as a function of distance from shore. For this reason, we examine two diversity measures of eDNA communities that have been widely used to reveal broad scale patterns based on macrobiota in many ecological systems. Finally, we identify the taxa represented in the eDNA communities, which span a range of life-history characteristics, and we expect that the spatial distribution of eDNA will most closely resemble the distribution of adults in taxa with low dispersal potential.

METHODS

There are seven discrete steps to our methodology: (1) Environmental sample collection, (2) isolation of particulates from water via filtration, (3) isolation of DNA from filter membrane, (4) amplification of target locus via PCR, (5) sequencing of amplicons, (6) bioinformatic translation of raw sequence data into tables of sequence abundance among samples, and (7) community ecological analyses of eDNA. We provide brief overviews of these steps here, and encourage the reader to review the fully detailed methods presented in [Supplemental Information 1](#).




Figure 1 Map of study area. Depth in meters below sea level is indicated by shading and 25 meter contours. Sampled locations are indicated by red points.

Environmental sampling

Starting from lower-intertidal patches of *Zostera marina*, we collected water samples at 1 m depth from 8 points (0, 75, 125, 250, 500, 1,000, 2,000, and 4,000 m) along three parallel transects separated by 1,000 m (24 sample locations total; Fig. 1). Samples were collected by attaching bottles to a PVC pole and lowering it over the side of a boat over the span of one hour on 27 June 2014. To destroy residual DNA on equipment used for field sampling and filtration, we washed with a 1:10 solution of household bleach (8.25% sodium hypochlorite; 7.25% available chlorine) and deionized water, followed by thorough rinsing with deionized water. Each environmental sample was collected in a clean 1 liter high-density polyethylene bottle, the opening of which was covered with 500 micrometer nylon mesh to prevent entry of larger particles. Immediately after collecting the sample, the mesh was replaced with a clean lid and the sample was held on ice until filtering.

Filtration

One liter from each water sample was filtered in the lab on a clean polysulfone vacuum filter holder fitted with a 47 millimeter diameter cellulose acetate membrane with 0.45 micrometer pores. Filter membranes were moved into 900 microliters of Longmire buffer (Longmire, Maltbie & Baker, 1997) using clean forceps and stored at room temperature

(*Renshaw et al., 2015*). To test for the extent of contamination attributable to laboratory procedures, we filtered three replicate 1 liter samples of deionized water. These samples were treated identically to the environmental samples throughout the remaining protocols.

DNA purification

DNA was purified from the membrane following a phenol:chloroform:isoamyl alcohol protocol following Renshaw (*Renshaw et al., 2015*). Preserved membranes were incubated at 65 °C for 30 min before adding 900 microliters of phenol:chloroform:isoamyl alcohol and shaking vigorously for 60 s. We conducted two consecutive chloroform washes by centrifuging at 14,000 rpm for 5 min, transferring the aqueous layer to 700 microliters chloroform, and shaking vigorously for 60 s. After a third centrifugation, 500 microliters of the aqueous layer was transferred to tubes containing 20 microliters 5 molar NaCl and 500 microliters 100% isopropanol, and frozen at –20 °C for approximately 15 h. Finally, all liquid was removed by centrifuging at 14,000 rpm for 10 min, pouring off or pipetting out any remaining liquid, and drying in a vacuum centrifuge at 45 °C for 15 min. DNA was resuspended in 200 microliters of ultrapure water. Four replicates of genomic DNA extracted from tissue of a species absent from the sampled environment (*Oreochromis niloticus*) served as positive control for the remaining protocols.

PCR amplification

We chose a primer set that amplifies an approximately 115 base pair (bp) region of the mitochondrial 16S rRNA gene in at least 10 metazoan phyla from this habitat, excludes non-metazoans, and resolves taxonomy to the family level in most cases using a public sequence database (*Kelly et al., 2016a*). We used a two-step polymerase chain reaction (PCR) protocol described by *O'Donnell et al. (2016)* to generate four replicate products from each DNA sample. In the first set of reactions, primers were identical in every reaction (forward: AGTTACYYTAGGGATAACAGCG; reverse: CCGGTCTGAACTCAGATCAYGT); primers in the second set of reactions included these same sequences but with 3 variable nucleotides (NNN) and an index sequence on the 5' end (see Sequencing Metadata). We used the program OligoTag (*Coissac, 2012*) to generate 30 unique 6-nucleotide index sequences differing by a minimum Hamming distance of 3 (see Sequencing Metadata). Indexed primers were assigned to samples randomly, with the identical index sequence on the forward and reverse primer to avoid errors associated with dual-indexed multiplexing (*Schnell, Bohmann & Gilbert, 2015*). In a UV-sterilized hood, we prepared 25 microliter reactions containing 18.375 microliters ultrapure water, 2.5 microliters 10× buffer, 0.625 microliters deoxynucleotide solution (8 millimolar), 1 microliter each forward and reverse primer (10 micromolar, obtained lyophilized from Integrated DNA Technologies; Coralville, IA, USA), 0.25 microliters Qiagen HotStar Taq polymerase, and 1.25 microliter genomic or eDNA template at 1:100 dilution in ultrapure water. PCR thermal profiles began with an initialization step (95 °C; 15 min) followed by cycles (40 and 20 for the first and second reaction, respectively) of denaturation (95 °C; 15 s), annealing (61 °C; 30 s), and extension (72 °C; 30 s). A total of 20 identical PCRs were conducted from each DNA extract using non-indexed primers; these were pooled into four groups of five in order to

ensure ample template for the subsequent PCR with indexed primers. In order to isolate the fragment of interest from primer dimer and other spurious fragments generated in the first PCR, we used the AxyPrep Mag FragmentSelect-I kit with solid-phase reversible immobilization (SPRI) paramagnetic beads at 2.5× the volume of PCR product (Axygen BioSciences, Corning, NY, USA). A 1:5 dilution in ultrapure water of the product was used as template for the second reaction. PCR products of the second reaction were purified using the Qiagen MinElute PCR Purification Kit (Qiagen, Hilden, Germany). Ultrapure water was used in place of template DNA and run along with each batch of PCRs to serve as a negative control for PCR; none of these produced visible bands on an agarose gel. In total, four separate replicates from each of 31 DNA samples were carried through the two-step PCR process for a total of 124 sequenced PCR products. These were combined with additional samples from other projects, totaling 345 samples for sequencing.

DNA sequencing

Up to 30 PCR products were combined according to their primer index in equal concentration into one of 14 pools, and 150 nanograms from each were prepared for library sequencing using the KAPA high-throughput library prep kit with real-time library amplification protocol (KAPA Biosystems, Wilmington, MA, USA). Each of these ligated sequencing adapters included an additional six base pair index sequence (NEXTflex DNA barcodes; BIOO Scientific, Austin, TX, USA). Thus, each PCR product was identifiable via its unique combination of index sequences in the sequencing adapters and primers. Fragment size distribution and concentration of each library was quantified using an Agilent 2100 BioAnalyzer. Libraries were pooled in equal concentrations and sequenced for 150 base pairs in both directions (PE150) using an Illumina NextSeq at the Stanford Functional Genomics Facility, where 20% PhiX Control v3 was added to act as a sequencing control and to enhance sequencing depth by increasing sequence diversity. Raw sequence data in fastq format is publicly available (see Data Availability).

Sequence Data Processing (Bioinformatics)

Detailed bioinformatic methods are provided in the supplemental material, and analysis scripts used from raw sequencer output onward can be found in the public project directory (see Analysis Scripts). Briefly, we performed five steps to process the sequence data: (1) Merge paired-end reads (*Zhang et al., 2014*), (2) eliminate low-quality reads (*Edgar, 2010; Rognes et al., 2016*), (3) eliminate PCR artifacts (chimeras) (*Edgar, 2010; Rognes et al., 2016; Martin, 2011*), (4) cluster reads by similarity into operational taxonomic units (OTUs) (*Mahé et al., 2014*), and (5) match observed sequences to taxon names (*Camacho et al., 2009; Chamberlain & Szöcs, 2013; Chamberlain et al., 2016*). Additionally, we checked for consistency among PCR replicates, excluded extremely rare sequences, and rescaled (rarefied) the data to account for differences in sequencing depth. The data for input to further analyses are a contingency table of the mean count of unique sequences, OTUs, or taxa present in each environmental sample.

Ecological analyses

After gathering the data, we use the eDNA community observed at each location to make inferences about the spatial patterning of eDNA communities. We use statistical tools from community ecology to assess the spatial structure of eDNA communities. We report similarity (1 - dissimilarity) rather than dissimilarity in all cases for ease of interpretation.

Objective 1: community similarity as a function of distance

Distance decay

To address our first objective and determine whether or not nearby samples are more similar than distant ones, we fit a nonlinear model to represent decreasing community similarity with distance. We calculated the pairwise Bray–Curtis similarity (1—Bray–Curtis dissimilarity) between eDNA communities using the R package *vegan* (Oksanen et al., 2016) and the great circle distance between sampling points using the Haversine method as implemented by the R package *geosphere* (Hijmans, 2016). This model is similar to the Michaelis–Menten function, but with an asymptote fixed at 0:

$$y_{ij} = \frac{AB}{B + x_{ij}} \quad (1)$$

Where the relationship between community similarity (y_{ij}) and spatial distance (x_{ij}) between observations i and j is determined by the similarity of samples at distance 0 (A), and the distance at which half the total change in similarity is achieved (B). This allows for samples collected very close together (near 0) to have similarity significantly less than one. We assessed model fit using the R function `nls` (R Core Team, 2016), using the `nl2sol` algorithm from the `Port` library to solve separable nonlinear least squares using analytically computed derivatives (<http://netlib.org/port/nsg.f>). We set bounds of 0 and 1 for the intercept parameter and a lower bound of 0 for the distance at half similarity; starting values of these parameters were 0.5 and $x_{\max}/2$, respectively. We calculated a 95% confidence interval for the parameters and the predicted values using a first-order Taylor expansion approach implemented by the function `predictNLS` in the R package `propagate` (Spiess, 2014).

There are other conceptually reasonable forms to expect the space-by-similarity relationship to take; we present these in the supplemental material along with alternative data subsets and similarity indices.

Objective 2: spatial distribution of diversity

Community classification

To determine the spatial distribution and variation of eDNA communities (objective 2), we used multivariate classification algorithms. We simultaneously assessed the existence of distinct community types and the membership of samples to those community types using an unsupervised classification algorithm known as partitioning around medoids (PAM; sometimes referred to as k -medoids clustering) (Kaufman & Rousseeuw, 1990), as implemented in the R package `cluster` (Maechler et al., 2016). The classification of samples to communities was made on the basis of their pairwise Bray–Curtis similarity, calculated using the function `vegdist` in the R package `vegan` (Oksanen et al., 2016). Other distance

metrics were evaluated but had no appreciable effect on the outcome of the analysis (Fig. S1). In order to choose an optimal number of clusters (K), we evaluated the distribution of silhouette widths, a measure of the similarity between each sample and its cluster compared to its similarity to other clusters. We repeated the analysis using fuzzy clustering (FANNY, (Kaufman & Rousseeuw, 1990)); however, the results were qualitatively similar to the results using PAM so we omit them here.

Aggregate measures of diversity

We calculated two measures of diversity, richness and evenness, to ask if aggregate metrics of the eDNA community showed evidence of spatial patterning. Richness is a measure of the number of distinct types of organisms present and so ranges from 1 (only one taxon observed) to S , the number of taxa observed across all samples. To calculate the evenness of the distribution of abundance of taxa in a sample, we used the complement of the Simpson (1949) index ($1 - \sum p_i^2$, where p_i is the proportional abundance of taxon i). The values of this index range from 0 to 1, with the value interpreted as the probability that two sequences randomly selected from the sample will belong to different taxa; thus, larger values of the index indicate more evenly divided communities (Magurran, 2004). We calculated Moran's I for both diversity metrics to test for spatial autocorrelation. We also tested for a linear effect of log-transformed distance from shore on each measure of diversity to ask how diversity changes over this strong environmental gradient.

Taxon and life history patterns

After assigning taxon names to the abundance data, we plotted the distribution in space of a selection of taxa to compare with our expectations on the basis of adult distributions (objective 2). Our aim was to understand where each taxon occurred in the greatest proportional abundance, and its distribution in space relative to that maximum. Thus, we rescaled each sample to proportional abundance, extracted the data from a single taxon, and scaled those values between 0 and 1. We collated life history characteristics for each of the major taxonomic groups recovered, including dispersal range of the gametes, larvae, and adults, adult habitat type and selectivity, and adult body size. For each life history stage of each taxon group, we made an order-of-magnitude approximation of the scale of dispersal. For example, internally fertilized species were assigned a gamete range of 0 km, while broadcast spawners were assigned a gamete range of 10 km. Similarly, adult range size was approximated as 0 km (sessile), 1 km (motile but not pelagic), or 10 km (highly mobile, pelagic). Variables were specified as 'multiple' for life history stages known to span more than 1 magnitude of range size. For groups to which sequences were annotated with high confidence, but for which life history strategy is diverse or poorly known (e.g., families in the phylum Nemertea), we used conservative, coarse approximations at a higher taxonomic rank (see Life History Data). These data were used to contextualize group-specific spatial distributions and inform expectations based on known adult distributions.

RESULTS

Sequence data processing (bioinformatics)

Preliminary sequence analysis strongly suggested that the observed variation among environmental samples reflects true variation in the environment, rather than variability due to lab protocols, for the following reasons (note that all value ranges are reported as mean plus and minus one standard deviation). First, all libraries passed the FastQC per-base sequence quality filter, generating a total of 371,576,190 reads passing filter generated in each direction. Second, samples in this study were represented by an adequate number of reads ($333,537.9 \pm 112,200.5$), with no individual sample receiving fewer than 130,402 reads. Third, there was a very low frequency of cross-contamination from other libraries into those reported here ($5e-005 \pm 8e-05$; max proportion 0.00034). Fourth, after scaling all samples to the same sequencing depth, OTUs with abundance greater than 178 reads (0.14% of a sample's reads) experienced no turnover among PCR replicates within a sample. Fifth, sequence abundances among PCR replicates within water samples were remarkably consistent. A single sample had low similarity among PCR replicates (0.659) after removing this outlier, the lowest mean similarity among replicates within a sample was 0.966. Overall similarities among PCR replicates within a sample were extremely high (0.976 ± 0.013), and far higher than those among samples (0.3 ± 0.16). Across PCR replicates, each sample was represented by at least 781425 reads in the raw data and contained between 111 and 443 rarefied OTUs (Fig. S2).

Ecological analyses

Distance decay

Physical proximity is a good predictor of eDNA community similarity: similarity decreased from 0.40 (95% CI [0.36–0.45]) to half that amount at 4,500 m (95% CI [2,900–7,500]) (Fig. 2).

Community classification

Despite a clear trend in community similarity as a function of spatial separation, the results from our classification analysis are difficult to interpret. The silhouette analysis indicated the presence of 8 distinct communities; however, the gain in mean silhouette width from 2 was small (0.1), and lacked a distinctive peak (Fig. 3), indicating substantial uncertainty in the clustering algorithm. Thus, we present the results of cluster assignment for both $K = 2$ and $K = 8$ to illustrate the range of results (Fig. 4). Excluding taxa which occur in only one site had no discernible effect on the outcome of the PAM analysis (number of clusters, assignment to clusters). While there was no distinct spatial divide indicating the presence of an inshore versus an offshore community, one of the two communities (at $K = 2$) occurred in only two out of 18 samples inside 1,000 m from shore, and never occurred within 125 m of shore, suggesting the presence of an inshore and offshore community.

Diversity in space

Sites offshore tend to be less rich and more even than those inshore (Fig. 5). Mean OTU richness declined by 1.42 per 1,000 m from a mean of 17.6 taxa (95% CI = 2.15) inshore to 11.9 taxa (95% CI = 4.31) at offshore locations ($p = 0.0415$; Fig. 5). Evenness increased by




Figure 2 Distance decay relationship of environmental DNA communities. Each point represents the Bray–Curtis similarity of a site sampled along three parallel transects comprising a 3,000 by 4,000 m grid. Blue dashed line represents fit of a nonlinear least squares regression (see ‘Methods’), and shading denotes the 95% confidence interval. Boxplot is comparisons within-sample across PCR replicates, separated by a vertical line at zero, where the central line is the median, the box encompasses the interquartile range, and the lines extend to 1.5 times the interquartile range. Boxplot outliers are omitted for clarity.




Figure 3 Silhouette widths from PAM analysis. Points are the width of the PAM silhouette of each sample at each number of clusters (K). Red line is the mean, blue line is the median. Boxes encompass the interquartile range with a line at the median, and the whiskers extend to 1.5 times the interquartile range. Boxplot outliers are omitted for clarity.




Figure 4 Cluster membership of sampled sites. Distance from onshore starting point is log scaled. Sites are colored and labeled by their assignment to a cluster by PAM analysis for number of clusters (K) chosen based on a priori expectations (A; 2) and mean silhouette width (B; 8).

.0666 per 1,000 m from 0.225 (95% CI = 0.0558) to 0.491 (95% CI = ± 0.112), indicating that sequence reads were less evenly distributed among taxa in offshore samples ($p \ll 0.05$; Fig. 5). The subset of data used for this analysis had no qualitative effect on the outcome of this analysis (Fig. S3). There was no evidence for spatial autocorrelation for any of the diversity metrics (Moran's I, $p > 0.05$; Fig. 6).

Taxon and life history patterns

We were able to assign a taxon name with confidence to 136 of 146 OTU sequences. The vast majority of sequences (97.6%) and OTUs (96.9%) were matched to organisms that have high potential for dispersal at either the gamete, larval, or adult stage, making it impossible to determine whether the source of that DNA was adults with well-documented spatial patterns (e.g., sessile nearshore specialists) or highly mobile early life history stages. Of the 6 OTUs for which dispersal is limited during all life history stages, only 2 occurred in more than two samples, precluding a quantitative comparison of spatial dispersion based on life history characteristics. These were assigned to *Cymatogaster aggregata*, a viviparous nearshore fish with internal fertilization, and *Cupolaconcha meroclista*, a sessile Vermetid gastropod with presumed internal fertilization and short larval dispersal (Strathmann & Strathmann, 2006; Phillips & Shima, 2010; Calvo & Templado, 2004). *Cymatogaster aggregata* was distinctly more abundant close to shore, with no sequences occurring




Figure 5 Aggregate diversity metrics of each site plotted against distance from shore. Both Simpson's Index of evenness (A) and richness (B) are shown, and have been computed from the mean abundance of unique DNA sequences found across 4 PCR replicates at each of 24 sites. Lines and bands illustrate the fit and 95% confidence interval of a linear model.




Figure 6 Aggregate measures of diversity at each sample site. Data are rarefied counts of mitochondrial 16S sequences collected from three parallel transects in Puget Sound, Washington, USA. Evenness (A) is the probability that two sequences drawn at random are different; richness (B) represents the total number of unique sequences from that location.




Figure 7 Distribution of eDNA from select taxa. Taxa represented are: Embiotocidae (A), *Cupolaconcha meroclista* (B), *Thysanoessa raschii* (C), *Mytilus* (D), Sessilia (E). Circles are colored and scaled by the proportion of that taxon's maximum proportional abundance; the largest circle is the same size in each of the panels, and occurs where that taxon contributed the greatest proportional abundance of reads to that sample.

in any sample beyond 250 m (Fig. 7). *Cupolaconcha meroclista* showed no such distinct spatial trend, occurring in nearly equal abundance at three sites, 75, 500, and 2,000 m from shore. An additional species that was highly abundant in the sequence data, the krill *Thysanoessa raschii*, has pelagic adults, highly seasonal reproduction, and sinking eggs; their distribution was consistent with our expectations based on a tendency of adults to aggregate offshore. Finally, the two most abundant taxa in the dataset were the mussel genus *Mytilus* and the Barnacle order Sessilia; the adults of both taxa are sessile and occur exclusively on hard intertidal substrata but have highly motile larvae. Because large-scale dispersal could not be ruled out for the vast majority of taxa, subsetting the community data by taxonomic group had no qualitative effect on the spatial patterning or diversity metrics, and we omit those results here.

DISCUSSION

Indirect surveys of organismal presence are a key development in ecosystem monitoring in the face of increased anthropogenic pressure and dwindling resources for ecological research. Monitoring of organisms using environmental DNA is an especially promising method, given the rapid pace of advancement in technological innovation and cost efficiency in the field of DNA sequencing and quantification. We document four key patterns: (1) eDNA communities far from one another tend to be less similar than those that are nearby, (2) distinct eDNA communities exist and are distributed in a non-random fashion, (3) diversity declines with distance from shore, and (4) spatial patterning of eDNA is associated with taxon-specific life history characteristics.

Communities far from one another tend to be less similar than those that are nearby

We demonstrate that distant locations have less-similar eDNA communities than proximate locations in Puget Sound, a dynamic marine environment. Our finding is in line with observations based on traditional surveys of terrestrial plants and fungi (*Nekola & White, 1999; Bahram et al., 2013; Condit, 2002; Chust et al., 2006*) and of microorganisms in freshwater (*Wetzel et al., 2012*), marine (*Chust et al., 2013*), and estuarine (*Martiny et al., 2011*) environments. To our knowledge, it is the first to report such a pattern using massively parallel sequencing of environmental DNA in the marine environment, and the first using any technique to describe this pattern from microbial metazoans. We note that the theoretical expectation is that samples at very close distance be nearly completely similar, while our samples separated by the 50 m were only 40% similar. We interpret this to reflect the highly dynamic nature of this environment, which could cause DNA to be distributed quickly from its source, eroding the rise in similarity at small distances. At the same time, community similarity decreased to very low levels at larger scales, indicating that DNA distribution is not completely unpredictable. This finding implies that the effectively sampled area of individual water samples for eDNA analysis is likely to be quite small (<100 m) in this nearshore environment. Our estimated distance-decay relationship does indicate that proximate samples are more similar than distant samples, but we suggest this pattern is partially obscured by other factors, including signal from mobile, microscopic life-stages.

Distinct eDNA communities exist and are distributed in a non-random fashion

We demonstrate strong evidence for distinct community types and the non-random spatial patterning of those communities. While the spatial distributions of communities is surprising if one were concerned only with the macroscopic life stages of metazoans, it indeed does align with the broader view that even offshore pelagic communities are comprised of and influenced by nearshore organisms. This result underscores the idea that areas immediately offshore act as ecotones, a mixing zone of taxa characteristic of benthic and pelagic environments. While there was no distinct break in community types between onshore and offshore sites, there was some clustering of community types that may be explained by oceanographic features such as nearshore eddies generated by strong tidal exchange in a steep bathymetric setting (*Yang & Khangaonkar, 2010*). It would be useful to better understand such features during the period of sampling, by way of oceanographic monitoring devices. Finally, the uncertainty in identification of the number of distinct clusters to best characterize the community underlines the difficulty of identifying community patterns with the number of taxonomic groups considered here. We suspect that the signature of eDNA from microscopic life-stages may explain our inability to easily detect spatial community level patterns that align with our initial expectations.

Richness declines and evenness increases with distance from shore

We found that richness declined while evenness increased with distance from shore. Such a pattern is consistent with many other ecosystems which show strong clines in

diversity metrics over environmental gradients. However, our study is novel in that it corroborates a cline well-known on macroscales for macrobiota on a much smaller spatial scale for microscopic animals, suggesting that there may be a self-similarity across scales in diversity patterning (Levin, 1992). The coastal ocean is a highly productive and diverse ecosystem, where biomass is concentrated most heavily along the bottom and shoreline (Ray, 1988). This differential in biomass concentration from the shoreline to open waters may contribute to the opposing trends we detected. Where particles (organisms, tissues, and cells) are sparse, fewer would be collected per sample of constant volume, thus decreasing the probability of drawing as many types (richness) and increasing the probability that any two particles originate from the same type (evenness). Intriguingly, the cline in diversity from inshore to offshore was not determined by shared changes in communities as one moved offshore; the classification analysis suggested a fair amount of differences among communities at a given offshore distance (Fig. 4).

Spatial patterning of eDNA is associated with taxon-specific life history

In contrast to our expectations, other taxa including species with sessile adult stages restricted to benthic hard substrates (e.g., barnacles, mussels) are among the most abundant taxa at sites furthest from shore. However, the larvae and gametes of these taxa are abundant, pelagic, and can be transported long distances by water movement (Strathmann, 1987). This indicates that we likely detected DNA of their pelagic phase gametes and larvae. It is always possible that DNA of adults was advected over long distances and detected offshore but in light of our results with krill and surfperch, we view this as unlikely. We interpret our results as evidence that the chaotic spatial distribution of eDNA communities (Fig. 4) results from our primers' affinity for many species which at some point exist as microscopic pelagic gametes or larvae. Our results emphasize that expected results based on easily visually observed individuals or detectable with traditional sampling gear such as nets may be very different from results using eDNA. This does caution that eDNA surveys may have different purposes and may not be directly comparable to existing surveys (Shelton et al., 2016).

We acknowledge that sampling artifacts may have affected our results. For example if entire multicellular individuals were captured in our samples, their DNA could be in much greater density than eDNA, affecting the observed community. Our sampling bottles excluded particles larger than 500 micrometers, but gametes and very small larvae could have gained entry. It is possible that even a single small individual, containing many thousand mitochondria, would overwhelm the signal of another species from which hundreds of cells had been sloughed from many, larger individuals. Data on larval size distribution at the time of sampling from each species in our data set would allow us to estimate the frequency of such events. Nevertheless, it is precisely the sensitivity to small particles that makes the eDNA approach powerful, so we are reluctant to recommend that aquatic eDNA sampling use finer pre-filtering. Instead, we emphasize the importance of designing and selecting primer sets that selectively amplify target organisms. In the case of the present study, in order to recover patterns matching our expectations, this would be non-transient, benthic marine organisms lacking any pelagic life stage.

The marker we chose for this study detects a wide variety of metazoans while excluding other more common taxa; however, it does not effectively discriminate among species within a higher group in all cases. Other markers, such as mitochondrial cytochrome c oxidase subunit 1 (COX1, COI, or COI) may provide adequate species-level resolution in some metazoan groups, but have other shortcomings including taxon dropout ([Deagle et al., 2014](#)) and amplification of more abundant non-metazoans, as we discovered in an accompanying study ([Kelly et al., 2016a](#)). Both have undesirable effects of biasing estimates of diversity. In our case, it is possible that the lumping of multiple species into one group underestimates the true richness of the group and of the entire sample, in turn obscuring true underlying patterns of diversity. In the case of COX1, well-documented primer biases cause failure to amplify some taxa, particularly in mixed samples, with the same result ([Deagle et al., 2014](#)). In fact, even surveys relying on traditional capture techniques (e.g., seine nets) and morphological characteristics are subject to biases imposed by the sampling gear (e.g., mesh size), the observer (e.g., taxonomic expertise), and organisms (e.g., morphologically cryptic species). Similarly, no single molecular marker adequately and effectively samples all taxa without bias ([Drummond et al., 2015](#)), and thus the choice of marker is an important and context-dependent one. Until whole-genome sequencing of individual cells is a reality, the tradeoffs between taxonomic breadth and resolution will continue to be problematic for metabarcoding studies, just as they are for more traditional ecological survey methods ([Kelly et al., 2016a](#)).

Our results also highlight the need for curated life-history databases. As technological advances increase the speed and throughput of DNA sequencing and sequence processing, making sense of these data in a timely manner requires that natural history data be stored in standard formats in centralized repositories. The rate at which we can make sense of high-throughput survey methods will be limited by our ability to collate auxiliary data. Databases such as Global Biodiversity Information Facility (GBIF), Encyclopedia of Life (EOL), and FishBase ([Parr et al., 2014](#); [Froese & Pauly, 2016](#)) contain records of taxonomy, occurrence, and other rudimentary data types, but there is no centralized, standardized repository for even basic natural history data such as body size. As NCBI's nucleotide and protein sequence database (GenBank) has facilitated transformative studies in diverse fields, an ecological analog would be a boon for biodiversity science.

Surveys based on eDNA are intensely scrutinized because of the danger that the final data are subject to complicated laboratory and bioinformatic procedures. Finding virtually no variability among lab and bioinformatic treatments from the point of PCR onward, we were confident our results represented actual field-based differences among samples. However, we note that one PCR replicate had a clear signal of contamination in that the sequence community was extremely similar to those from a different environmental sample. The source of this error is difficult to identify, but seems most likely to be an error during PCR preparation, either in assignment or pipetting during preparation of indexed primers. While the remainder of our results would be largely unchanged had we sequenced a single replicate per environmental sample, we believe the sequencing of PCR replicates is critical for ensuring data quality in eDNA sequencing studies.

While there is much work to be done before eDNA techniques can be confidently deployed as a standard method for ecological monitoring, this study serves as a first analysis of diversity at the fine spatial scales that are likely to be relevant to eDNA work in the field across a range of study systems.

ACKNOWLEDGEMENTS

We wish to thank Linda Park, Robert Morris, E. Virginia Armbrust, and James Kralj. The manuscript was improved by suggestions from editor Magnus Johnson, and reviewers Owen Wangenstein and Stephen Moss.

ADDITIONAL INFORMATION AND DECLARATIONS

Funding

This work was supported by a grant from the David and Lucile Packard Foundation to RPK (grant 2014-39827). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Grant Disclosures

The following grant information was disclosed by the authors:
David and Lucile Packard Foundation: 2014-39827.

Competing Interests

The authors declare there are no competing interests.

Author Contributions

- James L. O'Donnell conceived and designed the experiments, performed the experiments, analyzed the data, wrote the paper, prepared figures and/or tables, reviewed drafts of the paper.
- Ryan P. Kelly and Andrew Olaf Shelton conceived and designed the experiments, performed the experiments, reviewed drafts of the paper.
- Jameal F. Samhouri, Natalie C. Lowell and Gregory D. Williams performed the experiments, reviewed drafts of the paper.

DNA Deposition

The following information was supplied regarding the deposition of DNA sequences:
All sequence data are available from NCBI under BioProject [PRJNA338801](https://www.ncbi.nlm.nih.gov/bioproject/PRJNA338801).

Data Availability

The following information was supplied regarding data availability:
Analysis code and data (except sequence data) is available at: <https://doi.org/10.5281/zenodo.242976>.

Supplemental Information

Supplemental information for this article can be found online at <http://dx.doi.org/10.7717/peerj.3044#supplemental-information>.

REFERENCES

- Anderson MJ, Crist TO, Chase JM, Vellend M, Inouye BD, Freestone AL, Sanders NJ, Cornell HV, Comita LS, Davies KF, Harrison SP, Kraft NJB, Stegen JC, Swenson NG. 2011. Navigating the multiple meanings of beta diversity: a roadmap for the practicing ecologist. *Ecology Letters* 14(1):19–28
DOI 10.1111/j.1461-0248.2010.01552.x.
- Bahram M, Kõljalg U, Courty PE, Diédhiou AG, Kjølner R, Põlme S, Ryberg M, Veldre V, Tedersoo L. 2013. The distance decay of similarity in communities of ectomycorrhizal fungi in different ecosystems and scales. *Journal of Ecology* 101(5):1335–1344
DOI 10.1111/1365-2745.12120.
- Barnes MA, Turner CR, Jerde CL, Renshaw MA, Chadderton WL, Lodge DM. 2014. Environmental conditions influence eDNA persistence in aquatic systems. *Environmental Science and Technology* 48(3):1819–1827 DOI 10.1021/es404734p.
- Bell T. 2010. Experimental tests of the bacterial distance–decay relationship. *The ISME Journal* 4(11):1357–1365 DOI 10.1038/ismej.2010.77.
- Burns RE. 1985. The shape and form of Puget Sound. 1st edition. Seattle: Washington Sea Grant, 100.
- Calvo M, Templado J. 2004. Reproduction and development in a vermetid gastropod, *Vermetus triquetrus*. *Invertebrate Biology* 123(4):289–303
DOI 10.1111/j.1744-7410.2004.tb00163.x.
- Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K, Madden TL. 2009. BLAST+: architecture and applications. *BMC Bioinformatics* 10:421
DOI 10.1186/1471-2105-10-421.
- Chamberlain SA, Szöcs E. 2013. Taxize: taxonomic search and retrieval in R. *F1000Research* 2(0):Article 191 DOI 10.12688/f1000research.2-191.v2.
- Chamberlain SA, Szöcs E, Boettiger C, Ram K, Bartomeus I, Foster Z, O'Donnell JL. 2016. taxize: taxonomic information from around the web. R package. Available at <https://github.com/ropensci/taxize>.
- Chust G, Chave J, Condit R, Aguilar S, Lao S, Perez R. 2006. Determinants and spatial modeling of tree beta-diversity in a tropical forest landscape in Panama. *Journal of Vegetation Science* 17(1):83–92 DOI 10.1111/j.1654-1103.2006.tb02426.x.
- Chust G, Irigoien X, Chave J, Harris RP. 2013. Latitudinal phytoplankton distribution and the neutral theory of biodiversity. *Global Ecology and Biogeography* 22(5):531–543 DOI 10.1111/geb.12016.
- Coissac E. 2012. OligoTag: a program for designing sets of tags for next-generation sequencing of multiplexed samples. In: Pompanon F, Bonin A, eds. *Data production and analysis in population genomics SE—2. Methods in molecular biology*, vol. 888. New York: Humana Press, 13–31.
- Condit R. 2002. Beta-diversity in tropical forest trees. *Science* 295(5555):666–669
DOI 10.1126/science.1066854.

- Cowart DA, Pinheiro M, Mouchel O, Maguer M, Grall J, Miné J, Arnaud-Haond S. 2015. Metabarcoding is powerful yet still blind: a comparative analysis of morphological and molecular surveys of seagrass communities. *PLOS ONE* 10(2):e0117562 DOI 10.1371/journal.pone.0117562.
- Deagle BE, Jarman SN, Coissac E, Pompanon F, Taberlet P. 2014. DNA metabarcoding and the cytochrome c oxidase subunit I marker: not a perfect match. *Biology Letters* 10(9):1789–1793 DOI 10.1098/rsbl.2014.0562.
- Deiner K, Altermatt F. 2014. Transport distance of invertebrate environmental DNA in a natural river. *PLOS ONE* 9(2):e88786 DOI 10.1371/journal.pone.0088786.
- Dethier MN. 2010. Overview of the ecology of Puget Sound beaches. In: Shipman H, Dethier MN, Gelfenbaum G, Fresh KL, Dinicola RS, eds. *Puget Sound shorelines and the impacts of armoring—proceedings of a state of the science workshop*. 262.
- De Vargas C, Audic S, Henry N, Decelle J, Mahé F, Logares R, Lara E, Berney C, Le Bescot N, Probert I, Carmichael M, Poulain J, Romac S, Colin S, Aury J-M, Bittner L, Chaffron S, Dunthorn M, Engelen S, Flegontova O, Guidi L, Horák A, Jaillon O, Lima-Mendez G, Lukeš J, Malviya S, Morard R, Mulot M, Scalco E, Siano R, Vincent F, Zingone A, Dimier C, Picheral M, Searson S, Kandels-Lewis S, Acinas SG, Bork P, Bowler C, Gorsky G, Grimsley N, Hingamp P, Iudicone D, Not F, Ogata H, Pesant S, Raes J, Sieracki ME, Speich S, Stemmann L, Sunagawa S, Weissenbach J, Wincker P, Karsenti E. 2015. Eukaryotic plankton diversity in the sunlit ocean. *Science* 348(6237):1261605 DOI 10.1126/science.1261605.
- Drummond AJ, Newcomb RD, Buckley TR, Xie D, Dopheide A, Potter BC, Heled J, Ross HA, Tooman L, Grosser S, Park D, Demetras NJ, Stevens MI, Russell JC, Anderson SH, Carter A, Nelson N. 2015. Evaluating a multigene environmental DNA approach for biodiversity assessment. *GigaScience* 4(1):Article 46 DOI 10.1186/s13742-015-0086-1.
- Edgar RC. 2010. Search and clustering orders of magnitude faster than BLAST. *Bioinformatics* 26(19):2460–2461 DOI 10.1093/bioinformatics/btq461.
- Evans NT, Olds BP, Renshaw MA, Turner CR, Li Y, Jerde CL, Mahon AR, Pfrender ME, Lamberti GA, Lodge DM. 2016. Quantification of mesocosm fish and amphibian species diversity via environmental DNA metabarcoding. *Molecular Ecology Resources* 16(1):29–41 DOI 10.1111/1755-0998.12433.
- Ficetola GF, Miaud C, Pompanon F, Taberlet P. 2008. Species detection using environmental DNA from water samples. *Biology Letters* 4(4):423–425 DOI 10.1098/rsbl.2008.0118.
- Fonseca VG, Carvalho GR, Nichols B, Quince C, Johnson HF, Neill SP, Lambshead JD, Thomas WK, Power DM, Creer S. 2014. Metagenetic analysis of patterns of distribution and diversity of marine meiobenthic eukaryotes. *Global Ecology and Biogeography* 23(11):1293–1302 DOI 10.1111/geb.12223.
- Froese R, Pauly D. 2016. FishBase. Available at www.fishbase.org.
- Guardiola M, Uriz MJ, Taberlet P, Coissac E, Wangensteen OS, Turon X. 2015. Deep-sea, deep-sequencing: metabarcoding extracellular DNA from sediments of marine canyons. *PLOS ONE* 10(10):e0139633 DOI 10.1371/journal.pone.0139633.

- Guardiola M, Wangensteen OS, Taberlet P, Coissac E, Uriz MJ, Turon X. 2016.** Spatio-temporal monitoring of deep-sea communities using metabarcoding of sediment DNA and RNA. *PeerJ* 4:e2807 DOI 10.7717/peerj.2807.
- Guo H, Chamberlain SA, Elhaik E, Jalli I, Lynes AR, Marczak L, Sabath N, Vargas A, Więski K, Zelig EM, Pennings SC. 2015.** Geographic variation in plant community structure of salt marshes: species, functional and phylogenetic perspectives. *PLOS ONE* 10(5):e0127781 DOI 10.1371/journal.pone.0127781.
- Handelsman J, Rondon MR, Brady SF, Clardy J, Goodman RM. 1998.** Molecular biological access to the chemistry of unknown soil microbes: a new frontier for natural products. *Chemistry & Biology* 5(10):R245–R249 DOI 10.1016/S1074-5521(98)90108-9.
- Hijmans RJ. 2016.** geosphere: spherical trigonometry. Available at <https://cran.r-project.org/package=geosphere>.
- Hubbell S. 2001.** The unified neutral theory of biodiversity and biogeography. In: *Monographs in population biology*, vol. 32. 375 pp.
- Iverson V, Morris RM, Frazar CD, Berthiaume CT, Morales RL, Armbrust EV. 2012.** Untangling genomes from metagenomes: revealing an uncultured class of marine euryarchaeota. *Science* 335(6068):587–590 DOI 10.1126/science.1212665.
- Kaufman L, Rousseeuw PJ. 1990.** Finding groups in data: an introduction to cluster analysis. In: *Wiley Series in Probability and Statistics*. Hoboken: Wiley, 342.
- Kelly RP, Closek CJ, O'Donnell JL, Kralj JE, Shelton AO, Samhouri JF. 2016a.** Genetic and manual survey methods yield different and complementary views of an ecosystem. *Frontiers in Marine Science* 3:Article 283 DOI 10.3389/FMARS.2016.00283.
- Kelly RP, O'Donnell JL, Lowell NC, Shelton AO, Samhouri JF, Hennessey SM, Feist BE, Williams GD. 2016b.** Genetic signatures of ecological diversity along an urbanization gradient. *PeerJ* 4:e2444 DOI 10.7717/peerj.2444.
- Klymus KE, Richter CA, Chapman DC, Paukert C. 2015.** Quantification of eDNA shedding rates from invasive bighead carp *Hypophthalmichthys nobilis* and silver carp *Hypophthalmichthys molitrix*. *Biological Conservation* 183:77–84 DOI 10.1016/j.biocon.2014.11.020.
- Kozloff EN. 1973.** Seashore life of Puget Sound, the Strait of Georgia, and the San Juan Archipelago. Seattle: University of Washington Press, 282.
- Lallias D, Hiddink JG, Fonseca VG, Gaspar JM, Sung W, Neill SP, Barnes N, Ferrero T, Hall N, Lambshead PJD, Packer M, Thomas WK, Creer S. 2015.** Environmental metabarcoding reveals heterogeneous drivers of microbial eukaryote diversity in contrasting estuarine ecosystems. *The ISME Journal* 9(5):1208–1221 DOI 10.1038/ismej.2014.213.
- Levin SA. 1992.** The problem of pattern and scale in ecology. *Ecology* 73(6):1943–1967 DOI 10.2307/1941447.
- Longmire JL, Maltbie M, Baker RJ. 1997.** Use of lysis buffer in DNA isolation and its implication for museum collections. In: *Occasional papers Museum of Texas Tech University* 163. Lubbock: Texas Tech University.

- Maechler M, Rousseeuw P, Struyf A, Hubert M, Hornik K. 2016.** Cluster: cluster analysis basics and extensions. R package version 2.0.5. Available at <https://cran.r-project.org/web/packages/cluster/>.
- Magurran AE. 2004.** Measuring biological diversity. Oxford: Blackwell Science, Ltd, 264.
- Mahé F, Rognes T, Quince C, De Vargas C, Dunthorn M. 2014.** Swarm: robust and fast clustering method for amplicon-based studies. *PeerJ* 2:e593 DOI 10.7717/peerj.593.
- Martin M. 2011.** Cutadapt removes adapter sequences from high-throughput sequencing reads. *EMBnet.Journal* 17(1):10–12 DOI 10.14806/ej.17.1.200.
- Martiny JBH, Eisen JA, Penn K, Allison SD, Horner-Devine MC. 2011.** Drivers of bacterial beta diversity depend on spatial scale. *Proceedings of the National Academy of Sciences of the United States of America* 108(19):7850–7854 DOI 10.1073/pnas.1016308108.
- Nekola JC, White PS. 1999.** The distance decay of similarity in biogeography and ecology. *Journal of Biogeography* 26(4):867–878 DOI 10.1046/j.1365-2699.1999.00305.x.
- O'Donnell JL, Kelly RP, Lowell NC, Port JA. 2016.** Indexed PCR primers induce template-specific bias in large-scale DNA sequencing studies. *PLOS ONE* 11(3):e0148698.
- Oksanen J, Blanchet FG, Friendly M, Kindt R, Legendre P, McGlinn D, Minchin PR, O'Hara RB, Simpson GL, Solymos P, Stevens MHH, Szoecs E, Wagner H. 2016.** Vegan: community ecology package. Available at <https://cran.r-project.org/package=vegan>.
- Parr CS, Wilson N, Leary P, Schulz KS, Lans K, Walley L, Hammock JA, Goddard A, Rice J, Studer M, Holmes JTG, Corrigan RJ. 2014.** The encyclopedia of life v2: providing global access to knowledge about life on earth. *Biodiversity Data Journal* 2(2):e1079 DOI 10.3897/BDJ.2.e1079.
- Phillips NE, Shima JS. 2010.** Reproduction of the vermetid gastropod dendropoma maximum (Sowerby, 1825) in Moorea, French Polynesia. *Journal of Molluscan Studies* 76(2):133–137 DOI 10.1093/mollus/eyp049.
- Port JA, O'Donnell JL, Romero-Maraccini OC, Leary PR, Litvin SY, Nickols KJ, Yamahara KM, Kelly RP. 2016.** Assessing vertebrate biodiversity in a kelp forest ecosystem using environmental DNA. *Molecular Ecology* 25(2):527–541 DOI 10.1111/mec.13481.
- R Core Team. 2016.** R: a language and environment for statistical computing. Vienna: R Foundation for Statistical Computing. Available at <https://www.r-project.org/>.
- Ray GC. 1988.** Ecological diversity in coastal zones and oceans. In: Wilson EO, Peter FM, eds. *Biodiversity*. Washington, D.C.: National Academies Press (US).
- Renshaw MA, Olds BP, Jerde CL, McVeigh MM, Lodge DM. 2015.** The room temperature preservation of filtered environmental DNA samples and assimilation into a phenol–chloroform–isoamyl alcohol DNA extraction. *Molecular Ecology Resources* 15(1):168–176 DOI 10.1111/1755-0998.12281.
- Rognes T, Flouri T, Nichols B, Quince C, Mahé F. 2016.** VSEARCH: a versatile open source tool for metagenomics. *PeerJ* 4:e2584 DOI 10.7717/peerj.2584.

- Sassoubre LM, Yamahara KM, Gardner LD, Block BA, Boehm AB. 2016.** Quantification of environmental DNA (eDNA) shedding and decay rates for three marine fish. *Environmental Science & Technology* **50**(19):10456–10464 DOI [10.1021/acs.est.6b03114](https://doi.org/10.1021/acs.est.6b03114).
- Schnell IB, Bohmann K, Gilbert MTP. 2015.** Tag jumps illuminated—reducing sequence-to-sample misidentifications in metabarcoding studies. *Molecular Ecology Resources* **15**(6):1289–1303 DOI [10.1111/1755-0998.12402](https://doi.org/10.1111/1755-0998.12402).
- Shelton AO, O'Donnell JL, Samhouri JF, Lowell N, Williams GD, Kelly RP. 2016.** A framework for inferring biological communities from environmental DNA. *Ecological Applications* **26**(6):1645–1659 DOI [10.1890/15-1733.1](https://doi.org/10.1890/15-1733.1).
- Shogren AJ, Tank JL, Andruszkiewicz EA, Olds B, Jerde C, Bolster D. 2016.** Modelling the transport of environmental DNA through a porous substrate using continuous flow-through column experiments. *Journal of the Royal Society Interface* **13**(119):423–425 DOI [10.1098/rsif.2016.0290](https://doi.org/10.1098/rsif.2016.0290).
- Simpson EH. 1949.** Measurement of diversity. *Nature* **163**:688.
- Spiess A-N. 2014.** propagate: propagation of uncertainty. Available at <https://cran.r-project.org/package=propagate>.
- Strathmann MF. 1987.** Reproduction and development of marine invertebrates of the Northern Pacific Coast: data and methods for the study of eggs, embryos, and larvae. Seattle: University of Washington Press, 682.
- Strathmann MF, Strathmann RR. 2006.** A vermetid gastropod with complex intracapsular cannibalism of nurse eggs and sibling larvae and a high potential for invasion. *Pacific Science* **60**(1):97–108 DOI [10.1353/psc.2005.0062](https://doi.org/10.1353/psc.2005.0062).
- Strickland RM. 1983.** The fertile fjord: plankton in Puget Sound. Seattle: University of Washington Press, 160.
- Strickler KM, Fremier AK, Goldberg CS. 2015.** Quantifying effects of UV-B, temperature, and pH on eDNA degradation in aquatic microcosms. *Biological Conservation* **183**:85–92 DOI [10.1016/j.biocon.2014.11.038](https://doi.org/10.1016/j.biocon.2014.11.038).
- Thomsen PF, Kielgast J, Iversen LL, Møller PR, Rasmussen M, Willerslev E. 2012.** Detection of a diverse marine fish fauna using environmental DNA from seawater samples. *PLOS ONE* **7**(8):1–9 DOI [10.1371/journal.pone.0041732](https://doi.org/10.1371/journal.pone.0041732).
- Turner CR, Uy KL, Everhart RC. 2015.** Fish environmental DNA is more concentrated in aquatic sediments than surface water. *Biological Conservation* **183**:93–102 DOI [10.1016/j.biocon.2014.11.017](https://doi.org/10.1016/j.biocon.2014.11.017).
- Tyson GW, Chapman J, Hugenholtz P, Allen EE, Ram RJ, Richardson PM, Solovyev VV, Rubin EM, Rokhsar DS, Banfield JF. 2004.** Community structure and metabolism through reconstruction of microbial genomes from the environment. *Nature* **428**(6978):37–43 DOI [10.1038/nature02340](https://doi.org/10.1038/nature02340).
- Venter JC, Remington K, Heidelberg JF, Halpern AL, Rusch D, Eisen JA, Wu D, Paulsen I, Nelson KE, Nelson W, Fouts DE, Levy S, Knap AH, Lomas MW, Nealson K, White O, Peterson J, Hoffman J, Parsons R, Baden-Tillson H, Pfannkoch C, Rogers Y-H, Smith HO. 2004.** Environmental genome shotgun sequencing of the Sargasso Sea. *Science* **304**(5667):66–74 DOI [10.1126/science.1093857](https://doi.org/10.1126/science.1093857).

- Wetzel CE, De Bicudo DC, Ector L, Lobo EA, Soininen J, Landeiro VL, Bini LM. 2012.** Distance decay of similarity in neotropical diatom communities. *PLOS ONE* 7(9):e45071 DOI [10.1371/journal.pone.0045071](https://doi.org/10.1371/journal.pone.0045071).
- Yang Z, Khangaonkar T. 2010.** Multi-scale modeling of Puget Sound using an unstructured-grid coastal ocean model: from tide flats to estuaries and coastal waters. *Ocean Dynamics* 60(6):1621–1637 DOI [10.1007/s10236-010-0348-5](https://doi.org/10.1007/s10236-010-0348-5).
- Zhang J, Kobert K, Flouri T, Stamatakis A. 2014.** PEAR: a fast and accurate illumina paired-end reAd mergeR. *Bioinformatics* 30(5):614–620 DOI [10.1093/bioinformatics/btt593](https://doi.org/10.1093/bioinformatics/btt593).