

Meeting abstract

Open Access

## The G protein-coupled receptor identity of the frizzled proteins

VL Katanaev\* and S Buestorf

Address: University of Konstanz, Department of Biology, Konstanz, Switzerland

\* Corresponding author

from 12th Joint Meeting of the Signal Transduction Society (STS). Signal Transduction: Receptors, Mediators and Genes Weimar, Germany. 29–31 October 2008

Published: 26 February 2009

*Cell Communication and Signaling* 2009, **7**(Suppl 1):A19 doi:10.1186/1478-811X-7-S1-A19

This abstract is available from: <http://www.biosignaling.com/content/7/S1/A19>

© 2009 Katanaev and Buestorf; licensee BioMed Central Ltd.

Receptors of the Frizzled family initiate Wnt ligand-dependent signal transduction cascades controlling multiple steps in organism development and are highly conserved in animal evolution. Misactivation of the Wnt/Frizzled signaling underlies many cases of cancerogenesis. Frizzled receptors possess seven transmembrane domains and their signaling depends on trimeric G proteins in various organisms. However, as Frizzled proteins constitute a distinct group within the superfamily of G protein-coupled receptors (GPCR), and as Frizzled signaling can apparently be G protein-independent in some experimental setups, the GPCR nature of Frizzled receptors has been questioned. Here we demonstrate that human Frizzled receptors can directly bind the trimeric Go protein in a pertussis toxin-sensitive manner. Furthermore, addition of Wnt ligands elicits Frizzled-dependent guanine nucleotide exchange on Go. An excess of secreted Frizzled-related protein, a known antagonist of the Wnt/Frizzled pathways, inhibits Go activation, as does pretreatment of Go with pertussis toxin. These experiments provide a biochemical proof of the GPCR activities of Frizzled receptors. They also establish an in vitro assay of monitoring Frizzled activation by Wnt ligands, applicable for the high-throughput agonist/antagonist screening.