

RESEARCH

Open Access



A note on Cauchy-Lipschitz-Picard theorem

Zonghong Feng¹, Fengying Li^{2*}, Ying Lv³ and Shiqing Zhang¹*Correspondence:
lify0308@163.com²School of Economic Mathematics,
Southwestern University of Finance
and Economics, Chengdu, 61130,
ChinaFull list of author information is
available at the end of the article**Abstract**

In this note, we try to generalize the classical Cauchy-Lipschitz-Picard theorem on the global existence and uniqueness for the Cauchy initial value problem of the ordinary differential equation with global Lipschitz condition, and we try to weaken the global Lipschitz condition. We can also get the global existence and uniqueness.

MSC: 34A34; 34C25; 34C37**Keywords:** initial value problems; generalized Lipschitz condition; global existence and uniqueness

1 Introduction

In his famous book [1], Brezis gave a very sketchy and interesting proof on the classical Cauchy-Lipschitz-Picard theorem.

Theorem 1.1 *Let E be a Banach space and let $F : E \rightarrow E$ be a Lipschitz map, i.e., there is a constant L such that*

$$\|F(u) - F(v)\| \leq L\|u - v\|, \quad \forall u, v \in E. \quad (1.1)$$

Then, for any given $u_0 \in E$, there exists a unique solution $u \in C^1([0, +\infty), E)$ of the problem:

$$\begin{cases} \frac{du(t)}{dt} = F(u(t)), & \forall t \in (0, +\infty), \\ u(0) = u_0. \end{cases} \quad (1.2)$$

It is well known that if we only assume the local Lipschitz condition, we can only get the local existence and uniqueness for Cauchy initial value problems.

In this paper, we try to weaken the global Lipschitz condition, but we also want to get the global existence and uniqueness; we have the following theorem.

Theorem 1.2 *Let E be a Banach space (with norm $\|\cdot\|$) and let $F : [0, +\infty) \times E \rightarrow E$ be a map satisfying*

$$\|F(t, u) - F(t, v)\| \leq L(t) \cdot p(\|u - v\|), \quad \forall t \in [0, +\infty), u, v \in E,$$

where $L : [0, +\infty) \rightarrow [0, +\infty)$, $p : (0, +\infty) \rightarrow [0, +\infty)$ are continuous and there are $0 \leq a < +\infty$ such that

$$L = L(t) \leq a,$$

$$p = p(s) \leq s,$$

and $p(s)$ is an increasing function. Then, for any given $u_0 \in E$, there exists a unique solution $u \in C^1([0, +\infty); E)$ for the problem

$$\begin{cases} \frac{du(t)}{dt} = F(t, u(t)), & \forall t \in (0, +\infty), \\ u(0) = u_0. \end{cases} \tag{1.3}$$

Corollary 1.3 *In Theorem 1.2, if we take $L(s) = a > 0$, $p(s) = s$ or $p(s) = \ln s$, then the conditions and the results of Theorem 1.2 hold.*

2 The proof of Theorem 1.2

Lemma 2.1 ([1], Banach contraction mapping principle) *Let X be a nonempty complete metric space and let $T : X \rightarrow X$ be a strict contraction, i.e., there is $0 < k < 1$ such that $d(T(x), T(y)) \leq kd(x, y)$, $\forall x, y \in X$, then S has a unique fixed point $u = T(u)$.*

Lemma 2.2 (Gronwall [2]) *Let $x \in C^1[a, b]$. If $R[a, b]$ denotes the set of Riemann integrable functional on $[a, b]$; $\beta \in R[a, b]$, and*

$$x'(t) \leq \beta(t) \cdot x(t), \quad \forall t \in [a, b],$$

then

$$x(t) \leq x(a) \cdot e^{\int_a^t \beta(s) ds}, \quad \forall t \in [a, b].$$

The following lemma can be regarded as a natural generalization of the Gronwall inequality.

Lemma 2.3 *Let $x \in C^1[a, b]$, $\alpha, \beta \in R[a, b]$. If*

$$x'(t) \leq \alpha(t) + \beta(t)x(t), \quad \forall t \in [a, b],$$

then

$$x(t) \leq \left[x(a) + \int_a^t e^{-\int_0^s \beta(\tau) d\tau} \alpha(s) ds \right] \cdot e^{\int_a^t \beta(s) ds}, \quad \forall t \in [a, b].$$

Proof Let $v = e^{\int_a^t \beta(s) ds}$, $w = \frac{x}{v}$, then $v'(t) = \beta(t)v(t)$,

$$w'(t) = \frac{x'v - xv'}{v^2} \leq \frac{(\alpha + \beta x)v - \beta xv}{v^2} = \frac{\alpha v}{v^2} = \frac{\alpha}{v}.$$

Then

$$w(t) \leq \int_a^t e^{-\int_0^s \beta(\tau) d\tau} \alpha(s) ds + w(a),$$

$$\frac{x(t)}{v(t)} \leq x(a) + \int_a^t e^{-\int_0^s \beta(\tau) d\tau} \alpha(s) ds,$$

$$x(t) \leq e^{\int_a^t \beta(s) ds} \cdot \left[x(a) + \int_a^t e^{-\int_0^s \beta(\tau) d\tau} \alpha(s) ds \right].$$

Now to prove Theorem 1.2, we use some similar arguments to Brezis [1]. Let $k > 0$, which is to be determined, and assume

$$X = \left\{ u \in C([0, +\infty); E) \mid \sup_{t \geq 0} e^{-kt} \cdot \|u(t)\| < +\infty \right\}.$$

Then it is easy to see that X is a Banach space for the norm

$$\|u\|_X = \sup_{t \geq 0} e^{-kt} \cdot \|u(t)\|.$$

For $\forall u \in X$, we define

$$\Phi(u)(t) = u_0 + \int_0^t F(s, u(s)) ds.$$

Then u is a solution of (1.3) if and only if $\Phi(u) = u$, that is, u is a fixed point of Φ .

(1) We now show that, for every $u \in X$, $\Phi(u)$ also belongs to X .

In fact,

$$\begin{aligned} \|\Phi u\|_X &= \sup_{t \geq 0} e^{-kt} \|\Phi(u)(t)\| \\ &\leq \sup_{t \geq 0} e^{-kt} \cdot \|u_0\| + \sup_{t \geq 0} e^{-kt} \cdot \left\| \int_0^t F(s, u(s)) ds \right\|. \end{aligned}$$

We only need to prove

$$\sup_{t \geq 0} e^{-kt} \cdot \int_0^t \|F(s, u(s))\| ds < +\infty.$$

Notice that

$$\begin{aligned} \|F(s, u(s)) - F(0, u_0)\| &\leq L(s) \cdot p(\|u(s) - u_0\|), \\ \|F(s, u(s))\| &\leq L(s) \cdot p(\|u(s) - u_0\|) + \|F(0, u_0)\|. \end{aligned}$$

Since

$$\sup_{t \geq 0} e^{-kt} \cdot \int_0^t \|F(0, u_0)\| ds = \|F(0, u_0)\| \sup_{t \geq 0} e^{-kt} \cdot t < +\infty.$$

Hence we only need to prove

$$\sup_{t \geq 0} e^{-kt} \cdot \int_0^t L(s)p(\|u(s) - u_0\|) ds < +\infty.$$

Let

$$\varphi(t) = e^{-kt} \int_0^t L(s)p(\|u(s) - u_0\|) ds, \quad \varphi(0) = 0,$$

then

$$\begin{aligned} \varphi(t) &\leq \int_0^t [L(s)e^{ks}]e^{-ks}(\|u(s)\| + \|u_0\|) ds e^{-kt} \\ &\leq \left[\sup_{t \geq 0} e^{-ks} \|u(s)\| + \sup_{t \geq 0} e^{-ks} \|u_0\| \right] \sup_{t \geq 0} \left[\int_0^t L(s)e^{ks} ds e^{-kt} \right] \\ &\leq C_1 C_2 = C_3 < +\infty. \end{aligned}$$

Hence we have proved Φ is a self-mapping from X to X .

(2) We prove the contraction property of Φ . We have

$$\begin{aligned} \|\Phi u - \Phi v\| &\leq \int_0^t \|F(s, u(s)) - F(s, v(s))\| ds \leq \int_0^t L(s)p(\|u(s) - v(s)\|) ds, \\ \|\Phi u - \Phi v\|_X &\leq \sup_{t \geq 0} e^{-kt} \cdot \int_0^t L(s)p(\|u(s) - v(s)\|) ds \\ &\leq \sup_{t \geq 0} \int_0^t L(s)e^{ks} [e^{-ks} \|u(s) - v(s)\|] ds e^{-kt} \\ &\leq \sup_{t \geq 0} [e^{-ks} \|u(s) - v(s)\|] \sup_{t \geq 0} e^{-kt} \int_0^t L(s)e^{ks} ds \\ &\leq \sup_{t \geq 0} e^{-kt} \int_0^t L(s)e^{ks} ds \|u - v\|_X. \end{aligned}$$

Hence we have

$$\|\Phi u - \Phi v\|_X \leq \frac{L}{k} \|u - v\|_X, \quad \forall u, v \in X.$$

We can choose $k > L$, then we use Banach contraction mapping principle to find that (1.3) has at least one solution on $[0, +\infty)$.

Furthermore, by Gronwall’s inequality, we can get the uniqueness.

In fact, let u_1, u_2 be two solutions of (1.3), then

$$\begin{aligned} \|u_1(t) - u_2(t)\| &\leq \int_0^t \|F(s, u_1(s)) - F(s, u_2(s))\| ds \\ &\leq \int_0^t L(s)p(\|u_1(s) - u_2(s)\|) ds \\ &\leq \int_0^t L(s)(\|u_1(s) - u_2(s)\|) ds. \end{aligned}$$

By Gronwall’s inequality, we have

$$\|u_1(t) - u_2(t)\| \leq 0.$$

Hence for any $t \in [0, +\infty)$, we have $u_1(t) = u_2(t)$. □

3 Examples

Example 3.1

$$\begin{cases} \frac{du(t)}{dt} = F(t, u(t)) = \frac{1}{1+t+|u|}, & \forall t \in (0, +\infty), \\ u(0) = u_0. \end{cases} \tag{3.1}$$

Then

$$|F(t, u(t)) - F(t, v(t))| = \frac{\|v\| - \|u\|}{(1+t+|u|)(1+t+|v|)} \leq \|v\| - \|u\|.$$

By the triangle inequality, we have

$$|F(t, u(t)) - F(t, v(t))| \leq |u - v|.$$

So $F(t, u(t)) = \frac{1}{1+t+|u|}$ is Lipschitz with $a = 1$ and $p(s) = s$.

Example 3.2 Let $p_1(t) : [0, +\infty) \rightarrow [0, +\infty)$ continuous and there is $0 \leq a < +\infty$ such that

$$p_1(t) \leq a.$$

Let $p_2(t) : [0, +\infty) \rightarrow R$ continuous.

We consider

$$\begin{cases} \frac{du(t)}{dt} = F(t, u(t)) = p_1(t)|u - p_2(t)|, & \forall t \in (0, +\infty), \\ u(0) = u_0. \end{cases} \tag{3.2}$$

Then

$$|F(t, u(t)) - F(t, v(t))| = p_1(t)(\|u - p_2(t)\| - \|v - p_2(t)\|).$$

By the triangle inequality, we have

$$|F(t, u(t)) - F(t, v(t))| \leq p_1(t)|u - v|.$$

So $F(t, u(t)) = p_1(t)|u - p_2(t)|$ is Lipschitz with $L(t) = p_1(t)$ and $p(s) = s$.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

The research and writing of this manuscript was a collaborative effort from all the authors. All authors read and approved the final manuscript.

Author details

¹Department of Mathematics, Sichuan University, Chengdu, 610064, China. ²School of Economic Mathematics, Southwestern University of Finance and Economics, Chengdu, 61130, China. ³Department of Mathematics and Statistics, Southwest University, Chongqing, 400715, China.

Acknowledgements

The authors sincerely thank referee for his/her valuable comments. This paper was partially supported by NSFC (No. 11671278 and No. 11426181) and the National Research Foundation for the Doctoral Program of Higher Education of China under Grant No. 20120181110060.

Received: 6 July 2016 Accepted: 18 October 2016 Published online: 04 November 2016

References

1. Brezis, H: Functional Analysis, Sobolev Spaces and PDE, pp. 184-185. Springer, New York (2011)
2. Thomas, HG: Note on the derivatives with respect to a parameter of the solutions of a system of differential equations. *Ann. Math.* **20**, 292-296 (1919)

Submit your manuscript to a SpringerOpen[®] journal and benefit from:

- Convenient online submission
- Rigorous peer review
- Immediate publication on acceptance
- Open access: articles freely available online
- High visibility within the field
- Retaining the copyright to your article

Submit your next manuscript at ► springeropen.com
