

RESEARCH

Open Access

'*Candidatus Rickettsia asemoensis*' and *Wolbachia* spp. in *Ctenocephalides felis* and *Pulex irritans* fleas removed from dogs in Ecuador

José A Oteo^{1*}, Aránzazu Portillo¹, Francisco Portero², Jorge Zavala-Castro³, José M Venzal⁴ and Marcelo B Labruna⁵

Abstract

Background: Flea-borne infections are distributed worldwide. Up to date there are no reports about microorganisms associated to fleas in Ecuador.

Methods: Seventy-one *Pulex irritans* and 8 *Ctenocephalides felis* fleas were removed from dogs in two Ecuadorian areas (Pastaza and Chimborazo Provinces) in December 2012. DNA extracts were tested by polymerase chain reaction (PCR) assays targeting universal 16S rRNA, as well as screened for the presence of *Rickettsia* spp. (*gltA*, *htrA*, *ompB*, *sca4* and *ompA* genes) and *Bartonella* spp. (*rpoB*, *gltA* and ITS genes).

Results: Our results showed the presence of '*Candidatus Rickettsia asemoensis*' (highly similar to *R. felis*) in *C. felis* and *Wolbachia* spp. endosymbionts in *P. irritans* collected from animals in Ecuador. No fleas were found to be positive for any *Bartonella* species or *Yersinia pestis*.

Conclusions: Clinicians should be aware of the potential risk of this new *Candidatus Rickettsia* sp. and keep in mind other flea-borne infections since these flea species frequently bite humans.

Keywords: Fleas, *Pulex irritans*, *Ctenocephalides felis*, Ecuador, '*Candidatus Rickettsia asemoensis*', *Wolbachia* spp., *Bartonella* spp., *Yersinia pestis*, Plague

Background

Flea-borne diseases are worldwide-distributed emerging and re-emerging infections. Among them, plague, which is caused by *Yersinia pestis*, is the most severe human infection transmitted by fleas [1]. In South America, permanent plague foci exist among native rodent and flea populations in Bolivia, Brazil, Ecuador and Peru [2]. Rats have been the responsible hosts and from them, the disease has spread to other rodents. Ecuador is considered a plague 'hot-spot' since its introduction in 1908, and has experienced important outbreaks. Chimborazo Province has historically been a highly endemic area and the last fatal Ecuadorian cases of plague were reported there in 2004 [3,4].

In addition, fleas are vectors of murine typhus (caused by *Rickettsia typhi*), flea-borne spotted fever (caused by

Rickettsia felis) and harbour *Bartonella* spp. [1,5-7]. Recent evidence of murine typhus in Ecuador is lacking, but the disease may be endemic in localities where commensal rodents (*Rattus* spp.) are abundant. To the best of our knowledge, data about distribution of *R. felis* in this country are unknown and there are no reports describing *Bartonella* spp. in fleas from Ecuador. For these reasons, our interest was focused on the study of flea-borne agents in two Ecuadorian areas (one of them where the last plague outbreak occurred) using molecular biological methods [polymerase chain reaction (PCR) and DNA sequencing].

Methods

In December 2012, a total of 79 fleas were removed from dogs by members of Red Iberoamericana para la Investigación y Control de las Enfermedades Rickettsiales, Programa Iberoamericano de Ciencia y Tecnologías para el Desarrollo (RIICER, CYTED; no. 210RT0403). Fifty-two specimens (44 *Pulex irritans* and 8 *Ctenocephalides felis*)

* Correspondence: jaoteo@riojasalud.es

¹Departamento de Enfermedades Infecciosas, Hospital San Pedro-Centro de Investigación Biomédica de La Rioja (CIBIR), C/ Piqueras 98, 26006 Logroño (La Rioja), Spain

Full list of author information is available at the end of the article

were collected in the Estación Experimental Fátima (ESPOCH), Cantón Puyo, in Pastaza Province (01°24'34.6"S; 77°59'57.5"W), and 27 specimens (all *P. irritans*) were collected in Cantón Guamote, in Chimborazo Province (02°00'25.0"S; 78°47'09.3"W). The study areas had altitudes of 1,034 m and 3,657 m, respectively. After identification at the species level, samples were kept in 70% ethanol at room temperature before being tested. DNA of each arthropod was extracted by lysis with 0.7 M ammonium hydroxide and tested by PCR with the universal primers fD1 and rp2 [8]. This primer pair amplifies the main part of the 16S rRNA gene and has been used for the identification of *Y. pestis* as the causative agent of plague in India [9]. Samples were also screened for the presence of *Rickettsia* spp. with PCR assays targeting rickettsial citrate synthase (*gltA*) and 17 kDa antigen (*htrA*) genes [10,11]. In accordance with the taxonomic scheme [12], additional rickettsial genes (*ompB*, *sca4* and *ompA*) were tested to properly identify *Rickettsia*-positive specimens [13-17]. Moreover, *Bartonella* spp. was tested using RNA polymerase β -subunit-encoding gene (*rpoB*), *gltA* and intergenic spacer region gene (ITS) PCR primers, which amplify fragments of *Bartonella* genes [18-20].

Each PCR included positive controls consisting of *Bartonella henselae* DNA extracted from a cat flea (*C. felis*) from La Rioja - Spain, or *Rickettsia slovaca* strain S14ab DNA (obtained from Vero cells inoculated in our facility with a *Dermacentor marginatus* tick from La Rioja - Spain, and known to be infected with *R. slovaca*). Negative controls (DNA-free water) were included in all assays. Sequences generated by each pair of primers were then compared with those in GenBank using BLAST (www.ncbi.nlm.nih.gov/blast/Blast.cgi).

Results and discussion

PCR assays using universal eubacterial primers for 16S rRNA gene yielded amplicons of different intensity for 69 out of 79 fleas (8/8 *C. felis* and 35/44 *P. irritans* from Pastaza and 26/27 *P. irritans* from Chimborazo). All *C. felis* (n = 8) were also found to be infected with *Rickettsia* species using *gltA* and *htrA* as rickettsial PCR targets, whereas no evidence of *Rickettsia* spp. was found in *P. irritans*. Moreover, no sample was positive for *Bartonella* species as determined either by *rpoB*, *gltA* or ITS PCR assays. Positive and negative controls worked as expected in all cases.

Sequences of rickettsial 16S rRNA gene obtained from 7/8 *C. felis* (1303–1373 bp) showed the closest identity (99.6–99.9%) with '*Candidatus Rickettsia aseboensis*', a potentially new *Rickettsia* species according to the established criteria [12,21] (Table 1). In these samples, the percentage of identity with 16S rRNA gene of a validly published *Rickettsia* species reached 99.4% for *R. felis*

(accession no. NR074483). The 1387 bp-long sequence of 16S rRNA gene obtained from the remaining *C. felis* had the highest identity (97.1%) with a sequence from uncultured flea-associated bacterium [22], and showed 93.6% identity with *Snodgrassella alvi*, betaproteobacteria classified in the family *Neisseriaceae* and previously isolated from the bee gut [23] (Table 1).

Sequencing of *gltA* fragments (350 bp) identified *R. felis*-like rickettsiaceae in *C. felis* (100% identity with *Rickettsia* sp. genotype RF2125) [24] and showed 99.7% identity (349/350 bp) with '*Candidatus R. aseboensis*' (Table 1). The sequences of the 17-kDa amplicons (394 bp) were homologous (100% identical) to each other and to '*Ca. R. aseboensis*' as well as to other molecular isolates included in the *R. felis*-like genotype group (*Rickettsia* sp. SE313 detected in *Echidnophaga gallinacea* from Egypt and *Rickettsia* sp. cf1and5 detected in *C. felis* from USA) (Table 1).

The sequences of *ompB* and *sca4* amplicons (464 and 352 bp, respectively) were also 100% identical to '*Ca. R. aseboensis*' (Table 1). Unfortunately, *ompA* PCR primers did not yield amplicons of the expected size, and inconclusive sequences were obtained for this target gene.

Percentages of identity with validated species *R. felis* (accession no. CP000053) were 98.6, 96.2, 97.9 and 96.6% for *gltA*, *htrA*, *ompB* and *sca4* genes, respectively.

In addition, 59 out of 61 sequences of 16S rRNA gene obtained from 61 *P. irritans* specimens provided evidence of the presence of probable endosymbionts similar to those found within other arthropods and belonging to the genus *Wolbachia* (Table 2). Unfortunately, in two cases (corresponding to two *P. irritans* samples from Chimborazo) it was not possible to get a good-quality sequence to identify the bacteria.

In our study '*Candidatus R. aseboensis*' has been found in fleas that bite humans (*C. felis*) removed from dogs in Ecuador. This potential new species was previously detected in *Ctenocephalides canis* from Kenya and whether it is a human pathogen remains unknown [21]. '*Ca. R. aseboensis*' is highly similar to *Rickettsia* RF2125, a member of the *R. felis*-like genotype group that circulates in fleas from Uruguay [25]. Apart from epidemic typhus (caused by *Rickettsia prowazekii* and transmitted by lice), no data about human diseases associated with *Rickettsia* species have been published from Ecuador [26]. Nevertheless, in the Pastaza province (one of our sampling areas) cases of acute undifferentiated febrile illness compatible with rickettsioses have been reported [27].

Up to the present study, the presence of *R. felis* and/or *Bartonella* spp. has not been demonstrated in fleas from Ecuador. However, *R. felis* has been found in South American fleas from Brazil, Peru, Uruguay, Chile, Argentina and Colombia [11,25,28-31]. In addition, human

Table 1 Results of nucleotide sequence analysis corresponding to PCR products amplified from the 8 *Ctenocephalides felis* specimens of this study

Gene*	Flea gender ¹	Location	Length of sequence (bp)	% identity	Bacteria with closest identity in the BLAST search	GenBank accession no.
16S rRNA	1/2 M; 6/6 F	Pastaza	1303-1373	99.6-99.9	' <i>Candidatus Rickettsia asemoensis</i> ' (<i>Rickettsia</i> F30)	JN315967
	1/2 M; 0/6 F	Pastaza	1387	97.1	Uncultured flea-associated bacterium	EU137419
<i>gltA</i>	2/2 M; 6/6 F	Pastaza	350	93.6	<i>Snodgrassella alvi</i>	JQ746645
				100	<i>Rickettsia felis</i> -like (<i>Rickettsia</i> RF2125)	AF516333
<i>htrA</i>	2/2 M; 6/6 F	Pastaza	394	99.7	' <i>Candidatus Rickettsia asemoensis</i> ' (<i>Rickettsia</i> F30)	JN315968
				100	' <i>Candidatus Rickettsia asemoensis</i> ' (<i>Rickettsia</i> F30)/ <i>Rickettsia felis</i> -like (<i>Rickettsia</i> SE313/cf1and5)	JN315969/DQ166937/ AY953286
<i>ompB</i>	2/2 M; 6/6 F	Pastaza	464	100	' <i>Candidatus Rickettsia asemoensis</i> ' (<i>Rickettsia</i> F30)	JN315972
<i>sca4</i>	2/2 M; 6/6 F	Pastaza	352	100	' <i>Candidatus Rickettsia asemoensis</i> ' (<i>Rickettsia</i> F30)	JN315970

*Inconclusive sequences were obtained for *ompA* gene.

¹Number of positive specimens / Total number of specimens from each flea gender; M: Male; F: Female.

infection with *R. felis* in South America has been confirmed in Brazil by molecular methods [32], and human serological evidence of *R. felis* infection has been recently reported in Colombia [33]. Moreover, there are limited reports describing *Bartonella* spp. in fleas from South America. A molecular study conducted in a *Pulex* specimen found on a Peruvian person evidenced the presence of a potential new *Bartonella* species [5]. Years later, our research group detected *B. rochalimae*, *B. clarridgeiae*, and *B. henselae* in *P. irritans* and *C. felis* collected from cats and dogs in Chile, suggesting the role of fleas as possible vectors of *Bartonella* spp. [7]

Lastly, *Wolbachia* spp. are alphaproteobacteria included in the family *Anaplasmataceae* that were first detected in fleas in 2000 [34]. In this study, the detection rate of *Wolbachia* spp. in *P. irritans* was 83% (59/71). On the contrary, no *C. felis* analysed (0/8) showed evidence of carriage of *Wolbachia* endosymbionts. Previous studies had identified *Wolbachia* in around 20% of cat fleas [35,36]. It has been suggested that *R. felis* infection in fleas might diminish the richness of flea microbiota [37]. According to our data, there is a strong association of

C. felis with '*Ca. R. asemoensis*' whereas *P. irritans* is associated with *Wolbachia* spp. Nevertheless, the interaction of *Wolbachia* with *R. felis* or other related species, such as '*Ca. R. asemoensis*' in fleas needs further investigation.

Based on 16S rDNA analysis, the presence of *Y. pestis* DNA has not been demonstrated in our fleas despite *P. irritans* having been previously described as vectors of *Y. pestis* in Ecuador [2] and plague outbreaks have been repeatedly reported in the Chimborazo region [3,4], where some flea specimens were collected. The rodent flea *Xenopsylla cheopis*, which is the main vector, does not exist in the inter-Andean region of Ecuador (2,500-4,000 m above sea level) possibly, due to very sudden changes in the climatic conditions [38].

Conclusions

In summary, our result confirms the presence of '*Candidatus R. asemoensis*' and *Wolbachia* spp. in fleas removed from dogs in Ecuador. Clinicians should be aware of the potential risk of this new *Candidatus Rickettsia* sp. and keep in mind other flea-borne infections in areas where humans are exposed to fleas.

Table 2 Results of nucleotide sequence analysis corresponding to 59 PCR products amplified¹ from the 71 *Pulex irritans* of this study (44 specimens from Pastaza and 27 from Chimborazo)

Gene	Flea gender ²	Location	Length of sequence (bp)	% identity	Bacteria with closest identity in the BLAST search	Genbank
16S rRNA	9/16 M; 10/28 F	Pastaza	1287-1386	98.3-100	<i>Wolbachia</i> sp. wRi endosymbiont of <i>Drosophila simulans</i>	NR074437
	11/13 M; 13/14 F	Chimborazo				
<i>gltA</i>	4/16 M; 2/28 F	Pastaza	1338-1383	98.4-98.7	<i>Wolbachia</i> sp. endosymbiont of <i>Pseudolynchia canariensis</i>	DQ115538
	0/16 M; 2/28 F	Pastaza	1300-1372	98.3-99.3	<i>Wolbachia</i> sp. endosymbiont of <i>Gryllus crickets</i>	U83094
	0/16 M; 1/28 F	Pastaza	1305	99.3	<i>Wolbachia</i> sp. endosymbiont of <i>Gryllus ovisopis</i>	U83093
	0/16 M; 1/28 F	Pastaza	1374	98.3	<i>Wolbachia</i> sp. endosymbiont of <i>Curculio hachijoensis</i>	AB746399
	0/16 M; 6/28 F	Pastaza	1291-1368	98.3-99	<i>Wolbachia</i> sp. endosymbiont of <i>Kleidocerys resedae</i>	JQ726770

¹In two cases we did not obtain enough good-quality sequences to identify the bacteria, and we did not obtain amplicons for ten specimens.

²Number of positive specimens / Total number of specimens from each flea gender; M: Male; F: Female.

Competing interests

The authors declare they have no competing interests.

Authors' contribution

Designed the study: JMV, MBL, JAO, JZC. Collected and identified fleas: JMV, MBL, JAO, FP, JZC. Processed samples and analyzed sequences: AP. Analyzed the data: AP, JAO. Wrote the paper: AP, JAO. All authors read and approved the final version of the manuscript.

Acknowledgments

We are very grateful to Ana M. Palomar for her tireless help. We appreciate the support of the Red Iberoamericana para la Investigación y Control de las Enfermedades Rickettsiales, Programa Iberoamericano de Ciencia y Tecnologías para el Desarrollo (RIICER, CYTED; no. 210RT0403).

Author details

¹Departamento de Enfermedades Infecciosas, Hospital San Pedro-Centro de Investigación Biomédica de La Rioja (CIBIR), C/ Piqueras 98, 26006 Logroño (La Rioja), Spain. ²Escuela Superior Politécnica de Chimborazo, Riobamba, Ecuador. ³Universidad Autónoma de Yucatán. Centro de investigaciones regionales "Dr. Hideyo Noguchi", Mérida, Yucatán, Mexico. ⁴Laboratorio de Vectores y Enfermedades Transmitidas and Departamento de Parasitología Veterinaria, Facultad de Veterinaria, Universidad de la República, Regional Norte, Salto, Uruguay. ⁵Departamento de Medicina Veterinária Preventiva e Saúde Animal, Faculdade de Medicina Veterinária e Zootecnia, Universidade de São Paulo, São Paulo, Brazil.

Received: 2 May 2014 Accepted: 20 September 2014

Published online: 30 September 2014

References

1. Bitam I, Dittmar K, Parola P, Whiting M, Raoult D: Fleas and flea-borne diseases. *Int J Infect Dis* 2010, **14**:e667–e676.
2. Ruiz A: Plague in the Americas. *Emerg Infect Dis* 2001, **7**(Suppl. 3):539–540.
3. Gabastou JM, Proaño J, Vimos A, Jaramillo G, Hayes E, Gage K, Chu M, Guarnier J, Zaki S, Bowers J, Guillemard C, Tamayo H, Ruiz A: An outbreak of plague including cases with probable pneumonic infection, Ecuador, 1998. *Trans R Soc Trop Med Hyg* 2000, **94**:387–391.
4. Faccini-Martínez AA, Sotomayor HA: Reseña histórica de la peste en Suramérica: una enfermedad poco conocida en Colombia. *Biomédica* 2013, **33**:8–27.
5. Parola P, Shpynov S, Montoya M, López M, Houpiakian P, Zeaiter Z, Guerra H, Raoult D: First molecular evidence of new *Bartonella* spp. in fleas and a tick from Perú. *Am J Trop Med Hyg* 2002, **67**:135–136.
6. Blanco JR, Pérez-Martínez L, Vallejo M, Santibáñez S, Portillo A, Oteo JA: Prevalence of *Rickettsia felis*-like and *Bartonella* spp. in *Ctenocephalides felis* and *Ctenocephalides canis* from La Rioja (Northern Spain). *Ann N Y Acad Sci* 2006, **1078**:270–274.
7. Pérez-Martínez L, Venzal JM, González-Acuña D, Portillo A, Blanco JR, Oteo JA: *Bartonella rochalimae* and other *Bartonella* spp. in fleas, Chile. *Emerg Infect Dis* 2009, **15**:1150–1152.
8. Weisburg WG, Barns SM, Pelletier DA, Lane DJ: 16S ribosomal DNA amplification for phylogenetic study. *J Bacteriol* 1991, **173**:697–703.
9. Shivaji S, Bhanu NV, Aggarwal RK: Identification of *Yersinia pestis* as the causative organism of plague in India as determined by 16S rDNA sequencing and RAPD-based genomic fingerprinting. *FEMS Microbiol Lett* 2000, **189**:247–252.
10. Labruna MB, Whitworth T, Horta MC, Bouyer DH, McBride JW, Pinter A, Popov V, Gennari SM, Walker DH: *Rickettsia* species infecting *Amblyomma cooperi* ticks from an area in the State of Sao Paulo, Brazil, where Brazilian Spotted Fever is endemic. *J Clin Microbiol* 2004, **42**:90–98.
11. Oliveira RP, Galvão MAM, Mafra CL, Chamone CB, Calic SB, Silva SU, Walker DH: *Rickettsia felis* in *Ctenocephalides* spp. fleas, Brazil. *Emerg Infect Dis* 2002, **8**:317–319.
12. Fournier PE, Dumler S, Greub G, Zhang J, Wu Y, Raoult D: Gene sequence-based criteria for identification of new *Rickettsia* isolates and description of *Rickettsia heilongjiangensis* sp. nov. *J Clin Microbiol* 2003, **41**:5456–5465.
13. Choi YJ, Lee SH, Park KH, Koh YS, Lee KH, Baik HS, Choi MS, Kim IS, Jang WJ: Evaluation of PCR-based assay for diagnosis of spotted fever group rickettsiosis in human serum samples. *Clin Vaccine Immunol* 2005, **12**:759–763.
14. Sekeyova Z, Roux V, Raoult D: Phylogeny of *Rickettsia* spp. inferred by comparing sequences of 'gene D', which encodes an intracytoplasmic protein. *Int J Syst Evol Microbiol* 2001, **51**:1353–1360.
15. Jiang J, Blair PJ, Felices V, Moron C, Cespedes M, Anaya E, Schoeler GB, Sumner JW, Olson JG, Richards AL: Phylogenetic analysis of a novel molecular isolate of spotted fever group *Rickettsiae* from northern Peru: *Candidatus Rickettsia andeanae*. *Ann N Y Acad Sci* 2005, **1063**:337–342.
16. Regnery RL, Spruill CL, Plikaytis BD: Genotypic identification of rickettsiae and estimation of intraspecies sequence divergence for portions of two rickettsial genes. *J Bacteriol* 1991, **173**:1576–1589.
17. Roux V, Fournier PE, Raoult D: Differentiation of spotted fever group rickettsiae by sequencing and analysis of restriction fragment length polymorphism of PCR-amplified DNA of the gene encoding the protein rOmpA. *J Clin Microbiol* 1996, **34**:2058–2065.
18. Renesto P, Gouvenet J, Drancourt M, Roux V, Raoult D: Use of *rpoB* gene analysis for detection and identification of *Bartonella* species. *J Clin Microbiol* 2001, **39**:430–437.
19. Norman AF, Regnery R, Jameson P, Greene C, Krause DC: Differentiation of *Bartonella*-like isolates at the species level by PCR-restriction fragment length polymorphism in the citrate synthase gene. *J Clin Microbiol* 1995, **33**:1797–1803.
20. Jensen WA, Fall M, Rooney J, Kordick DL, Breitschwerdt EB: Identification and differentiation of *Bartonella* species using a single step PCR assay. *J Clin Microbiol* 2000, **38**:1717–1722.
21. Jiang J, Maina AN, Knobel DL, Cleaveland S, Laudisoit A, Wamburu K, Ogola E, Parola P, Breiman RF, Njenga MK, Richards AL: Molecular detection of *Rickettsia felis* and *Candidatus Rickettsia asemboensis* in fleas from human habitats, Asembo, Kenya. *Vector Borne Zoonotic Dis* 2013, **13**:550–558.
22. Jones RT, McCormick KF, Martin AP: Bacterial communities of *Bartonella*-positive fleas: diversity and community assembly patterns. *Appl Environ Microbiology* 2008, **74**:1667–1670.
23. Kwong WK, Moran NA: Cultivation and characterization of the gut symbionts of honey bees and bumble bees: description of *Snodgrassella alvi* gen. nov., sp. nov., a member of the family Neisseriaceae of the Betaproteobacteria, and *Gilliamella apicola* gen. nov., sp. nov., a member of Orbaceae fam. nov., Orbales ord. nov., a sister taxon to the order 'Enterobacteriales' of the Gammaproteobacteria. *Int J Syst Evol Microbiol* 2013, **63**:2008–2018.
24. Parola P, Sanogo OY, Lerdtusnee K, Zeaiter Z, Chauvancy G, Gonzalez JP, Miller RS, Telford SR 3rd, Wongsrichanalai C, Raoult D: Identification of *Rickettsia* spp. and *Bartonella* spp. in fleas from the Thai-Myanmar border. *Ann N Y Acad Sci* 2003, **990**:173–181.
25. Venzal JM, Pérez-Martínez L, Félix ML, Portillo A, Blanco JR, Oteo JA: Prevalence of *Rickettsia felis* in *Ctenocephalides felis* and *Ctenocephalides canis* from Uruguay. *Ann N Y Acad Sci* 2006, **1078**:305–308.
26. Labruna MB, Mattar S, Nava S, Bermudez S, Venzal JM, Dolz G, Abarca K, Romero L, de Sousa R, Oteo JA, Zavala-Castro J: Rickettsioses in Latin America, Caribbean, Spain and Portugal. *Rev MVZ Córdoba* 2011, **16**:2435–2457.
27. Manock SR, Jacobsen KH, de Bravo NB, Russell KL, Negrete M, Olson JG, Sanchez JL, Blair PJ, Smalligan RD, Quist BK, Espín JF, Espinoza WR, McCormick F, Fleming LC, Kochel T: Etiology of acute undifferentiated febrile illness in the Amazon basin of Ecuador. *Am J Trop Med Hyg* 2009, **81**:146–151.
28. Blair PJ, Jiang J, Schoeler GB, Moron C, Anaya E, Cespedes M, Cruz C, Felices V, Guevara C, Mendoza L, Villaseca P, Sumner JW, Richards AL, Olson JG: Characterization of spotted fever group rickettsiae in flea and tick specimens from northern Peru. *J Clin Microbiol* 2004, **42**:4961–4967.
29. Labruna MB, Ogrzewalska M, Moraes-Filho J, Lepe P, Gallegos JL, López J: *Rickettsia felis* in Chile. *Emerg Infect Dis* 2007, **13**:1794–1795.
30. Nava S, Pérez-Martínez L, Venzal JM, Portillo A, Santibáñez S, Oteo JA: *Rickettsia felis* in *Ctenocephalides felis* from Argentina. *Vector Borne Zoonotic Dis* 2008, **8**:465–466.
31. Ramírez-Hernández A, Montoya V, Martínez A, Pérez JE, Mercado M, de la Ossa A, Vélez C, Estrada G, Correa MI, Duque L, Ariza JS, Henao C, Valbuena G, Hidalgo M: Molecular detection of *Rickettsia felis* in different flea species from Caldas, Colombia. *Am J Trop Med Hyg* 2013, **89**:453–459.
32. Raoult D, La Scola B, Enea M, Fournier PE, Roux V, Fenollar F, Galvão MA, de Lamballerie X: A flea-associated *Rickettsia* pathogenic for humans. *Emerg Infect Dis* 2001, **7**:73–81.

33. Hidalgo M, Montoya V, Martínez A, Mercado M, De la Ossa A, Vélez C, Estrada G, Pérez JE, Faccini-Martínez AA, Labruna MB, Valbuena G: **Flea-borne rickettsioses in the North of Caldas province, Colombia.** *Vector Borne Zoonotic Dis* 2013, **13**:289–294.
34. Jeyaprakash A, Hoy MA: **Long PCR improves *Wolbachia* DNA amplification: *wsp* sequences found in 76% of sixty-three arthropod species.** *Insect Mol Biol* 2000, **9**:393–405.
35. Gorham CH, Fang QQ, Durden LA: ***Wolbachia* endosymbionts in fleas (Siphonaptera).** *J Parasitol* 2003, **89**:283–289.
36. Rolain JM, Franc M, Davoust B, Raoult D: **Molecular detection of *Bartonella quintana*, *B. koehlerae*, *B. henselae*, *B. clarridgeiae*, *Rickettsia felis*, and *Wolbachia pipientis* in cat fleas, France.** *Emerg Infect Dis* 2003, **9**:338–342.
37. Pornwiroon W, Kearney MT, Husseneder C, Foil LD, Macaluso KR: **Comparative microbiota of *Rickettsia felis*-uninfected and -infected colonized cat fleas, *Ctenocephalides felis*.** *ISME J* 2007, **1**:394–402.
38. Sáenz Vera C: **DDT in the prevention of plague in Ecuador.** *Bull World Health Organ* 1953, **9**:615–618.

doi:10.1186/s13071-014-0455-0

Cite this article as: Oteo *et al.*: '*Candidatus Rickettsia asemboensis*' and *Wolbachia* spp. in *Ctenocephalides felis* and *Pulex irritans* fleas removed from dogs in Ecuador. *Parasites & Vectors* 2014 **7**:455.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

