

Numerical Solution of the Time Dependent 3D Schrödinger Equation Describing Tunneling of Atoms from Anharmonic Traps

Ilyas Ishmukhamedov^{1,2,3,*}, Altay Ishmukhamedov^{2,4,**}, and Vladimir Melezhhik^{1,5,***}

¹Bogoliubov Laboratory of Theoretical Physics, Joint Institute for Nuclear Research, 141980 Dubna, Moscow Region, Russian Federation

²Al-Farabi Kazakh National University, Almaty 050040, Republic of Kazakhstan

³Institute of Nuclear Physics, the Ministry of Energy of the Republic of Kazakhstan, Almaty 050032, Republic of Kazakhstan

⁴K.I. Satpaev Institute of Geological Sciences, Almaty 050010, Republic of Kazakhstan

⁵Peoples' Friendship University of Russia (RUDN University), 6 Miklukho-Maklaya St., 117198 Moscow, Russian Federation

Abstract. We present an efficient numerical method for the integration of the 3D Schrödinger equation. A tunneling problem of two interacting bosonic atoms confined in a 1D anharmonic trap has been successfully solved by means of this method. We demonstrate fast convergence of the final results with respect to spatial and temporal grid steps. The computational scheme is based on the operator-splitting technique with the implicit Crank-Nicolson algorithm on spatial sixth-order finite-differences. The computational time is proportional to the number of spatial grid points.

1 Introduction

Pair correlation plays a major role in quantum tunneling and there are a only few works in the context of ultracold physics which accurately take into account the interatomic interaction [1–7]. In the present paper, we consider a tunneling of two interacting atoms from an anharmonic optical trap. We solve this problem by means of a numerical method which is based on the split-operator approximation [1–3, 8, 9], the Crank-Nicolson scheme and the sixth-order spatial finite-differences.

2 Method

We consider the time-dependent Schrödinger equation (throughout $\hbar = \omega = m = 1$ units are used)

$$i \frac{\partial \psi(x_1, x_2, t)}{\partial t} = H(x_1, x_2) \psi(x_1, x_2, t), \quad (1)$$

with the Hamiltonian

$$H(x_1, x_2) = H_1(x_1) + H_2(x_2) + V^{(aa)}(x_1 - x_2), \quad (2)$$

* e-mail: i.ishmukhamedov@mail.ru

** e-mail: ishmukhamedov.altay@gmail.com

*** e-mail: melezhhik@theor.jinr.ru

where $V^{(aa)}$ describes the atom-atom interaction and $H_j(x_j)$ denote the single-atom Hamiltonians

$$H_j(x_j) = -\frac{1}{2} \frac{\partial^2}{\partial x_j^2} + V^{(at)}(x_j), \quad j = 1, 2, \quad (3)$$

in which $V^{(at)}$ describes the atom-trap interaction. The partial derivatives in (3) are approximated by sixth-order finite-differences.

To integrate (1) we use the split-operator method the basic ideas of which have been proposed in [8] and which has been developed in [2, 3, 9] in connection with the solution of confined ultracold atom-atom collisions in waveguide-like traps:

$$\begin{aligned} \psi(x_1, x_2, t + \Delta t) = & \exp \left\{ -i \frac{\Delta t}{2} V^{(aa)}(x_1 - x_2) \right\} \exp \{-i\Delta t H_1(x_1)\} \exp \{-i\Delta t H_2(x_2)\} \\ & \times \exp \left\{ -i \frac{\Delta t}{2} V^{(aa)}(x_1 - x_2) \right\} \psi(x_1, x_2, t). \end{aligned} \quad (4)$$

The actions of the operators $\exp \{-i\Delta t H_j(x_j)\}$ are approximated by the implicit Crank-Nicolson scheme, which maintains the accuracy order of the split-operator scheme (4) to $\mathcal{O}(\Delta t^3)$.

When considering a problem of particle tunneling it is necessary to correctly model the outgoing wave function at $x_1, x_2 \rightarrow \pm\infty$. This can be done by introducing into the original Hamiltonian (2) a complex absorbing potential (CAP) in the form suggested in [10]:

$$W(x_j) = w_c(|x_j| - x_c)^2 \theta(|x_j| - x_c), \quad j = 1, 2, \quad (5)$$

where $\theta(x)$ is the Heaviside step function. The values of interest of the parameters w_c and x_c are fixed after Eq. (8).

3 Results

The efficiency of our method is demonstrated in the solution of the topical problem of tunneling of cold atoms [1]. The atom-trap interaction is chosen of the standing-wave form:

$$V^{(at)}(x) = \begin{cases} 1/(2\beta^2) \cdot \sin^2(\beta x), & |x| \leq \pi/\beta, \\ 0, & |x| > \pi/\beta, \end{cases} \quad (6)$$

where $\beta = 2\pi/\lambda \cdot \sqrt{\hbar/(m\omega)}$ (in standard units), λ and ω represent the trap wavelength and frequency correspondingly. For $\beta = 0.427471$, this models an actual atom-trap interaction [1, 11, 12] in modern experiments [13] on ultracold atoms (figure 1).




Figure 1. Trap potential $V^{(at)}(x)$.

The atom-atom interaction is taken of Gaussian form:

$$V^{(aa)}(x_1 - x_2) = -V_0 \exp \left\{ -\frac{(x_1 - x_2)^2}{2r_0^2} \right\}, \quad (7)$$

where the range of the interaction is fixed to $r_0 = 0.1$ [4, 14] and the interaction strength V_0 is varied.

We calculate the total probability $P(t)$ in the box $|x_1, x_2| \leq 20$. For a wide range of values V_0 , $P(t)$ decays exponentially. This allows us to extract the tunneling rate γ as

$$\gamma = -\frac{1}{P(t)} \frac{dP(t)}{dt}. \quad (8)$$

We find that the parameters of the CAP (5), $w_c = -0.1$ and $x_c = 10$, lie in the domain where the tunneling rate γ does not change significantly under the variation of these parameters.

To solve (1), the value of the initial wave function is to be known. We fix it by replacing the trap (6) with the “closed” sextic (i.e., the power series expansion of (6) up to x_j^6) trapping potential [1, 15] and calculating the eigenstates of the doubly excited relative and center-of-mass motion states of the stationary Schrödinger equation by using the numerical scheme from [12].

Figure 2 shows the fast convergence of the errors

$$\text{error}(\delta) = \frac{|\gamma(\delta \rightarrow 0) - \gamma(\delta)|}{\gamma(\delta \rightarrow 0)}, \quad \delta = \Delta x, \Delta t. \quad (9)$$

and tunneling rates γ with the spatial Δx and temporal Δt steps.




Figure 2. (a) Dependence of the relative errors of the computed γ values (closed circles, left side scale) and dependence of the tunneling rate γ (open circles, right side scale) on the spatial grid step Δx . (b) Dependence of the relative errors of the computed γ values (closed circles, left side scale) and dependence of the tunneling rate γ (open circles, right side scale) on the temporal grid step Δt . Outputs are reported for three values of the $V^{(aa)}$ potential: $V_0 = 9$ and -42.6 (doubly excited relative motion state at $t \leq 0$) and $V_0 = 4.2$ (doubly excited center-of-mass motion state at $t \leq 0$).

In figure 3 one can see that the computational time grows linearly with the number of spatial grid points N , according to the sweeping method theory [16].

4 Conclusion

We have solved a tunneling problem of two cold interacting atoms by using a numerical method. The interatomic coupling strength is varied in a wide range – from strong attraction to strong repulsion.




Figure 3. Dependence of the CPU time (in minutes) for the derivation of the tunneling rate γ (open circles) at $V_0 = 9$ (doubly excited relative motion state at $t \leq 0$) on the number of spatial grid points $N = N_1 \times N_2$ (N_j defines the grid along x_j). The solid line is a fit function. The computation has been performed until the temporal variable reaches the value $t = 10$ using the server at the Bogoliubov Laboratory of Theoretical Physics, JINR (i7a: 8-core Intel E5-2690 @2.9-3.8GHz, 128 GB ECC RAM, GPU GTX680).

The method is based on the split-operator approximation of the time-evolution operator and Crank-Nicolson approximation of the single-particle Hamiltonians. We have found that the error of the method reduces quite fast with reducing the step sizes of the spatial and temporal grids. The method can be extended to more complicated problems with more degrees of freedom.

Acknowledgement

The work was financially supported by the Ministry of Education and Science of the Russian Federation (Agreement No. 02.a003.21.0008) and by the Grant of the Plenipotentiary Representative of the Republic of Kazakhstan to JINR.

References

- [1] I.S. Ishmukhamedov and V.S. Melezhik, *Phys. Rev. A* **95**, 062701 (2017)
- [2] V.S. Melezhik, J.I. Kim, and P. Schmelcher, *Phys. Rev. A* **76**, 053611 (2007)
- [3] V.S. Melezhik, *EPJ Web Conf.* **108**, 01008 (2016)
- [4] S.E. Gharashi and D. Blume, *Phys. Rev. A* **92**, 033629 (2015).
- [5] P. M. Krassovitskiy and F. M. Pen'kov, *J. Phys. B: At. Mol. Opt. Phys.* **47**, 225210 (2014)
- [6] V. O. Nesterenko, A. N. Novikov, and E. Suraud, *Laser Phys.* **24**, 125501 (2014)
- [7] A. U. J. Lode, A. I. Streltsov, O. E. Alon, H.-D. Meyer, and L. S. Cederbaum, *J. Phys. B* **42**, 044018 (2009)
- [8] G.I. Marchuk, *Methods of Numerical Mathematics* (Springer-Verlag, New York, 1975) Sec. 4.3.3
- [9] V.S. Melezhik, *Atoms and Molecules in Strong External Fields* (Plenum, New-York and London, 1998) pp. 89–94
- [10] U. V. Riss and H.-D. Meyer, *J. Phys. B* **26**, 4503 (1993)
- [11] S.-G. Peng, H. Hu, X.-J. Liu, and P.D. Drummond, *Phys. Rev. A* **84**, 043619 (2011)
- [12] I.S. Ishmukhamedov, D.T. Aznabayev, and S.A. Zhaugasheva, *Phys. Part. Nuclei Lett.* **12**: 680 (2015)
- [13] E. Haller, M. J. Mark, R. Hart, J. G. Danzl, L. Reichsollner, V. Melezhik, P. Schmelcher, and H.-C. Nägerl, *Physical Review Letters*, **104**, 153203, (2010)
- [14] I.S. Ishmukhamedov, D.S. Valiolda, and S.A. Zhaugasheva, *Phys. Part. Nuclei Lett.* **11**: 238 (2014)
- [15] S. Grishkevich and A. Saenz, *Phys. Rev A* **80**, 013403 (2009)
- [16] I.M. Gelfand and S.V. Fomin, *Calculus of Variations*, (Dover Publications, New York, 2000)