

Research Article

Some Embeddings into the Morrey and Modified Morrey Spaces Associated with the Dunkl Operator

Emin V. Guliyev¹ and Yagub Y. Mammadov^{1,2}

¹ Institute of Mathematics and Mechanics, Azerbaijan National Academy of Sciences, Baku, Azerbaijan

² Nakhchivan Teacher-Training Institute, Azerbaijan

Correspondence should be addressed to Yagub Y. Mammadov, yagubmammadov@yahoo.com

Received 29 October 2009; Revised 5 February 2010; Accepted 2 March 2010

Academic Editor: Ağacik Zafer

Copyright © 2010 E. V. Guliyev and Y. Y. Mammadov. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

We consider the generalized shift operator, associated with the Dunkl operator $\Lambda_\alpha(f)(x) = (d/dx)f(x) + ((2\alpha+1)/x)((f(x)-f(-x))/2)$, $\alpha > -1/2$. We study some embeddings into the Morrey space (D -Morrey space) $L_{p,\lambda,\alpha}$, $0 \leq \lambda < 2\alpha + 2$ and modified Morrey space (modified D -Morrey space) $\tilde{L}_{p,\lambda,\alpha}$ associated with the Dunkl operator on \mathbb{R} . As applications we get boundedness of the fractional maximal operator M_β , $0 \leq \beta < 2\alpha + 2$, associated with the Dunkl operator (fractional D -maximal operator) from the spaces $L_{p,\lambda,\alpha}$ to $L_\infty(\mathbb{R})$ for $p = (2\alpha + 2 - \lambda)/\beta$ and from the spaces $\tilde{L}_{p,\lambda,\alpha}(\mathbb{R})$ to $L_\infty(\mathbb{R})$ for $(2\alpha + 2 - \lambda)/\beta \leq p \leq (2\alpha + 2)/\beta$.

1. Introduction

On the real line, the Dunkl operators are differential-difference operators introduced in 1989 by Dunkl [1] and are denoted by Λ_α , where α is a real parameter $> -1/2$. These operators are associated with the reflection group \mathbb{Z}_2 on \mathbb{R} . Rösler in [2] shows that the Dunkl kernel verifies a product formula. This allows us to define the Dunkl translations τ_x , $x \in \mathbb{R}$.

In the theory of partial differential equations, together with weighted $L_{p,w}(\mathbb{R}^n)$ spaces, Morrey spaces $L_{p,\lambda}(\mathbb{R}^n)$ play an important role. Morrey spaces were introduced by Morrey in 1938 in connection with certain problems in elliptic partial differential equations and calculus of variations (see [3]). Later, Morrey spaces found important applications to Navier-Stokes [4, 5] and Schrödinger [6–8] equations, elliptic problems with discontinuous coefficients [9, 10], and potential theory [11–13]. An exposition of the Morrey spaces can be found in the book [14].

In the present work, we study some embeddings into the D -Morrey and modified D -Morrey spaces. As applications we give boundedness of the fractional D -maximal operator in the D -Morrey and modified D -Morrey spaces.

The paper is organized as follows. In Section 2, we present some definitions and auxiliary results. In Section 3, we give some embeddings into the D -Morrey and modified D -Morrey spaces. In Section 4, we prove the boundedness of the fractional D -maximal operator M_β from the spaces $L_{p,\lambda,\alpha}$ to $L_\infty(\mathbb{R})$ for $p = (2\alpha + 2 - \lambda)/\beta$ and from the spaces $\tilde{L}_{p,\lambda,\alpha}(\mathbb{R})$ to $L_\infty(\mathbb{R})$ for $(2\alpha + 2 - \lambda)/\beta \leq p \leq (2\alpha + 2)/\beta$.

2. Preliminaries

On the real line, we consider the first-order differential-difference operator defined by

$$\Lambda_\alpha(f)(x) := \frac{d}{dx}f(x) + \frac{2\alpha + 1}{x} \left(\frac{f(x) - f(-x)}{2} \right), \quad \alpha > -\frac{1}{2}, \quad (2.1)$$

which is called the Dunkl operator. For $\lambda \in \mathbb{C}$, the Dunkl kernel $E_\alpha(\lambda \cdot)$ on \mathbb{R} was introduced by Dunkl in [1] (see also [15–17]) and is given by

$$E_\alpha(\lambda x) = j_\alpha(i\lambda x) + \frac{\lambda x}{2(\alpha + 1)} j_{\alpha+1}(i\lambda x), \quad x \in \mathbb{R}, \quad (2.2)$$

where j_α is the normalized Bessel function of the first kind and order α [18], defined by

$$j_\alpha(z) := 2^\alpha \Gamma(\alpha + 1) \frac{J_\alpha(z)}{z^\alpha} = \Gamma(\alpha + 1) \sum_{n=0}^{\infty} \frac{(-1)^n (z/2)^{2n}}{n! \Gamma(n + \alpha + 1)}, \quad z \in \mathbb{C}. \quad (2.3)$$

The Dunkl kernel $E_\alpha(\lambda \cdot)$ is the unique analytic solution on \mathbb{R} of the initial problem for the Dunkl operator (see [1]).

Let μ_α be the weighted Lebesgue measure on \mathbb{R} given by

$$d\mu_\alpha(x) := \frac{|x|^{2\alpha+1}}{2^{\alpha+1} \Gamma(\alpha + 1)} dx. \quad (2.4)$$

For every $1 \leq p \leq \infty$, we denote by $L_{p,\alpha}(\mathbb{R}) = L_p(d\mu_\alpha)$ the spaces of complex-valued functions f , measurable on \mathbb{R} such that

$$\begin{aligned} \|f\|_{L_{p,\alpha}} &:= \left(\int_{\mathbb{R}} |f(x)|^p d\mu_\alpha(x) \right)^{1/p} < \infty \quad \text{if } p \in [1, \infty), \\ \|f\|_{L_{\infty,\alpha}} &:= \operatorname{ess\,sup}_{x \in \mathbb{R}} |f(x)| \quad \text{if } p = \infty. \end{aligned} \quad (2.5)$$

For $1 \leq p < \infty$, we denote by $WL_{p,\alpha}(\mathbb{R})$ the weak $L_{p,\alpha}$ spaces defined as the set of locally integrable functions $f(x)$, $x \in \mathbb{R}$ with the finite norm

$$\|f\|_{WL_{p,\alpha}} := \sup_{r>0} r(\mu_\alpha\{x \in \mathbb{R} : |f(x)| > r\})^{1/p}. \quad (2.6)$$

Note that

$$L_{p,\alpha}(\mathbb{R}) \subset WL_{p,\alpha}(\mathbb{R}), \quad \|f\|_{WL_{p,\alpha}} \leq \|f\|_{L_{p,\alpha}} \quad \forall f \in L_{p,\alpha}. \quad (2.7)$$

For all $x, y, z \in \mathbb{R}$, we put

$$W_\alpha(x, y, z) := (1 - \sigma_{x,y,z} + \sigma_{z,x,y} + \sigma_{z,y,x}) \Delta_\alpha(x, y, z), \quad (2.8)$$

where

$$\sigma_{x,y,z} := \begin{cases} \frac{x^2 + y^2 - z^2}{2xy} & \text{if } x, y \in \mathbb{R} \setminus \{0\}, \\ 0 & \text{otherwise,} \end{cases} \quad (2.9)$$

and Δ_α is the Bessel kernel given by

$$\Delta_\alpha(x, y, z) := \begin{cases} d_\alpha \frac{\left(\left[(|x| + |y|)^2 - z^2 \right] \left[z^2 - (|x| - |y|)^2 \right] \right)^{\alpha-1/2}}{|xyz|^{2\alpha}} & \text{if } |z| \in A_{x,y}, \\ 0 & \text{otherwise,} \end{cases} \quad (2.10)$$

where $d_\alpha = (\Gamma(\alpha + 1))^2 / (2^{\alpha-1} \sqrt{\pi} \Gamma(\alpha + 1/2))$ and $A_{x,y} = [||x| - |y||, |x| + |y|]$.

In the sequel we consider the signed measure $\nu_{x,y}$, on \mathbb{R} , given by

$$\nu_{x,y} := \begin{cases} W_\alpha(x, y, z) d\mu_\alpha(z) & \text{if } x, y \in \mathbb{R} \setminus \{0\}, \\ d\delta_x(z) & \text{if } y = 0, \\ d\delta_y(z) & \text{if } x = 0. \end{cases} \quad (2.11)$$

For $x, y \in \mathbb{R}$ and f being a continuous function on \mathbb{R} , the Dunkl translation operator τ_x is given by

$$\tau_x f(y) := \int_{\mathbb{R}} f(z) d\nu_{x,y}(z). \quad (2.12)$$

Using the change of variable $z = \Psi(x, y, \theta) = \sqrt{x^2 + y^2 - 2xy \cos \theta}$, we have also (see [2])

$$\tau_x f(y) = C_\alpha \int_0^\pi \left[f(\Psi) + f(-\Psi) + \frac{x+y}{\Psi} (f(\Psi) - f(-\Psi)) \right] d\nu_\alpha(\theta), \quad (2.13)$$

where $d\nu_\alpha(\theta) = (1 - \cos \theta) \sin^{2\alpha} \theta d\theta$ and $C_\alpha = \Gamma(\alpha + 1) / 2\sqrt{\pi} \Gamma(\alpha + 1/2)$.

Proposition 2.1 (see Soltani [16]). For all $x \in \mathbb{R}$ the operator τ_x extends to $L_{p,\alpha}(\mathbb{R})$, $p \geq 1$ and we have for $f \in L_{p,\alpha}(\mathbb{R})$,

$$\|\tau_x f\|_{L_{p,\alpha}} \leq 4\|f\|_{L_{p,\alpha}}. \quad (2.14)$$

Let $B(0, t) =] -t, t[$, $t > 0$ and $\mu_\alpha(] -t, t[) = b_\alpha t^{2\alpha+2}$, where $b_\alpha = 2^{-\alpha-1}((\alpha+1)\Gamma(\alpha+1))^{-1}$. For $L_{1,\alpha}^{\text{loc}}(\mathbb{R})$ (the space of locally integrable functions on \mathbb{R}), we consider

$$Mf(x) := \sup_{r>0} (\mu_\alpha B(0, r))^{-1} \int_{B(0, r)} \tau_x |f|(y) d\mu_\alpha(y). \quad (2.15)$$

Theorem 2.2 (see [19]). (1) If $f \in L_{1,\alpha}(\mathbb{R})$, then for every $\beta > 0$,

$$\mu_\alpha \{x \in \mathbb{R} : Mf(x) > \beta\} \leq \frac{C}{\beta} \|f\|_{L_{1,\alpha}}, \quad (2.16)$$

where $C > 0$ is independent of f .

(2) If $f \in L_{p,\alpha}(\mathbb{R})$, $1 < p \leq \infty$, then $Mf \in L_{p,\alpha}(\mathbb{R})$ and

$$\|Mf\|_{L_{p,\alpha}} \leq C_p \|f\|_{L_{p,\alpha}}, \quad (2.17)$$

where $C_p > 0$ is independent of f .

Corollary 2.3. If $f \in L_{1,\alpha}^{\text{loc}}(\mathbb{R})$, then

$$\lim_{r \rightarrow 0} (\mu_\alpha B(0, r))^{-1} \int_{B(0, r)} \tau_x f(y) d\mu_\alpha(y) = f(x) \quad (2.18)$$

for a.e. $x \in \mathbb{R}$.

3. Some Embeddings into the D -Morrey and Modified D -Morrey Spaces

Definition 3.1 (see [20]). Let $1 \leq p < \infty$, $0 \leq \lambda \leq 2\alpha + 2$, and $[t]_1 = \min\{1, t\}$, $t > 0$. We denote by $L_{p,\lambda,\alpha}(\mathbb{R})$ Morrey space ($\equiv D$ -Morrey space) and by $\tilde{L}_{p,\lambda,\alpha}(\mathbb{R})$ the modified Morrey space (\equiv modified D -Morrey space), associated with the Dunkl operator as the set of locally integrable functions $f(x)$, $x \in \mathbb{R}$, with the finite norms

$$\begin{aligned} \|f\|_{L_{p,\lambda,\alpha}} &:= \sup_{x \in \mathbb{R}, t > 0} \left(t^{-\lambda} \int_{B(0, t)} \tau_x |f|^p(y) d\mu_\alpha(y) \right)^{1/p}, \\ \|f\|_{\tilde{L}_{p,\lambda,\alpha}} &:= \sup_{x \in \mathbb{R}, t > 0} \left([t]_1^{-\lambda} \int_{B(0, t)} \tau_x |f|^p(y) d\mu_\alpha(y) \right)^{1/p}, \end{aligned} \quad (3.1)$$

respectively.

If $\lambda < 0$ or $\lambda > 2\alpha + 2$, then $\tilde{L}_{p,\lambda,\alpha}(\mathbb{R}) = \Theta$, where Θ is the set of all functions equivalent to 0 on \mathbb{R} .

Note that

$$\begin{aligned} L_{p,\alpha}(\mathbb{R}) \subset \tilde{L}_{p,0,\alpha}(\mathbb{R}) &= L_{p,0,\alpha}(\mathbb{R}), \\ \|f\|_{\tilde{L}_{p,0,\alpha}} &= \|f\|_{L_{p,0,\alpha}} \leq 4\|f\|_{L_{p,\alpha}}, \end{aligned} \tag{3.2}$$

$$\begin{aligned} \tilde{L}_{p,\lambda,\alpha}(\mathbb{R}) \subset L_{p,\alpha}(\mathbb{R}), \quad \|f\|_{L_{p,\alpha}} &\leq \|f\|_{\tilde{L}_{p,\lambda,\alpha}}, \\ \tilde{L}_{p,\lambda,\alpha}(\mathbb{R}) \subset L_{p,\lambda,\alpha}(\mathbb{R}), \quad \|f\|_{L_{p,\lambda,\alpha}} &\leq \|f\|_{\tilde{L}_{p,\lambda,\alpha}}. \end{aligned} \tag{3.3}$$

Definition 3.2 (see [19]). Let $1 \leq p < \infty$, $0 \leq \lambda \leq 2\alpha + 2$. We denote by $WL_{p,\lambda,\alpha}(\mathbb{R})$ the weak D -Morrey space and by $W\tilde{L}_{p,\lambda,\alpha}(\mathbb{R})$ the modified weak D -Morrey space as the set of locally integrable functions $f(x)$, $x \in \mathbb{R}$ with finite norms

$$\begin{aligned} \|f\|_{WL_{p,\lambda,\alpha}} &:= \sup_{r>0} r \sup_{x \in \mathbb{R}, t>0} \left(t^{-\lambda} \mu_\alpha \{y \in B(0, t) : \tau_x |f|(y) > r\} \right)^{1/p}, \\ \|f\|_{W\tilde{L}_{p,\lambda,\alpha}} &:= \sup_{r>0} r \sup_{x \in \mathbb{R}, t>0} \left([t]_1^{-\lambda} \mu_\alpha \{y \in B(0, t) : \tau_x |f|(y) > r\} \right)^{1/p}, \end{aligned} \tag{3.4}$$

respectively.

We note that

$$\begin{aligned} L_{p,\lambda,\alpha}(\mathbb{R}) \subset WL_{p,\lambda,\alpha}(\mathbb{R}), \quad \|f\|_{WL_{p,\lambda,\alpha}} &\leq \|f\|_{L_{p,\lambda,\alpha}}, \\ \tilde{L}_{p,\lambda,\alpha}(\mathbb{R}) \subset W\tilde{L}_{p,\lambda,\alpha}(\mathbb{R}), \quad \|f\|_{W\tilde{L}_{p,\lambda,\alpha}} &\leq \|f\|_{\tilde{L}_{p,\lambda,\alpha}}. \end{aligned} \tag{3.5}$$

Lemma 3.3 (see [20]). *Let $1 \leq p < \infty$. Then*

$$\begin{aligned} L_{p,2\alpha+2,\alpha}(\mathbb{R}) &= L_\infty(\mathbb{R}), \\ \|f\|_{L_{p,2\alpha+2,\alpha}} &= 4b_\alpha^{1/p} \|f\|_{L_\infty}. \end{aligned} \tag{3.6}$$

Lemma 3.4. *Let $1 \leq p < \infty$, $0 \leq \lambda \leq 2\alpha + 2$. Then*

$$\begin{aligned} \tilde{L}_{p,\lambda,\alpha}(\mathbb{R}) &= L_{p,\lambda,\alpha}(\mathbb{R}) \cap L_{p,\alpha}(\mathbb{R}), \\ \max \{ \|f\|_{L_{p,\lambda,\alpha}}, \|f\|_{L_{p,\alpha}} \} &\leq \|f\|_{\tilde{L}_{p,\lambda,\alpha}} \leq \max \{ \|f\|_{L_{p,\lambda,\alpha}}, 4\|f\|_{L_{p,\alpha}} \}. \end{aligned} \tag{3.7}$$

Proof. Let $f \in \tilde{L}_{p,\lambda,\alpha}(\mathbb{R})$. Then by (3.3) we have

$$\begin{aligned} \tilde{L}_{p,\lambda,\alpha}(\mathbb{R}) &\subset_{>} L_{p,\lambda,\alpha}(\mathbb{R}) \cap L_{p,\alpha}(\mathbb{R}), \\ \max\{\|f\|_{L_{p,\lambda,\alpha}}, \|f\|_{L_{p,\alpha}}\} &\leq \|f\|_{\tilde{L}_{p,\lambda,\alpha}}. \end{aligned} \quad (3.8)$$

Let $f \in L_{p,\lambda,\alpha}(\mathbb{R}) \cap L_{p,\alpha}(\mathbb{R})$. Then

$$\begin{aligned} \|f\|_{\tilde{L}_{p,\lambda,\alpha}} &= \sup_{x \in \mathbb{R}, t > 0} \left([t]_1^{-\lambda} \int_{B(0,t)} \tau_x |f|^p(y) d\mu_\alpha(y) \right)^{1/p} \\ &= \max \left\{ \sup_{x \in \mathbb{R}, 0 < t \leq 1} \left(t^{-\lambda} \int_{B(0,t)} \tau_x |f|^p(y) d\mu_\alpha(y) \right)^{1/p}, \right. \\ &\quad \left. \sup_{x \in \mathbb{R}, t > 1} \left(\int_{B(0,t)} \tau_x |f|^p(y) d\mu_\alpha(y) \right)^{1/p} \right\} \leq \max\{\|f\|_{L_{p,\lambda,\alpha}}, 4\|f\|_{L_{p,\alpha}}\}. \end{aligned} \quad (3.9)$$

Therefore, $f \in \tilde{L}_{p,\lambda,\alpha}(\mathbb{R})$ and the embedding $L_{p,\lambda,\alpha}(\mathbb{R}) \cap L_{p,\alpha}(\mathbb{R}) \subset_{>} \tilde{L}_{p,\lambda,\alpha}(\mathbb{R})$ is valid.

Thus $\tilde{L}_{p,\lambda,\alpha}(\mathbb{R}) = L_{p,\lambda,\alpha}(\mathbb{R}) \cap L_{p,\alpha}(\mathbb{R})$. \square

From Lemmas 3.3 and 3.4 for $1 \leq p < \infty$, we have

$$\tilde{L}_{p,2\alpha+2,\alpha}(\mathbb{R}) = L_\infty(\mathbb{R}) \cap L_{p,\alpha}(\mathbb{R}). \quad (3.10)$$

Lemma 3.5. *Let $0 \leq \lambda \leq 2\alpha + 2$. Then*

$$L_{(2\alpha+2)/(2\alpha+2-\lambda),\alpha}(\mathbb{R}) \subset_{>} L_{1,\lambda,\alpha}(\mathbb{R}), \quad \|f\|_{L_{1,\lambda,\alpha}} \leq 4b_\alpha^{\lambda/(2\alpha+2)} \|f\|_{L_{(2\alpha+2)/(2\alpha+2-\lambda),\alpha}}. \quad (3.11)$$

Proof. The embedding is a consequence of Hölder's inequality and Proposition 2.1. Indeed,

$$\begin{aligned} \|f\|_{L_{1,\lambda,\alpha}} &= \sup_{x \in \mathbb{R}, t > 0} t^{-\lambda} \int_{B(0,t)} \tau_x |f|(y) d\mu_\alpha(y) \\ &\leq \sup_{x \in \mathbb{R}, t > 0} t^{-\lambda} (\mu_\alpha B(0,t))^{\lambda/(2\alpha+2)} \left(\int_{B(0,t)} \tau_x |f|^{(2\alpha+2)/(2\alpha+2-\lambda)} d\mu_\alpha(y) \right)^{(2\alpha+2-\lambda)/(2\alpha+2)} \\ &\leq 4b_\alpha^{\lambda/(2\alpha+2)} \|f\|_{L_{(2\alpha+2)/(2\alpha+2-\lambda),\alpha}}. \end{aligned} \quad (3.12)$$

\square

On the D -Morrey spaces, the following embedding is valid.

Lemma 3.6 (see [20]). *Let $0 \leq \lambda < 2\alpha + 2$ and $0 \leq \beta < 2\alpha + 2 - \lambda$. Then for $p = (2\alpha + 2 - \lambda)/\beta$,*

$$L_{p,\lambda,\alpha}(\mathbb{R}) \subset L_{1,2\alpha+2-\beta,\alpha}(\mathbb{R}), \quad \|f\|_{L_{1,2\alpha+2-\beta,\alpha}} \leq b_\alpha^{1/p'} \|f\|_{L_{p,\lambda,\alpha}}, \quad (3.13)$$

where $1/p + 1/p' = 1$.

On the modified D -Morrey spaces, the following embedding is valid.

Lemma 3.7. *Let $0 \leq \lambda < 2\alpha + 2$ and $0 \leq \beta < 2\alpha + 2 - \lambda$. Then for $(2\alpha + 2 - \lambda)/\beta \leq p \leq (2\alpha + 2)/\beta$,*

$$\tilde{L}_{p,\lambda,\alpha}(\mathbb{R}) \subset L_{1,2\alpha+2-\beta,\alpha}(\mathbb{R}), \quad \|f\|_{L_{1,2\alpha+2-\beta,\alpha}} \leq b_\alpha^{1/p'} \|f\|_{\tilde{L}_{p,\lambda,\alpha}}. \quad (3.14)$$

Proof. Let $0 < \lambda < 2\alpha + 2$, $0 < \beta < 2\alpha + 2 - \lambda$, $f \in \tilde{L}_{p,\lambda,\alpha}(\mathbb{R})$, and $(2\alpha + 2 - \lambda)/\beta \leq p \leq (2\alpha + 2)/\beta$. By the Hölder's inequality, we have

$$\begin{aligned} \|f\|_{L_{1,2\alpha+2-\beta,\alpha}} &= \sup_{x \in \mathbb{R}, t > 0} [t]_1^{\beta-2\alpha-2} \int_{B(0,t)} \tau_x |f|(y) d\mu_\alpha(y) \\ &\leq b_\alpha^{1/p'} \sup_{x \in \mathbb{R}, t > 0} \left([t]_1 t^{-1} \right)^{-(2\alpha+2)/p'} [t]_1^{\beta-(2\alpha+2-\lambda)/p} \\ &\quad \times \left([t]_1^{-\lambda} \int_{B(0,t)} \tau_x |f|^p(y) d\mu_\alpha(y) \right)^{1/p} \\ &= b_\alpha^{1/p'} \sup_{x \in \mathbb{R}, t > 0} \left([t]_1 t^{-1} \right)^{2\alpha+2-\beta} \left([t]_1 t^{-1} \right)^{-(2\alpha+2)/p'} [t]_1^{\beta-(2\alpha+2-\lambda)/p} \\ &\quad \times \left([t]_1^{-\lambda} \int_{B(0,t)} \tau_x |f|^p(y) d\mu_\alpha(y) \right)^{1/p} \\ &\leq b_\alpha^{1/p'} \|f\|_{\tilde{L}_{p,\lambda,\alpha}} \sup_{t > 0} \left([t]_1 t^{-1} \right)^{(2\alpha+2)/p-\beta} [t]_1^{\beta-(2\alpha+2-\lambda)/p}. \end{aligned} \quad (3.15)$$

Note that

$$\begin{aligned} \sup_{t > 0} \left([t]_1 t^{-1} \right)^{(2\alpha+2)/p-\beta} [t]_1^{\beta-(2\alpha+2-\lambda)/p} &= \max \left\{ \sup_{0 < t \leq 1} t^{\beta-(2\alpha+2-\lambda)/p}, \sup_{t > 1} t^{\beta-(2\alpha+2)/p} \right\} < \infty \\ &\text{iff } \frac{2\alpha + 2 - \lambda}{\beta} \leq p \leq \frac{2\alpha + 2}{\beta}. \end{aligned} \quad (3.16)$$

Therefore, $f \in L_{1,2\alpha+2-\beta,\alpha}(\mathbb{R})$ and

$$\|f\|_{L_{1,2\alpha+2-\beta,\alpha}} \leq b_\alpha^{1/p'} \|f\|_{\tilde{L}_{p,\alpha}}. \quad (3.17)$$

□

4. Some Applications

In this section, using the results of Section 3, we get the boundedness of the fractional D -maximal operator in the D -Morrey and modified D -Morrey spaces.

For $0 \leq \beta < 2\alpha + 2$, we define the fractional maximal functions

$$M_\beta f(x) := \sup_{t>0} (\mu_\alpha B(0,t))^{-1+\beta/(2\alpha+2)} \int_{B(0,t)} \tau_x |f|(y) d\mu_\alpha(y), \quad (4.1)$$

$$M_{p,\beta} f(x) := (M_\beta |f|^p)^{(1/p)}(x).$$

In the case $\beta = 0$, we denote $M_{p,0}f$ by $M_p f$. Note that $M_1 f = Mf$.

Lemma 4.1. *Let $1 \leq p < \infty$, $0 \leq \beta < 2\alpha + 2$, and $f \in L_{p,2\alpha+2-\beta,\alpha}(\mathbb{R})$. Then $M_{p,\beta} f \in L_\infty(\mathbb{R})$ and the following equality*

$$\|M_{p,\beta} f\|_{L_\infty} = b_\alpha^{(\beta/(2\alpha+2)-1)(1/p)} \|f\|_{L_{p,2\alpha+2-\beta,\alpha}} \quad (4.2)$$

is valid.

Proof.

$$\begin{aligned} \|M_{p,\beta} f\|_{L_\infty} &= b_\alpha^{(\beta/(2\alpha+2)-1)(1/p)} \sup_{x \in \mathbb{R}, t > 0} \left(t^{\beta-2\alpha-2} \int_{B(0,t)} \tau_x |f|^p(y) d\mu_\alpha(y) \right)^{1/p} \\ &= b_\alpha^{(\beta/(2\alpha+2)-1)(1/p)} \|f\|_{L_{p,2\alpha+2-\beta,\alpha}}. \end{aligned} \quad (4.3)$$

□

Taking $\beta = 0$ in Lemma 4.1 and using Lemma 3.3, we get for $M_p f$ the following result.

Corollary 4.2. *Let $1 \leq p < \infty$. Then*

$$\|M_p f\|_{L_\infty} = 4 \|f\|_{L_\infty}. \quad (4.4)$$

Lemma 4.3. *Let $1 \leq p < \infty$, $0 \leq \beta < 2\alpha + 2$, and $f \in \tilde{L}_{p,2\alpha+2-\beta,\alpha}(\mathbb{R})$. Then $M_{p,\beta} f \in L_\infty(\mathbb{R})$ and the following equality*

$$\|M_{p,\beta} f\|_{L_\infty} = b_\alpha^{(\beta/(2\alpha+2)-1)(1/p)} \|f\|_{\tilde{L}_{p,2\alpha+2-\beta,\alpha}} \quad (4.5)$$

is valid.

Corollary 4.4. Let $0 \leq \lambda < 2\alpha + 2$ and $0 \leq \beta < 2\alpha + 2 - \lambda$. Then the operator M_β is bounded from $L_{p,\lambda,\alpha}$ to L_∞ for $p = (2\alpha + 2 - \lambda)/\beta$. Moreover,

$$\|M_\beta f\|_{L_\infty} = b_\alpha^{\beta/(2\alpha+2)-1} \|f\|_{L_{1,2\alpha+2-\beta,\alpha}} \leq b_\alpha^{\beta/(2\alpha+2)-1/p} \|f\|_{L_{p,\lambda,\alpha}}. \quad (4.6)$$

Corollary 4.5. $1 \leq p < \infty$, $0 \leq \lambda < 2\alpha + 2$, $0 \leq \beta < 2\alpha + 2 - \lambda$. Then the operator M_β is bounded from $\tilde{L}_{p,\lambda,\alpha}$ to L_∞ for $(2\alpha + 2 - \lambda)/\beta \leq p \leq (2\alpha + 2)/\beta$. Moreover,

$$\|M_\beta f\|_{L_\infty} = b_\alpha^{\beta/(2\alpha+2)-1} \|f\|_{\tilde{L}_{1,2\alpha+2-\beta,\alpha}} \leq b_\alpha^{\beta/(2\alpha+2)-1/p} \|f\|_{\tilde{L}_{p,\lambda,\alpha}}. \quad (4.7)$$

Acknowledgment

The authors express their thank to the referees for careful reading, helpful comments and suggestions on the manuscript of this paper.

References

- [1] C. F. Dunkl, "Integral kernels with reflection group invariance," *Canadian Journal of Mathematics*, vol. 43, no. 6, pp. 1213–1227, 1991.
- [2] M. Rösler, "Bessel-type signed hypergroups on \mathbb{R} ," in *Probability Measures on Groups and Related Structures, XI (Oberwolfach, 1994)*, H. Heyer and A. Mukherjea, Eds., pp. 292–304, World Scientific, River Edge, NJ, USA, 1995.
- [3] C. B. Morrey Jr., "On the solutions of quasi-linear elliptic partial differential equations," *Transactions of the American Mathematical Society*, vol. 43, no. 1, pp. 126–166, 1938.
- [4] A. L. Mazzucato, "Besov-Morrey spaces: function space theory and applications to non-linear PDE," *Transactions of the American Mathematical Society*, vol. 355, no. 4, pp. 1297–1364, 2003.
- [5] M. E. Taylor, "Analysis on Morrey spaces and applications to Navier-Stokes and other evolution equations," *Communications in Partial Differential Equations*, vol. 17, no. 9-10, pp. 1407–1456, 1992.
- [6] C. Pérez, "Two weighted norm inequalities for Riesz potentials and uniform L_p -weighted Sobolev inequalities," *Indiana University Mathematics Journal*, vol. 39, no. 1, pp. 31–44, 1990.
- [7] A. Ruiz and L. Vega, "Unique continuation for Schrödinger operators with potential in Morrey spaces," *Publicacions Matemàtiques*, vol. 35, no. 1, pp. 291–298, 1991.
- [8] Z. Shen, "The periodic Schrödinger operators with potentials in the Morrey class," *Journal of Functional Analysis*, vol. 193, no. 2, pp. 314–345, 2002.
- [9] L. Caffarelli, "Elliptic second order equations," *Rendiconti del Seminario Matematico e Fisico di Milano*, vol. 58, pp. 253–284, 1988.
- [10] G. Di Fazio, D. K. Palagachev, and M. A. Ragusa, "Global Morrey regularity of strong solutions to the Dirichlet problem for elliptic equations with discontinuous coefficients," *Journal of Functional Analysis*, vol. 166, no. 2, pp. 179–196, 1999.
- [11] D. R. Adams, "A note on Riesz potentials," *Duke Mathematical Journal*, vol. 42, no. 4, pp. 765–778, 1975.
- [12] D. R. Adams, "Choquet integrals in potential theory," *Publicacions Matemàtiques*, vol. 42, no. 1, pp. 3–66, 1998.
- [13] F. Chiarenza and M. Frasca, "Morrey spaces and Hardy-Littlewood maximal function," *Rendiconti di Matematica e delle sue Applicazioni*, vol. 7, no. 3-4, pp. 273–279, 1987.
- [14] A. Kufner, O. John, and S. Fučík, *Function Spaces. Monographs and Textbooks on Mechanics of Solids and Fluids; Mechanics: Analysis*, Noordhoff International, Leyden, The Netherlands; Academia, Prague, Czech Republic, 1977.
- [15] M. Sifi and F. Soltani, "Generalized Fock spaces and Weyl relations for the Dunkl kernel on the real line," *Journal of Mathematical Analysis and Applications*, vol. 270, no. 1, pp. 92–106, 2002.
- [16] F. Soltani, " L_p -Fourier multipliers for the Dunkl operator on the real line," *Journal of Functional Analysis*, vol. 209, no. 1, pp. 16–35, 2004.

- [17] K. Trimèche, "Paley-Wiener theorems for the Dunkl transform and Dunkl translation operators," *Integral Transforms and Special Functions*, vol. 13, no. 1, pp. 17–38, 2002.
- [18] G. N. Watson, *A Treatise on the Theory of Bessel Functions*, Cambridge University Press, Cambridge, UK, 1966.
- [19] V. S. Guliyev and Y. Y. Mammadov, "On fractional maximal function and fractional integrals associated with the Dunkl operator on the real line," *Journal of Mathematical Analysis and Applications*, vol. 353, no. 1, pp. 449–459, 2009.
- [20] Y. Y. Mammadov, "Some embeddings into the Morrey spaces associated with the Dunkl operator on \mathbb{R} ," in *Proceedings of NASA, Embedding Theorems. Harmonic Analysis*, pp. 258–269, 2007.



Hindawi

Submit your manuscripts at
<http://www.hindawi.com>

