

## ORAL PRESENTATION

## Open Access

# Low reliability of the Risser sign in consecutive radiographs: a case series

Stefano Negrini\*, Salvatore Atanasio, Sabrina Donzelli, Monia Lusini, Salvatore Minnella, Francesco Negrini, Fabio Zaina

From 10th International Conference on Conservative Management of Spinal Deformities - SOSORT 2013 Annual Meeting  
Chicago, IL, USA. 8-11 May 2013

## Background

The low reliability of the Risser sign has been described in previous studies; however, the test/retest reliability of the Risser sign in different radiographs has not been examined or reported.

## Purpose

The goal of this report was to present data collected from studying a group of adolescent idiopathic scoliosis (AIS) patients whose Risser sign decreased in two consecutive X rays.

## Methods

Case series using the European Risser test.

## Results

By chance, a reduction of the Risser test was discovered in one patient. Subsequently, other cases have been searched to see if this was an exceptional situation. In one year, we found three more cases.

## Conclusions and discussion

At this stage, the following explanatory hypotheses can be drawn:

- Technical radiological differences (exposure, machine)
- Variation of pelvis positioning
- Postural changes influencing the pelvis
- Brace compression on the pelvis

According to our study results, all of the hypotheses include data both in favor and against the reliability of the Risser sign. Since the Risser sign is a 2D evaluation

**Table 1**

	1	2	3	4
Age	16.1	14.8	14.4	15.9
Height (cm)	0	+ 1	+1.5	0
Weight (kg)	+1	0	+1.5	0
Risser	4 to 2	1 to 0	2 to 1	4 to 3
Brace	Sforzesco	PASB	SpineCor	Sforzesco
Hours/day	23	12	20	16
Bracing months	4	6	4	22
Months between x-rays	7	6	6	14
In brace x-ray	No	No	Yes	No
Change of x-ray machine	No	No	Yes	Yes
°Cobb	-23	-4	-7	+4
Change of rotation of the pelvis	++	+	?	+
C7+L3 distance (cm)	-3	0	+0.5	-0.5
ATR (°)	-8	-3	-3	+3

of a 3D phenomenon, pelvis repositioning could perhaps be the most plausible explanation.

This case series is open to the possibility that the Risser sign is even less reliable than originally considered. Unfortunately, this result cannot be checked experimentally due to ethical reasons. Nevertheless, observational designs could be considered in the future.

Published: 18 September 2013

## References

1. Hammond KE, Dierckman BD, Burnworth L, Meehan PL, Oswald TS: **Inter-observer and intra-observer reliability of the Risser sign in a metropolitan scoliosis screening program.** *J Pediatr Orthop* 2011, **31**(8):e80-e84.
2. Kotwicki T: **Improved accuracy in Risser sign grading with lateral spinal radiography.** *Eur Spine J* 2008, **17**(12):1676-1685.

\* Correspondence: [stefano.negrini@isico.it](mailto:stefano.negrini@isico.it)  
University of Brescia - IRCCS Don Gnocchi, Milan, Italy

3. Reem J, Carney J, Stanley M, Cassidy J: **Risser sign inter-rater and intra-rater agreement: is the Risser sign reliable?** *Skeletal Radiol* 2009, **38**(4):371-375.

doi:10.1186/1748-7161-8-S2-O18

**Cite this article as:** Negrini *et al.*: Low reliability of the Risser sign in consecutive radiographs: a case series. *Scoliosis* 2013 **8**(Suppl 2):O18.

**Submit your next manuscript to BioMed Central  
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at  
[www.biomedcentral.com/submit](http://www.biomedcentral.com/submit)

