

Animal (2016), 10:11, pp 1899–1910 © The Animal Consortium 2016
doi:10.1017/S1751731116000707



An approach to holistically assess (dairy) farm eco-efficiency by combining Life Cycle Analysis with Data Envelopment Analysis models and methodologies

A. D. Soteriades^{1†}, P. Faverdin^{2,3}, S. Moreau⁴, T. Charroin⁴, M. Blanchard⁵ and A. W. Stott¹

¹Scotland's Rural College, Future Farming Systems Group, Edinburgh, UK; ²INRA, UMR 1348 PEGASE, F-35590 St-Gilles, France; ³Agrocampus-Ouest, UMR 1348 PEGASE, F-35000 Rennes, France; ⁴Institut de l'Élevage, F-75000 Paris, France; ⁵CIRAD, UMR 0868 SELMET, Bobo-Dioulasso, Burkina Faso

(Received 20 October 2015; Accepted 21 March 2016; First published online 29 April 2016)

Eco-efficiency is a useful guide to dairy farm sustainability analysis aimed at increasing output (physical or value added) and minimizing environmental impacts (EIs). Widely used partial eco-efficiency ratios (EIs per some functional unit, e.g. kg milk) can be problematic because (i) substitution possibilities between EIs are ignored, (ii) multiple ratios can complicate decision making and (iii) EIs are not usually associated with just the functional unit in the ratio's denominator. The objective of this study was to demonstrate a 'global' eco-efficiency modelling framework dealing with issues (i) to (iii) by combining Life Cycle Analysis (LCA) data and the multiple-input, multiple-output production efficiency method Data Envelopment Analysis (DEA). With DEA each dairy farm's outputs and LCA-derived EIs are aggregated into a single, relative, bounded, dimensionless eco-efficiency score, thus overcoming issues (i) to (iii). A novelty of this study is that a model providing a number of additional desirable properties was employed, known as the Range Adjusted Measure (RAM) of inefficiency. These properties altogether make RAM advantageous over other DEA models and are as follows. First, RAM is able to simultaneously minimize EIs and maximize outputs. Second, it indicates which EIs and/or outputs contribute the most to a farm's eco-inefficiency. Third it can be used to rank farms in terms of eco-efficiency scores. Thus, non-parametric rank tests can be employed to test for significant differences in terms of eco-efficiency score ranks between different farm groups. An additional DEA methodology was employed to 'correct' the farms' eco-efficiency scores for inefficiencies attributed to managerial factors. By removing managerial inefficiencies it was possible to detect differences in eco-efficiency between farms solely attributed to uncontrollable factors such as region. Such analysis is lacking in previous dairy studies combining LCA with DEA. RAM and the 'corrective' methodology were demonstrated with LCA data from French specialized dairy farms grouped by region (West France, Continental France) and feeding strategy (regardless of region). Mean eco-efficiency score ranks were significantly higher for farms with <10% and 10% to 30% maize than farms with >30% maize in the total forage area before correcting for managerial inefficiencies. Mean eco-efficiency score ranks were higher for West than Continental farms, but significantly higher only after correcting for managerial inefficiencies. These results helped identify the eco-efficiency potential of each region and feeding strategy and could therefore aid advisors and policy makers at farm or region/sector level. The proposed framework helped better measure and understand (dairy) farm eco-efficiency, both within and between different farm groups.

Keywords: eco-efficiency, composite indicators, managerial inefficiency, uncontrollable factors, French dairy farm data

Implications

Dairying contributes significantly to society (employment, economy, nutritional value of dairy products, etc.) at the cost of several environmental impacts. Therefore, improvements in dairy farm 'eco-efficiency' are essential to ensure more output with fewer impacts. This study introduced a modelling framework to measure, analyse and understand dairy farm eco-efficiency in much more depth than previously published assessments.

The framework was demonstrated with data from French specialized dairy farms. This framework can be a powerful tool for improving the sustainability of dairy farming systems, especially when multiple, conflicting objectives (multiple-output maximization v. multiple-impact minimization) are involved.

Introduction

Facing the environmental impacts (EIs) of agriculture, the challenge to satisfy the demands of a growing and more affluent global population, the scarcity of natural resources

† E-mail: andreas.soteriades@sruc.ac.uk

and the consequences of climate change, agricultural policies are increasingly directed towards 'sustainable intensification' of agriculture (Foresight, 2011). Consequently, the dairy industry (along with other sectors) is required to comply with several policies promoting environmentally sustainable and resource use-efficient production (Casey and Holden, 2005). This necessitates the application of tools to measure dairy farm performance in terms of resource use efficiency and productivity, increased product quantity and value and minimization of EIs. Such a tool is 'eco-efficiency', originally developed for the business sector; it is expressed as a ratio of product or service value to EI (Economic and Social Commission for Asia and the Pacific [ESCAP], 2009).

In dairy studies, eco-efficiency is usually expressed as the ratio of an EI per some functional unit such as kg milk or ha land (e.g. Casey and Holden, 2005; van Calker *et al.*, 2008; Basset-Mens *et al.*, 2009; Guerci *et al.*, 2013; Bava *et al.*, 2014). To calculate the EIs dairy studies (including the aforementioned) are increasingly using Life Cycle Analysis (LCA), an internationally standardized method for estimating the EIs of agricultural products from a global perspective (Bava *et al.*, 2014). Using LCA, some studies have been confined to comparing different dairy systems in terms of several eco-efficiency indicators defined by two or more functional units (e.g. Basset-Mens *et al.*, 2009). Others have examined the relationships between eco-efficiency ratios and related factors (e.g. farming intensity, farm self-sufficiency) by employing multivariate methods such as regression (Casey and Holden, 2005) and principal component analysis (Bava *et al.*, 2014). Other studies have focussed on expressing the relative importance of several eco-efficiency indicators based on different stakeholder weighting schemes (see van Calker *et al.*, 2008).

There are six main comments to be made on the approaches to dairy farm eco-efficiency in the aforementioned studies. First, analyses involving multiple partial eco-efficiency ratios ignore the substitution possibilities that might exist between different EIs. That is, farms performing moderately for several EIs tend to be overlooked in favour of farms performing exceptionally well for one EI (Kuosmanen and Kortelainen, 2005). Second, with these ratios the allocation of EIs to products is challenging as dairy farms generally produce other products too, such as meat. Third, incommensurability between several criteria expressed by multiple eco-efficiency ratios rather than a single performance index can complicate decision making (Kuosmanen and Kortelainen, 2005). Fourth, analyses with methods such as regression and principal component analysis are subject to the method chosen to normalize/standardize eco-efficiency ratios expressed in different units. Fifth, assigning subjective weights to indicators (e.g. the eco-efficiency ratios) has been debated in the literature (Kuosmanen and Kortelainen, 2005). Sixth, allowance should be made for the fact that there exist factors affecting eco-efficiency that are beyond managerial control, such as the different bio-physical conditions under which farms operate (see Bogetoft and Otto, 2011; Jan *et al.*, 2012).

All six aforementioned limitations can be overcome with the productive efficiency method of Data Envelopment Analysis (DEA; see Cooper *et al.*, 2007), employed in this study. DEA is a relative, multiple-input, multiple-output efficiency measurement method calculating single aggregated efficiency indices for each dairy farm by assessing the whole production system, including EIs. Importantly, with DEA no allocation of EIs to specific products is required because the farm is assessed as a whole, multiple-input, multiple-output entity. Most DEA models are not affected by the different measurement units of the data and their weighting schemes are endogenous, that is, 'data-driven' (e.g. the model of Cooper *et al.*, 1999 employed in this study). DEA methodologies correcting for managerial inefficiencies and accounting for uncontrollable factors are available, such as that of Brockett and Golany (1996) adopted in this study.

DEA has been applied in several dairy studies for the calculation of eco-efficiency. For example, Jan *et al.* (2012) and subsequently Pérez Urdiales *et al.* (2015) used the DEA eco-efficiency model of Kuosmanen and Kortelainen (2005) to define a dairy farm eco-efficiency ratio. This ratio equalled the amount of (physical or monetary) dairy farm output to an aggregate EI index calculated as a weighted summation of all EIs considered in their study. This ratio was then maximized by minimizing the aggregate EIs for the given production levels. Importantly, the EIs in Jan *et al.* (2012) were LCA-derived. In fact, efficiency studies are increasingly recognizing the advantages of combining LCA with DEA as the former can capture EIs using detailed, cradle-to-grave data (e.g. land use required for the production of feed imported in the dairy farm plus on-farm land use), while the latter has the aforementioned advantages (Vázquez-Rowe and Iribarren, 2015).

The objective of this study was to propose a framework combining LCA with DEA that not only overcomes the six aforementioned issues, but also improves the measurement and understanding of farm eco-efficiency using dairying as exemplar. This will guide farming practice to greater yet sustainable production (sustainable intensification) as advocated for example by the UK Foresight report (2011). The DEA model employed, known as the range adjusted measure (RAM) of inefficiency (Cooper *et al.*, 1999), has several desirable properties, for example it allows for the ranking of farms in terms of eco-efficiency performance. Moreover, it seeks to maximize eco-efficiency by simultaneously minimizing EIs and maximizing production. Furthermore, it can identify the factors contributing the most to inefficiency, such as excess EIs and/or under-produced outputs. A method to isolate managerial inefficiency from uncontrollable factors was also demonstrated. That way, it was possible to compare different dairy systems in terms of eco-efficiency solely under the influence of uncontrollable, rather than managerial, factors. The exercise was run using detailed LCA data for French specialized dairy farms. Region was considered as the uncontrollable factor in this study due to the remarkable differences between West and Continental

France in terms of farm structure and bio-physical conditions (Gac *et al.*, 2010b). The results helped identify the eco-efficiency potential of each region and feeding strategy and could therefore aid advisors and policy makers at farm or region/sector level.

Material and methods

Data

LCA was used to estimate several important midpoint impacts of dairy farming systems. It was conducted using the DIAPASON database resulting from a partnership involving voluntary participation of farmers, the Chambers of Agriculture (France) and the French Livestock Institute. This database contains detailed information on technical and economic operations of nearly 500 farms each year throughout France (Charroin *et al.*, 2005).

Environmental performance was assessed by indicators of pressure from agricultural activity on the environment considering midpoint impact indicators of LCA. The frontier of the farm system was limited to the farm, considered as a system dedicated to agricultural products (crops, milk, meat) at farm gate. Impacts associated with these products beyond the farm gate were not considered in this study. The limits of the system included the whole farm and all the inputs of the farming system. The system and its main processes are described in Figure 1.

The different EIs considered in this study were midpoint impacts consistent with the CML 2001 methodology (Guinée *et al.*, 2002) with some specific equations to estimate the emissions. They concern global warming potential and non-renewable energy according to the greenhouse gas emissions GES'TIM methodology (Gac *et al.*, 2010a) and non-renewable energy use (Béguin *et al.*, 2008) and based on the Fourth Assessment Report of the Intergovernmental Panel on Climate Change (IPCC, 2007). Eutrophication was calculated as a unique impact according to the CML 2001 methodology (Guinée *et al.*, 2002) and

acidification using equations from the European Monitoring and Evaluation Programme/Core Inventory of Air Emissions in Europe (EMEP/CORINAIR, 2002). Table 1 summarizes the inventory of all the emissions considered to calculate the different impacts.

The factors applied to the nitrogen (N), phosphorus and carbon fluxes (calculated with the DIAPASON database), generated estimates of EIs. Dry matter intake and mineral excretion in the faeces and urine of animals were calculated according to physiological needs (milk production, weight after calving) using equations proposed by CORPEN (Comité d'orientation pour de pratiques agricoles respectueuses de l'environnement, 1999) taking into account the farmers' feeding practices (types of forages and concentrates). The carbon (C) storage of permanent grassland that was taken into account was up to 500 kg C/ha per year (Gac *et al.*, 2010b). On-farm N leaching was estimated using the N farm surplus, including symbiotic fixation (based on a fixed proportion of legumes for permanent grassland), but after removing losses of ammonia and organic N storage in soils assumed as 10% of C storage (with C : N ratio of 10), which represents 50 kg N/ha per year in permanent pasture. The impact values of inputs were derived from the LCA database 'ecoinvent' (Nemecek and Kägi, 2007) and Gac *et al.* (2010a). Because the whole farm was chosen as the functional unit, all farm products were considered simultaneously in this analysis, therefore no allocation of emissions to the different products was applied.

Finally, 185 dairy farms (specialized dairy farms according to the widely recognized Farm Accounts Data Network (FADN) typology) located in different French lowland regions in 2007 and 2008 were kept in this study. The different farms were classified into two main groups according to climate zone and specialization: oceanic specialized systems (OSS; West France, consisting of the following regions: Basse-Normandie, Bretagne, Haute-Normandie, Pays de la Loire, Poitou-Charente) and continental specialized systems (CSS; Continental France, consisting of the following regions: Alsace, Centre, Champagne-Ardenne, Franche-Comté,




Figure 1 Description of the dairy farming system used for the Life Cycle Analysis (LCA) calculations.

Table 1 On-farm emissions due to different sources of the farming system

	Source and emissions																						
	Fertilization						Other sources																
	Enteric		Building		Storage		Organic		Min.		Grazing		Crop residue		Indirect		Combustion						
CH ₄	N ₂ O	NH ₃	NO	CH ₄	N ₂ O	NH ₃	NO	N ₂ O	NO	NH ₃	NO	CH ₄	N ₂ O	NH ₃	NO	N ₂ O	CO ₂	SO ₂	PO ₄	N leaching (NO ₃)	P runoff (PO ₄)		
Milk yield (fat %)	X																						
Livestock units	X																						
% time indoors		X																					
Building type		X																					
Manure type			X																				
Animal N exc.				X																			
MS (time and volume)					X																		
N/P fertilization									X														
Energy use																		X					
Inputs																			X				
Farm N surplus																				X			
Crop rot.																					X		

Min. = mineral; N = nitrogen; P = phosphorous; Animal N exc. = animal nitrogen excretion; MS = manure storage; Crop rot. = crop rotation.

Lorraine, Rhône-Alpes). The second dimension of the typology, crossed with the first dimension, concerned the type of feeding strategies, based on the area of maize silage in the total forage area of the farm: <10%, 10% to 30%, >30% maize. Other farm classes were not considered due to insufficient number of farms in the class. Table 2 summarizes the five EIs and three outputs used in this study per system for the years 2007 and 2008.

Data envelopment analysis

DEA is a non-stochastic, non-parametric technique that benchmarks different decision-making units (DMUs) performing the same task in terms of their capacity to convert inputs into outputs. DEA calculates dimensionless and aggregated efficiency indices without requiring *a priori* assumptions on the importance of each variable for the DMUs' performance, making it a particularly attractive multiple-criteria tool. DEA constructs an efficient frontier, that is, a convex, piece-wise linear surface over observed data points against which all DMUs are benchmarked (or 'enveloped'). Figure 2 represents an efficient frontier ABC for the single-EI, single-output case. The efficient frontier comprises of the best performers (DMUs A, B and C in Figure 2) and the performance of all other DMUs (e.g. DMU D in Figure 2) is evaluated by deviations from the frontier line (Cooper *et al.*, 2007). This is a fundamental difference between DEA and methods such as regression as the latter reflects 'average' or 'central tendency' behaviour (Cooper *et al.*, 2007) and is unable to provide a holistic characterization of DMUs within a multiple-objective assessment. Convexity in DEA allows for the interpolation from observed DMUs to 'virtual' DMUs with input-output profiles between the observations, allowing us to rely on fewer actual observations. These 'virtual' DMUs are derived as convex combinations of inputs and outputs of observed DMUs. Convexity can be illustrated in Figure 2 as follows. Any line connecting any two points belonging to, or being placed below, the frontier would also be placed on or below the frontier, and never outside this space (i.e. above the frontier). The points these lines comprise of can represent both observed and 'virtual' DMUs. See Bogetoft and Otto (2011) for a theoretical background on convexity in DEA.

Data envelopment analysis in the eco-efficiency context

As mentioned in the introduction, eco-efficiency measurement with DEA is advantageous for three main reasons: (i) several EIs are aggregated into a single index, (ii) substitution possibilities between EIs are not left unaccounted for and (iii) no allocation of EIs to specific outputs is required. Points (i) to (iii) can be expressed in the DEA context by minimizing the denominator of the following ratio:

$$\text{Eco-efficiency} = \max \left\{ \frac{\text{Output}}{\text{Weighted sum of EIs}} \right\} \quad (1)$$

subject to a number of constraints (see Kuosmanen and Kortelainen, 2005). In ratio 1 the output can be expressed in monetary or physical terms. The weights summing the various EIs are calculated by the DEA model itself so one

Table 2 Statistics of dairy farm environmental impacts and outputs per system, in both years 2007 and 2008

Data	CSS (<i>n</i> = 59)				OSS (<i>n</i> = 126)			
	Min	Max	Mean	SD	Min	Max	Mean	SD
EI								
Non-renewable energy (10 ³ MJ)	580	5256	1643	846	343	4223	1406	709
Land use (ha)	48	351	133	67	48	268	101	43
Eutrophication (kg PO ₄)	625	10 890	3200	2241	425	10 070	3200	2058
Acidification (kg SO ₂)	2189	11 780	4728	1982	1543	8413	3798	1419
GWP (kg CO ₂)	163 500	1 431 000	535 000	257 097	91 400	1 330 000	507 200	218 404
Outputs								
Milk (kg protein)	2210	10 540	5218	1957	2080	10 900	5195	1907
Meat (kg live weight)	0	73 410	21 700	13 401	0	92 210	23 330	11 644
Crops (10 ³ MJ)	614	10 930	3488	2683	0	8152	2142	1848

CSS = continental specialized systems; OSS = oceanic specialized systems; EI = environmental impact; GWP = global warming potential.




Figure 2 An efficient frontier ABC in the case of a single environmental impact (EI) and a single output. Inefficient decision-making unit D seeks maximal EI reduction and output expansion and thus is projected on ABC at point B.

need not rely on subjective, pre-defined weight choices for the importance of each EI. Specifically, the DEA model maximizing ratio 1, selects the most self-favourable weights for each DMU so that its eco-efficiency is maximized. These weights cancel out the (often) different measurement units of the EIs, making the DEA model 'units invariant' (Cooper *et al.*, 2007).

Despite its usefulness, there are two main limitations with the eco-efficiency DEA model of Kuosmanen and Kortelainen (2005). First, ratio 1 can only be maximized by minimizing the EIs for the *given* output levels. In other words, simultaneous minimization of EIs and maximization of output is not possible. For example, DMU D in Figure 2 would have to move horizontally towards the frontier to become efficient, ignoring any potential increases in its output. Because eco-efficiency expresses the idea of firms (e.g. dairy farms)

providing 'more' to society with less EIs, it is desirable to use a DEA model allowing for simultaneous adjustments in EIs and output. Second, full eco-efficiency can only be achieved by minimizing all EIs by the *same* proportion. A DEA model should be able to identify those EIs generating the most detrimental excess (or 'slack' in the DEA terminology) to a DMU's eco-'inefficiency'.

Both aforementioned limitations can be overcome with the use of so-called 'additive' DEA models (see Cooper *et al.*, 2007). These models are able to simultaneously, and non-proportionally, minimize EIs and maximize output for a given DMU. In such a case, DMU D in Figure 2 would move towards point B. The term 'additive' is attributed to the fact that these models' objective functions involve summations of all input and output slacks in order to identify all potential sources of inefficiency. In Figure 2 this summation is

represented by the vector heading from point D towards point B and equals the maximal sum of the EI slack and the Output slack. As will be shown below, this summation of all slacks in the objective function departs from the ratio form of ratio 1. However, it is consistent with the idea of maximizing output while minimizing EIs and thus has been adopted in past eco-efficiency studies (see Ramli and Munisamy, 2015 and the related studies they cite). This study employed the RAM additive model (Cooper *et al.*, 1999), presented below. RAM and its variants have been used in several eco-efficiency studies of industries other than dairy, see Ramli and Munisamy (2015).

Range adjusted measure of inefficiency. Suppose that there are n DMUs (e.g. dairy farms) each using m inputs (or EIs in the case of this study) to produce s outputs, denoted as x_i ($i = 1, \dots, m$) and y_r ($r = 1, \dots, s$), respectively. The RAM inefficiency score of the j th DMU, denoted as DMU_o , is given by the following linear programme (Cooper *et al.*, 1999):

$$\rho^* = \max_{\lambda_j, s_{io}, s_{ro}} \frac{1}{m+s} \left(\sum_{i=1}^m \frac{s_{io}}{R_i} + \sum_{r=1}^s \frac{s_{ro}}{R_r} \right) \quad (2)$$

subject to

$$x_{io} = \sum_{j=1}^n x_{ij}\lambda_j + s_{io} \quad (i = 1, \dots, m)$$

$$y_{ro} = \sum_{j=1}^n y_{rj}\lambda_j - s_{ro} \quad (r = 1, \dots, s)$$

$$\sum_{j=1}^n \lambda_j = 1$$

$$\lambda_j, s_{io}, s_{ro} \geq 0$$

where x_{io} and y_{ro} are the inputs and outputs of DMU_o respectively; s_{io} and s_{ro} are the input and output slacks, respectively (Note: input slacks represent overused inputs, i.e. DMU_o could have produced the same amount of output using less input. Output slacks represent output shortfalls, i.e. DMU_o could have produced more output given its current input use.); λ_j is a scalar which, when positive, indicates that DMU_j has been used as a reference (i.e. benchmark) by DMU_o ; and $R_i = \max_j \{x_{ij}\} - \min_j \{x_{ij}\}$, $R_r = \max_j \{y_{rj}\} - \min_j \{y_{rj}\}$ represent the ranges in inputs and outputs, respectively, common across *all* DMUs. The ranges act as a 'data-driven' weighting scheme, a more objective one compared to methods where the weights are (subjectively) pre-defined by the user. These weights normalize the slacks and make RAM units invariant. The objective function represents the average proportion of the inefficiencies that the ranges show to be possible in each input and output (Cooper *et al.*, 1999).

The constraint $\sum_{j=1}^n \lambda_j = 1$ is the 'variable returns-to-scale' specification (see Cooper *et al.*, 2007) which ensures that a farm is only compared to farms of similar size. This specification was desirable in this study as DEA works with absolute values rather than ratios.

Model 2 is run n times, once for each DMU. When DMU_o is efficient all its slacks equal zero as this means that it does not need to further reduce its inputs and increase its outputs to become efficient (e.g. DMUs A, B and C in Figure 2). In this case RAM inefficiency ρ^* in model 2 equals 0, indicating that DMU_o is 100% efficient. If DMU_o is inefficient, one can identify through the slack values (which in this case are non-proportional) the inputs and desirable outputs contributing the most to its inefficiency. For an inefficient DMU (e.g. DMU D in Figure 2) any choice of input resulting in $x_{io} > \sum_{j=1}^n x_{ij}\lambda_j$ means that with some combination of inputs *other* DMUs (identified by the non-zero λ_j values) could have improved this input in amount by $s_{io} = x_{io} - \sum_{j=1}^n x_{ij}\lambda_j$ without worsening any other input or output (Brockett *et al.*, 2004). Consider, for example a DMU on ABC with coordinates (2.7, 3) as opposed to DMU D with coordinates (7, 3) in Figure 2. The same applies for the desirable outputs and their shortfalls $s_{ro} = \sum_{j=1}^n y_{rj}\lambda_j - y_{ro}$. In this case consider a DMU with coordinates (7, 5.7) as opposed to DMU D in Figure 2. In either case RAM inefficiency ρ^* is greater than 0, indicating that DMU_o is inefficient.

Because $\sum_{j=1}^n \lambda_j = 1$ in model 2 it follows that $s_{io} = \sum_{j=1}^n (x_{io} - x_{ij})\lambda_j \leq \sum_{j=1}^n R_i\lambda_j = R_i$ and similarly $s_{ro} \leq R_r$ and thus $0 \leq \rho^* \leq 1$. Hence, the measure of *inefficiency* ρ^* in model 2 can be easily converted to a measure of *efficiency* as follows:

$$\text{RAM efficiency} = 1 - \rho^* \quad (3)$$

RAM efficiency 3 is bounded by 0 and 1. Unity indicates that the DMU under evaluation is efficient while values < 1 imply that it is inefficient.

Two very attractive properties of RAM are the following: (i) RAM uses the ranges as a common weighting scheme across *all* DMUs; and (ii) RAM is strongly monotone in the slacks, that is, holding any other inputs and outputs constant, an increase (decrease) in any of its inputs (outputs) will increase the inefficiency score for an inefficient DMU. Model 1 does not carry properties (i) to (ii).

Properties (i) to (ii) allow for a full ranking of inefficient DMUs in terms of their RAM efficiency score 3 (Cooper *et al.*, 1999). (Not all DEA models carry this property. For example, with ratio 1 one cannot say that a DMU with a score of 0.8 is more eco-efficient than a DMU with a score of 0.7 because the EI weights are DMU-specific and will generally differ between DMUs.) This was strongly desirable in the current study so as to determine whether farms ranked higher in terms of eco-efficiency in a specific region or under a certain feeding strategy.

DEA variables. This study used the five EIs and three outputs in Table 2 for the calculation of eco-efficiency with RAM,

namely non-renewable energy use, land use, eutrophication, acidification, global warming potential and milk, meat and crop production. With DEA, increasing the number of variables also increases the number of efficient DMUs which can be quite problematic with small sample sizes. A rough rule of thumb is to choose $n \geq \max\{m \times s, 3 \times (m + s)\}$ (Cooper *et al.*, 2007, p. 116). The rule of thumb was satisfied in this study: $n = 185 \geq \max\{m \times s, 3 \times (m + s)\} = 24$.

Testing for differences in eco-efficiency between regions and feeding strategies

Differences in dairy farm eco-efficiency scores between regions and feeding strategies were tested for using the non-parametric Kruskal–Wallis test (see Conover, 1999), also known as ‘non-parametric Kruskal–Wallis one-way ANOVA by ranks’ (Sheskin, 1997). The Kruskal–Wallis test is employed with ordinal (rank-order) data in hypothesis testing involving a design with two or more independent samples (Sheskin, 1997). That is, dairy farms were ranked in terms of their eco-efficiency scores and differences between groups were tested based on each group’s average rank. The null hypothesis is that all of the populations are identical against the alternative that at least one of the populations tends to yield larger observations than at least one of the other observations (Conover, 1999). When at least three groups are compared the Kruskal–Wallis test cannot indicate which pairs of groups significantly differ (provided that significant differences occur). The *post*-Kruskal–Wallis non-parametric rank test known as Dunn’s test (see Sheskin, 1997) was therefore employed to identify specific differences between the three feeding strategies.

Choosing non-parametric tests over the parametric one-way ANOVA and its *post hoc* tests was done for two reasons. First, the theoretical distribution of efficiency scores in DEA is generally unknown so a convention in the DEA literature is to use non-parametric tests (Brockett and Golany, 1996; Cooper *et al.*, 2007; Bogetoft and Otto, 2011). Second, because RAM can be used to rank DMUs, it lends itself to the rankings that underlie non-parametric rank statistics (Brockett *et al.*, 2004). Both tests employed in this study operate based on the rank transformation approach; that is, the data are replaced by their ranks and then the usual parametric tests (e.g. *t* test, *F* test, etc.) are applied on the ranks. (Tied observations [e.g. when at least two DMUs are eco-efficient] are given the average rank of the tied scores.) Therefore, these tests are not affected by outliers or skewed data. See Conover (1999).

Examining the effect of region on eco-efficiency

The bio-physical conditions under which dairy farms operate largely differ between West and Continental France. Regional differences in eco-efficiency were therefore tested. It would seem appropriate to pool farms from both regions in one dataset, run the RAM model and then test for differences between regions with the Kruskal–Wallis test. Such practice, however, would reveal any differences between regions under the *observed* levels of EIs and output (i.e. the EI

and output values outlined in Table 2). This means that inefficiencies attributed to both managerial and regional factors would not allow inefficient farms to operate under their full potential. Indeed, the risk of amalgamating both sources of inefficiency (managerial and regional) is to grant inadvertently some bad managers (farmers) good eco-efficiency scores when they are only benefitting from operating under particularly favourable bio-physical conditions (see Brockett and Golany, 1996). Removing EI and output managerial inefficiencies (i.e. slacks) was therefore essential before comparing the two regions in terms of eco-efficiency. This was done by adopting the methodology of Brockett and Golany (1996) which involved the following four steps:

1. Run two separate DEA exercises, one for CSS only and one for OSS only with model 2.
2. Using the optimal EI and output slacks obtained from the previous step make the necessary reductions in EIs and outputs so that inefficient DMUs in each group become efficient. This is done using the following formulas:

$$\begin{aligned}\hat{x}_{io} &= x_{io} - s_{io}^* \quad (i = 1, \dots, m) \\ \hat{y}_{ro} &= y_{ro} + s_{ro}^* \quad (r = 1, \dots, s)\end{aligned}\quad (4)$$

where the asterisks (*) denote optimality. (For example, let us assume that Figure 2 represents OSS farms. With formulas 4 the OSS farm D would have been projected onto the OSS efficient frontier at point B.) Now managerial inefficiency has been eliminated within OSS and CSS and both are operating ‘up to the boundary of the capabilities which the evidence showed was possible for [OSS and CSS]’ (Cooper *et al.*, 2007, p. 238).

3. Pool all DMUs deriving from the previous step and run a new DEA exercise with model 2.
4. Test for significant differences between the systems’ efficiency scores using non-parametric rank statistics, i.e. the Kruskal–Wallis test.

Following the steps above it was possible to compare the two regions in terms of eco-efficiency. It should be noted, however, that the DMUs were then evaluated not based on their *actual* levels of EIs and output, but on their *efficient* ones. Because this methodology corrects for any managerial inefficiencies present in DMUs, from this point it is referred to as the ‘corrective’ methodology.

Putting all methods together

Figure 3 summarizes the methodology employed in this study. Phase 1 did not apply the ‘corrective’ methodology and involved two steps. In Step 1.1 the EIs and outputs for each farm were fed into RAM and the eco-efficiency scores were obtained. Note that in this step DMUs from *both* CSS and OSS were pooled before the RAM was run. Step 1.2a tested for differences in eco-efficiency scores between the two systems and between the three feeding strategies with non-parametric rank tests. Moreover, the EI and output




Figure 3 Description of the modelling framework adopted in this study. DEA: Data Envelopment Analysis. LCA = Life Cycle Analysis.

slacks were compared between systems in Step 1.2b. Phase 2 applied the 'corrective' methodology and involved four steps. In Step 2.1 the RAM model was run for *each* system (CSS, OSS). In Step 2.2 the EIs and outputs of each farm in each system were projected onto their efficient levels with the formulae in 4. In Step 2.3 the RAM model was re-run for the *whole* sample (both CSS and OSS) using the projected data from Step 2.2. Step 2.4 tested for differences in the new eco-efficiency scores between the two systems and between the three feeding strategies with non-parametric rank tests. Unlike Phase 1, in Phase 2, systems and feeding strategies were exposed to the full eco-efficiency potential that the data showed to be possible for these groups.

There are distinct differences between Phase 1 and 2. Although Phase 1 did not differentiate between regional and managerial factors, it helped to evaluate the 185 French specialized farms under their *observed* levels of EIs and outputs, as reported in Table 2. In other words, Phase 1 evaluated farms 'as they *actually* performed' and not 'as they *could* be performing', as in the 'corrective' methodology described in Phase 2. Phase 1 is therefore useful for efficiency comparisons between and within farms in terms of the whole population, without correcting for potential systematic differences between groups (defined by region in this case). Phase 2 is appropriate for testing the hypothesis that systematic unavoidable differences between groups will affect efficiency performance. Phases 1 and 2 are therefore independent but complementary. See Brockett *et al.* (2004) who also conducted their analysis in two stages analogous to the two Phases employed here.

All calculations were run with the R language (<http://www.R-project.org/>). The R function for RAM was developed by the first author of this article. The Kruskal–Wallis test is available in the standard version of R. Dunn's test is available by the R package 'dunn.test' (<https://cran.r-project.org/web/packages/dunn.test/dunn.test.pdf>).

Results

Eco-efficiency scores and slacks per system and feeding strategy when accounting for managerial inefficiencies
The results for the eco-efficiency scores and slacks presented in this sub-section were calculated *before* applying the 'corrective' methodology (Phase 1 in Figure 3).

Eco-efficiency scores. Statistics for the eco-efficiency scores and their mean ranks per system and feeding strategy are presented in Table 3. The mean, median and mean ranks of eco-efficiency scores were higher for OSS than CSS. However, the Kruskal–Wallis test did not identify significant differences between CSS and OSS in terms of the eco-efficiency scores' mean ranks ($P = 0.105$). The three feeding strategies ranked as follows in terms of mean, median and mean ranks of eco-efficiency scores: (<10% maize) > (10% to 30% maize) > (>30% maize). The Kruskal–Wallis test identified significant differences between the three feeding strategies in terms of the eco-efficiency scores' mean ranks ($P = 0.001$). Specific differences were identified with Dunn's test. Differences were significant between DMUs with <10% maize and >30% maize in the total forage area ($P < 0.001$) and between DMUs with 10% to 30% maize and >30% maize ($P = 0.011$). No differences were found between DMUs with >10% maize and 10% to 30% maize in the total forage area ($P = 0.083$).

EI and output slacks. Table 4 summarizes the optimal EI and output slacks from model 2 per system, expressed as proportions of their respective ranges i.e. s_{io}^*/R_i ($i = 1, \dots, m$) and s_{ro}^*/R_r ($r = 1, \dots, s$). That way, it was possible to 'decompose' the eco-efficiency scores in Table 3 in order to detect the EIs and outputs with the highest relative contribution to a DMU's inefficiency. (Averaging each system's input and output inefficiencies in Table 4 and then subtracting them from 1 equals the mean efficiency scores presented in Table 3.) The EIs with the highest contribution to CSS systems' inefficiency were eutrophication potential, land use and acidification potential. By contrast, eutrophication potential was the EI with the by-far-largest contribution to OSS systems' inefficiency. In terms of output inefficiency, meat and milk were by far the largest contributors to the inefficiency of both OSS and CSS. Notably, for both OSS and CSS the mean input inefficiencies were much higher than the mean output inefficiencies.

Eco-efficiency scores per system and feeding strategy after eliminating managerial inefficiencies
The eco-efficiency results per system and feeding strategy presented in this section were obtained *after* eliminating all

Table 3 Statistics for eco-efficiency scores per system and feeding strategy before removal of managerial inefficiencies

	Eco-efficiency scores					
	Min	Max	Median	Mean	SD	Mean rank
System						
CSS	0.840	1.000	0.934	0.938	0.047	83.814
OSS	0.762	1.000	0.950	0.949	0.050	97.302
Feeding strategy						
<10% maize ¹	0.841	1.000	0.966	0.964	0.038	113.795 ^a
10% to 30% maize ¹	0.840	1.000	0.954	0.950	0.045	98.596 ^a
>30% maize ¹	0.762	1.000	0.930	0.932	0.053	78.310 ^b

CSS = continental specialized systems; OSS = oceanic specialized systems.
^{a,b}Values within a column with different superscripts differ significantly at $P < 0.05$.

¹Maize area as % of total forage area on farm.

Table 4 Mean slack values per system expressed as a proportion of their corresponding ranges

	CSS	OSS
Environmental impacts		
Non-renewable energy	0.066	0.060
Land use	0.100	0.041
Eutrophication	0.107	0.141
Acidification	0.090	0.053
GWP	0.060	0.069
Mean	0.085	0.073
Outputs		
Crops	0.003	0.007
Milk	0.033	0.019
Meat	0.040	0.022
Mean	0.025	0.016

CSS = continental specialized systems; OSS = oceanic specialized systems;
 GWP = global warming potential.

managerial inefficiencies (i.e. slacks) from the 59 CSS farms and 126 OSS farms, based on the 'corrective' methodology (Phase 2 in Figure 3). Statistics for the eco-efficiency scores and their mean ranks per system and feeding strategy are presented in Table 5. The mean and mean ranks of eco-efficiency scores were higher for OSS than CSS and the medians of both systems equalled 1. The Kruskal–Wallis test identified significant differences between the eco-efficiency scores' mean ranks of the two systems ($P < 0.001$). The three feeding strategies had almost-equal mean and equal median eco-efficiency scores. The Kruskal–Wallis test did not identify significant differences between feeding strategies in terms of mean ranks of the eco-efficiency scores ($P = 0.767$).

Discussion

This study is aimed at researchers, advisors and policy makers searching for tools that can address the challenges of increasing farm output and reducing EIs, especially given

Table 5 Statistics for eco-efficiency scores per system and feeding strategy after removal of managerial inefficiencies

	Eco-efficiency scores					
	Min	Max	Median	Mean	SD	Mean rank
System						
CSS	0.908	1.000	0.995	0.985	0.022	67.059 ^a
OSS	0.890	1.000	1.000	0.994	0.018	105.147 ^b
Feeding strategy						
<10% maize ¹	0.934	1.000	1.000	0.991	0.017	88.614
10% to 30% maize ¹	0.928	1.000	1.000	0.993	0.016	94.991
>30% maize ¹	0.890	1.000	1.000	0.991	0.024	93.946

CSS = continental specialized systems; OSS = oceanic specialized systems;
 GWP = global warming potential.

^{a,b}Values within a column with different superscripts differ significantly at $P < 0.05$.

¹Maize area as % of total forage area on farm.

the recent trend towards sustainable intensification of agriculture (see Foresight, 2011). Our framework contributes to the stream of literature employing methodologies able to capture several aspects in order to ensure that development is in fact 'sustainable'. Dairy farming was used as an exemplar to demonstrate the framework, which is expandable to other agricultural settings.

Not 'just LCA' but 'DEA and LCA'

According to recent guidelines by the Livestock Environmental Assessment and Performance Partnership (LEAP, 2015, p. 6), '[i]n order to prevent shift of burden from [one] environmental issue to another, no environmental improvement option should be recommended without having [...] assessed [...] the effects on resource use and those other EIs targeted as relevant for livestock supply chains [...]'. In other words, the LEAP guidelines themselves implicitly acknowledge the issue of substitution possibilities between LCA eco-efficiency ratios, mentioned in the introduction to this study. The implications of this issue can be demonstrated by looking at the results of LCA eco-efficiency studies comparing dairy farms with different proportions of land devoted to maize silage (e.g. Basset-Mens *et al.*, 2009; Rotz *et al.*, 2010). According to these studies, because grassland requires less fertilization than arable land, lower impacts from eutrophication, acidification, greenhouse gas emissions and non-renewable energy use have been observed on grass-based farms. However, arable crops such as maize silage have higher yields per hectare. It is therefore impossible to conclude that a particular feeding strategy has a higher eco-efficiency potential than another one, unless all feeding strategies are evaluated at the *aggregate* level, as was done in this study. Indeed, feeding the LCA variables into the RAM model showed that the eco-efficiency of farms with >30% maize was lower, favouring more grass-based systems.

Regional differences

Higher eco-efficiency scores were expected for OSS systems over CSS because the bio-physical conditions in West France are more favourable. Specifically, the climate conditions in West France favour the production of high quality forages which are essential for dairy production. These differences in climate conditions between West and Continental France were implicitly examined in this study by removing managerial inefficiencies from CSS and OSS with the 'corrective' methodology. Indeed, Jan *et al.* (2012) emphasized that DEA results should be interpreted with care as inefficiencies may be attributed to factors that are beyond managerial control. Hence, removing managerial factors with the 'corrective' methodology revealed each system's true eco-efficiency potential that the projected data showed to be possible, solely as a result of the different bio-physical conditions between West and Continental France. OSS systems then ranked significantly higher, on average, than CSS in terms of eco-efficiency scores (Table 5).

Identifying specific sources of eco-'inefficiency'

Examining the slacks (Table 4) can help prioritize the reduction (increase) of those EIs (outputs) most responsible for the eco-inefficiency of CSS and OSS. For example, CSS systems had a quite large acidification slack. In fact, in CSS systems cows are generally offered more protein concentrates, potentially to avoid any protein shortages, which tends to increase ammonia emissions (Faverdin *et al.*, 2014). It is noteworthy that CSS also had a large land use slack (Table 4). These systems devoted a larger part of on-farm land to crop production at the expense of lower milk and meat production than OSS (compare mean crops-milk and crops-meat ratios per system, which can be easily derived from Table 2). This, in turn, explains the lower crops slack, and higher milk and meat slacks, of CSS in comparison with OSS (Table 4). Finally, note that for both systems the largest slack was eutrophication, as opposed to the relatively low global warming potential slacks. This agrees with the findings of Bava *et al.* (2014) that livestock systems are often responsible for important local EIs.

Methodological aspects

Eco-efficiency as a relative measure to improve sustainability. It can be argued that improving eco-efficiency does not guarantee sustainability. Because eco-efficiency is a relative measure, improvements can be achieved if either EIs are reduced or outputs are increased. Furthermore, the absolute environmental pressure can still exceed the ecosystem's carrying capacity (Kuosmanen and Kortelainen, 2005). For example, there is a high concentration of dairy farms in West France and the main production regions are located near environmentally sensitive areas (Chatellier and Pflimlin, 2006). Thus, although OSS systems had higher eco-efficiency, this does not necessarily mean that they operated within the local ecosystem's carrying capacity.

Nevertheless, eco-efficiency is often cost-effective so it makes economic sense to exploit it to the utmost (Kuosmanen and Kortelainen, 2005). In this study the RAM model helped identify such options through the relative EI and output slacks (Table 4). Prioritizing those EIs and outputs with the largest relative slacks can result in notable eco-efficiency improvements. This is advantageous because policies targeted at eco-efficiency improvements tend to be easier to adopt, and politically easier to implement, than policies restricting the level of economic activity (Kuosmanen and Kortelainen, 2005).

Comparing RAM with alternative methods. This study considered RAM's ranking property as one of its main advantages. Besides RAM, there are several promising methods to rank DMUs. See the reviews by Adler *et al.* (2002) and Markovits-Somogyi (2011) regarding the methods mentioned hereafter. Other ranking methods missing from both reviews exist, such as the 'global efficiencies' (GLE) approach by Despotis (2002) which, like RAM, uses a common weighting scheme across all DMUs. These ranking methods can be roughly classified as having at least one of the following characteristics: (i) they require modifications to the original DEA model (e.g. when imposing weights restrictions); (ii) they involve supplementary analyses with tools such as multivariate statistics (e.g. canonical correlation analysis for ranking) or multiple-criteria decision making (e.g. GLE), which translates to additional computational time and/or coding effort; (iii) the original DEA model cannot be easily solved (e.g. fuzzy DEA); and (iv) there is no correspondence between the DMUs' efficiency scores and their ranks (e.g. GLE). While some of these issues can be dealt with fairly easily (e.g. the weights restrictions), to the best of our knowledge, RAM is the only simple, readily available linear DEA model with a ranking property that does not involve (i to iv). Note that RAM can only rank inefficient DMUs. In fact, ranking efficient DMUs was not desirable here because rankings can differ between methods (see Adler *et al.*, 2002), possibly affecting the results of the non-parametric rank statistics.

Additive models (such as RAM) are not the only DEA models able to simultaneously minimize EIs (and/or inputs) and maximize output. Another example is the directional distance function (DDF) whereby the minimization of EIs and inputs, and maximization of outputs, is made via a 'direction vector' that reflects different stakeholder preferences. For example, the direction vector may be set to minimize EIs for the given outputs, maximize outputs for the given EIs or do both simultaneously. Several other choices are also possible (see Beltrán-Estevé *et al.*, 2014; Berre *et al.*, 2014). For instance, Berre *et al.* (2014) argued that a sustainable intensification scenario would seek to reduce pollution and increase outputs with a possible *increase* in inputs. The RAM model can also allow for input increases because it can handle negative values (see Cooper *et al.*, 1999): simply assign a negative sign to the inputs to be increased.

DDFs are advantageous over RAM when the objective is not only to calculate the input and output adjustments necessary for a DMU to operate efficiently, but also to determine how 'far' these adjustments are from an input–output combination maximizing profits (provided that input and output prices are known) for this particular DMU (Färe and Grosskopf, 2000). This 'allocation' problem cannot be modelled with RAM. Nonetheless, RAM is appropriate when it is desirable to decompose efficiency scores into variable-specific scores through the slacks (as was done here) because, unlike DDFs, RAM does not assume proportional adjustments in inputs and outputs (some recently developed DDFs that relax this assumption have in fact an additive structure; see Chen *et al.*, 2015). Note that there are several normalization options for the slacks (other than by division by the variables' ranges as was done here) that create opportunities for further analyses (Cooper *et al.*, 1999 discuss a range of choices). For example, when input prices are known, input slacks can be 'priced' to determine the proportion of each input's cost to the total cost (see Soteriades *et al.*, 2015).

Finally, we draw attention to the alternative definitions of 'data-driven' weights in models 1 and 2. In model 1 the weights are calculated by the model itself. This may result in large weights for EIs of secondary importance, leaving a negligible or zero weight for more important EIs (Kuosmanen and Kortelainen, 2005). This can be fixed by restricting *a priori* the weights' values to admissible ranges (see Kuosmanen and Kortelainen, 2005). By contrast, with RAM (model 2) the weights are not *calculated* but *given*, because the model uses the variable's ranges as weights, which are always non-zero. Therefore, reliance on subjective weights restrictions as in model 1 is not necessary with RAM.

Choice of DEA variables. Choice of input and output variables used is a key aspect of DEA methodology. Past studies on dairy farm eco-efficiency with DEA often use one aggregate output indicator to avoid too many DMUs on the efficient frontier. For example, Pérez Urdiales *et al.* (2015) defined output as economic value added [(milk sales + value of on-farm consumption of milk) – direct costs]. On the other hand, Jan *et al.* (2012) argued that economic value added might bias the results as an increase in the market price of a given commodity would lead to higher eco-efficiency. Instead, they aggregated all farm outputs into a single output of digestible energy content. However, with this method it is assumed that any form of energy in human diets can be substituted by any other, provided that energy requirements are met. Also, milk, meat and crops have different nutritional values in addition to energy content. Therefore, in this study it was deemed more appropriate to keep milk, meat and crops as three separate outputs.

Furthermore, in this study the eco-efficiency measure did not include operational inputs (e.g. labour, capital, on-farm electricity use) and 'undesirable' outputs (e.g. kg CO₂-equivalents, wastewater) because the idea was to aggregate *altogether* the two elements used in

LCA ratios: EIs and outputs. In other words, we were concerned with the EIs rather than the amount of operational inputs and undesirable outputs of DMUs (see Jan *et al.*, 2012, p. 715, but also Kuosmanen and Kortelainen, 2005). An alternative way of conducting eco-efficiency analysis by also involving operational inputs and undesirable outputs is with the 'LCA + DEA method' (see Vázquez-Rowe and Iribarren, 2015). With LCA + DEA, 'target' LCA impacts are obtained by adjusting the operational inputs to their optimal values via DEA and re-performing the LCA exercise. Therefore, in LCA + DEA the DEA exercise is an intermediate step that helps determine the DMUs' benchmarks and thus the target EIs. Alternatively, target EIs can be obtained directly from RAM's optimal slacks. This reduces potential dimensionality issues because the set of DEA variables will generally be smaller than that with LCA + DEA (Jan *et al.*, 2012, p. 715).

Conclusion

Combining LCA with RAM, the 'corrective' methodology and non-parametric rank tests can significantly improve (dairy) farm eco-efficiency assessments compared to previous studies using partial ratios or coupling LCA with DEA. The modelling framework was demonstrated with LCA data for French specialized dairy farms. Results showed that OSS systems ranked higher, on average, than CSS systems in terms of eco-efficiency. Also, the average eco-efficiency rank of farms with lower proportions of maize silage in the total forage area was higher, on average, than farms with higher proportions of maize. These results helped identify the eco-efficiency potential of each region and feeding strategy and could therefore aid advisors and policy makers at farm or region/sector level. This demonstration also highlights the capacity of the proposed multiple-EI, multiple-output framework to measure and understand eco-efficiency, and to compare different groups, which makes it a promising multiple-criteria tool towards the achievement of greater yet sustainable agricultural production.

Acknowledgements

The authors would like to express their gratitude to two anonymous referees whose constructive comments helped significantly improve an earlier version of this manuscript. The authors take responsibility for any errors that may remain. We acknowledge feedback from staff at SRUC (Scotland), INRA (St Gilles, France) and Idele (Le Rheu, France) and also from Dr Philippe Lecomte (CIRAD, France) and Dr Marc Metzger (University of Edinburgh). Likewise, thank you to Professor Mette Asmild (University of Copenhagen) and Professors Jean-Philippe Boussemart and Hervé Leleu at IÉSEG (Lille, France) for the fruitful discussions on DEA, as well as to Dr Jiayi Liu and Dr Giles Innocent (BioSS, Scotland) for their tips in statistics. This project is jointly funded by SRUC (via Scottish Government's Rural Affairs and the Environment Strategic Research programme) and the PHASE division of INRA.

References

- Adler N, Friedman L and Sinuany-Stern Z 2002. Review of ranking methods in the data envelopment analysis context. *European Journal of Operational Research* 140, 249–265.
- Basset-Mens C, Ledgard S and Boyes M 2009. Eco-efficiency of intensification scenarios for milk production in New Zealand. *Ecological Economics* 68, 1615–1625.
- Bava L, Sandrucci A, Zucali M, Guerci M and Tamburini A 2014. How can farming intensification affect the environmental impact of milk production? *Journal of Dairy Science* 97, 4579–4593.
- Béguin E, Bonnet J, Dollé J-B, Charroin T and Ferrand M 2008. Les différents postes de consommation en énergie et les pistes d'économie en élevage bovin laitier. *Rencontres Recherches Ruminants* 15, 217–220.
- Beltrán-Estevé M, Picazo-Tadeo A and Reig-Martínez E 2014. A metafrontier directional distance function approach to assessing eco-efficiency. *Journal of Productivity Analysis* 41, 69–83.
- Berre D, Blancard S, Boussemart JP, Leleu H and Tillard E 2014. Finding the right compromise between productivity and environmental efficiency on high input tropical dairy farms: a case study. *Journal of Environmental Management* 146, 235–244.
- Bogetoft P and Otto L 2011. *Benchmarking with DEA, SFA, and R*. Springer, New York, NY, USA.
- Brockett PL, Cooper WW, Golden LL, Rousseau JJ and Wang YY 2004. Evaluating solvency versus efficiency performance and different forms of organization and marketing in US property – liability insurance companies. *European Journal of Operational Research* 154, 492–514.
- Brockett PL and Golany B 1996. Using rank statistics for determining programmatic efficiency differences in data envelopment analysis. *Management Science* 42, 466–472.
- Casey JW and Holden NM 2005. The relationship between greenhouse gas emissions and the intensity of milk production in Ireland. *Journal of Environmental Quality* 34, 429–436.
- Charroin T, Palazon R, Madeline Y, Guillaumin A and Tchakerian E 2005. Le système d'information des Réseaux d'Élevage français sur l'approche globale de l'exploitation. Intérêt et enjeux dans une perspective de prise en compte de la durabilité. *Rencontres Recherches Ruminants* 12, 335–338.
- Chatellier V and Pflimlin A 2006. Dairy systems in the European regions of the Atlantic area. In *Proceedings of the Final Seminar of the Green Dairy Project*, 13 to 14 December 2006, Rennes, France pp. 11–30.
- Chen PC, Yu MM, Chang CC, Hsu SH and Managi S 2015. The enhanced Russell-based directional distance measure with undesirable outputs: numerical example considering CO₂ emissions. *Omega* 53, 30–40.
- Conover WJ 1999. *Practical nonparametric statistics*, 3rd edition. John Wiley & Sons Inc, New York, NY, USA.
- Cooper WW, Park KS and Pastor JT 1999. RAM: a range adjusted measure of inefficiency for use with additive models, and relations to other models and measures in DEA. *Journal of Productivity Analysis* 11, 5–42.
- Cooper WW, Seiford LM and Tone K 2007. *Data envelopment analysis: a comprehensive text with models, applications, references and DEA-Solver software*, 2nd edition. Springer Science + Business Media, LLC, New York, NY, USA.
- CORPEN- Comité d'orientation pour de pratiques agricoles respectueuses de l'environnement 1999. Estimation des flux d'azote, de phosphore et de potassium associés aux vaches laitières et à leur système fourrager : influence de l'alimentation et du niveau de production. Ministère de l'Écologie, de l'Énergie, du Développement durable et de la Mer, Ministère de l'Agriculture, France.
- Despotis DK 2002. Improving the discriminating power of DEA: focus on globally efficient units. *Journal of the Operational Research Society* 53, 314–323.
- EMEP/CORINAIR- European Monitoring and Evaluation Programme/Core Inventory of Air Emissions in Europe 2002. *Atmospheric emission inventory guidebook*, 3rd edition. European Environment Agency, Copenhagen, Denmark.
- ESCAP- Economic and Social Commission for Asia and the Pacific 2009. *Eco-efficiency indicators: measuring resource-use efficiency and the impact of economic activities on the environment*. ESCAP, Bangkok, Thailand.
- Färe R and Grosskopf S 2000. Theory and applications of directional distance functions. *Journal of Productivity Analysis* 13, 93–103.
- Faverdin P, Baratte C, Perbost R, Thomas S, Ramat E and Peyraud J-L 2014. A web tool to simply assess feed autonomy and nitrogen excretion of dairy herds: CowNex. *Rencontres Recherches Ruminants* 21, 391–394.
- Foresight 2011. *The future of food and farming*. The Government Office for Science, London, UK.
- Gac A, Deltour L, Cariolle M, Dollé J-B, Espagnol S, Flénet F, Guingand N, Lagadec S, Le Gall A, Lellahi A, Malaval C, Ponchant P and Tailleur A 2010a. GES'TIM, Guide méthodologique pour l'estimation des impacts des activités agricoles sur l'effet de serre. Version 1.2. Institut de l'Élevage, Paris, France.
- Gac A, Manneville V, Raison C, Charroin T and Ferrand M 2010b. L'empreinte carbone des élevages d'herbivores : présentation de la méthodologie d'évaluation appliquée à des élevages spécialisés lait et viande. *Rencontres Recherches Ruminants* 17, 335–342.
- Guerci M, Bava L, Zucali M, Sandrucci A, Penati C and Tamburini A 2013. Effect of farming strategies on environmental impact of intensive dairy farms in Italy. *Journal of Dairy Research* 80, 300–308.
- Guinée JB, Gorrié M, Heijungs R, Huppes G, Kleijn R, Koning de A, Oers van L, Wegener Sleswijk A, Suh S, Udo de Haes HA, Bruijn de H, Duin van R and Huijbregts MAJ 2002. *Handbook on life cycle assessment. Operational guide to the ISO standards*. I: LCA in perspective. IIa: Guide. IIb: Operational annex. III: Scientific background. Kluwer Academic Publishers, Dordrecht, the Netherlands.
- IPCC- Intergovernmental Panel on Climate Change 2007. *Climate Change 2007: the physical science basis. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change*. Cambridge University Press, Cambridge, UK.
- Jan P, Dux D, Lips M, Alig M and Dumondel M 2012. On the link between economic and environmental performance of Swiss dairy farms of the alpine area. *International Journal of Life Cycle Assessment* 17, 706–719.
- Kuosmanen T and Kortelainen M 2005. Measuring eco-efficiency of production with data envelopment analysis. *Journal of Industrial Ecology* 9, 59–72.
- LEAP- Livestock Environmental Assessment and Performance Partnership 2015. *Developing sound tools for transition to sustainable food and agriculture. LEAP partnership life cycle assessment guidelines on livestock supply chains: methodological notes*. Draft version 1 for public review. LEAP, Rome, Italy.
- Markovits-Somogyi R 2011. Ranking efficient and inefficient decision making units in data envelopment analysis. *International Journal for Traffic and Transport Engineering* 1, 245–256.
- Nemecek T and Kägi T 2007. *Life cycle inventories of Swiss and European agricultural production systems. Final report ecoinvent V2.0 No. 15a*. Agroscope Reckenholz-Taenikon Research Station ART, Zurich, Switzerland.
- Pérez Urdiales M, Oude Lansink A and Wall A 2015. Eco-efficiency among dairy farmers: the importance of socio-economic characteristics and farmer attitudes. *Environmental and Resource Economics*, doi:10.1007/s10640-015-9885-1, Published online by Springer Netherlands 20 February 2015.
- Ramli NA and Munisamy S 2015. Eco-efficiency in greenhouse emissions among manufacturing industries: a range adjusted measure. *Economic Modelling* 47, 219–227.
- Rotz CA, Montes F and Chianese DS 2010. The carbon footprint of dairy production systems through partial life cycle assessment. *Journal of Dairy Science* 93, 1266–1282.
- Sheskin DJ 1997. *Handbook of parametric and nonparametric statistical procedures*. CRC Press LLC, Boca Raton, FL, USA.
- Soteriades AD, Faverdin P, March M and Stott AW 2015. Improving efficiency assessments using additive data envelopment analysis models: an application to contrasting dairy farming systems. *Agricultural and Food Science* 24, 235–248.
- van Calster KJ, Berentsen PBM, Giesen GWJ and Huirne RBM 2008. Maximising sustainability of Dutch dairy farming systems for different stakeholders: a modelling approach. *Ecological Economics* 65, 407–419.
- Vázquez-Rowe I and Iribarren D 2015. Review of life-cycle approaches coupled with data envelopment analysis: launching the CFP. *The Scientific World Journal*. doi.org/10.1155/2015/813921.