

Hindawi Publishing Corporation
Autism Research and Treatment
Volume 2012, Article ID 724072, 5 pages
doi:10.1155/2012/724072

Research Article

Identification of Four Novel Synonymous Substitutions in the X-Linked Genes *Neurologigin 3* and *Neurologigin 4X* in Japanese Patients with Autistic Spectrum Disorder

Kumiko Yanagi,¹ Tadashi Kaname,¹ Keiko Wakui,² Ohiko Hashimoto,³ Yoshimitsu Fukushima,² and Kenji Naritomi¹

¹ Department of Medical Genetics, Graduate School of Medicine, University of The Ryukyus, 207 Uehara, Nishihara, Okinawa 903-0215, Japan

² Department of Medical Genetics, School of Medicine, Shinshu University, 3-1-1 Asahi, Matsumoto 390-8621, Japan

³ Faculty of Nursing and Rehabilitation, Aino University, 4-5-4 Higashi-Ohta, Ibaraki 567-0012, Japan

Correspondence should be addressed to Tadashi Kaname, tkaname@med.u-ryukyu.ac.jp

Received 22 January 2012; Revised 6 April 2012; Accepted 23 May 2012

Academic Editor: Bennett L. Leventhal

Copyright © 2012 Kumiko Yanagi et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Mutations in the X-linked genes *neurologigin 3* (*NLGN3*) and *neurologigin 4X* (*NLGN4X*) were first implicated in the pathogenesis of X-linked autism in Swedish families. However, reports of mutations in these genes in autism spectrum disorder (ASD) patients from various ethnic backgrounds present conflicting results regarding the etiology of ASD, possibly because of genetic heterogeneity and/or differences in their ethnic background. Additional mutation screening study on another ethnic background could help to clarify the relevance of the genes to ASD. We scanned the entire coding regions of *NLGN3* and *NLGN4X* in 62 Japanese patients with ASD by polymerase chain reaction-high-resolution melting curve and direct sequencing analyses. Four synonymous substitutions, one in *NLGN3* and three in *NLGN4X*, were identified in four of the 62 patients. These substitutions were not present in 278 control X-chromosomes from unrelated Japanese individuals and were not registered in the database of Single Nucleotide Polymorphisms build 132 or in the Japanese Single Nucleotide Polymorphisms database, indicating that they were novel and specific to ASD. Though further analysis is necessary to determine the physiological and clinical importance of such substitutions, the possibility of the relevance of both synonymous and nonsynonymous substitutions with the etiology of ASD should be considered.

1. Introduction

Neurologigin 3 (*NLGN3*) and *Neurologigin 4X* (*NLGN4X*) are members of neuroligins expressed in the postsynaptic neurons and mediate transsynaptic signaling by interacting their ligand, neuroligins [1]. Mutations in the X-linked genes *NLGN3* and *NLGN4X* (GenBank accession numbers NM_181303.1 and NM_020742.2, resp.) were first reported as being involved in X-linked autistic spectrum disorder (ASD; MIM#300425 and MIM#300495) in Swedish families [2]. Some reports have also indicated that *NLGN3* and *NLGN4X* are responsible for ASD. Mutations in these genes including 5 missense mutations [2–5], two small in-del mutations, which lead to a premature stop codon in the transcript [2, 6], an exon skipping mutation [7], and a large deletion

[8], have been identified in ASD patients (Table 1). In vitro experiments using *NLGN3* and *NLGN4X* proteins carrying amino acid changes that were identified in ASD patients indicated that the gene mutations could cause ASD by a loss-of-function mechanism [5, 9]. *Neurologigin 3* p.R451C knock-in mice and *neurologigin 4X*-deficient mice exhibited autism-related behaviors [10, 11]. In addition, recent mutation screening studies revealed that the p.K378R substitution in the *NLGN4X* gene was common in individuals of different ethnicities, such as Portuguese and Greek [3, 4]. A recent association study regarding rare variants in *NLGN3* and *NLGN4X* also supported the relationship between a specific *NLGN3* haplotype and the etiology of ASD in Chinese Han male population [12]. Moreover, two synonymous variants are specifically found in ASD patients [13, 14].

TABLE 1: Previously identified sequence variations in coding regions in ASD patients.

Gene	NT ¹ change	A-A ² change	Mutation type	Ethnic background	Ref. ³	NCBI
<i>NLGN3</i>	c.222C>T	p.Y74Y	Synonymous	Finnish	[13]	NM_181303.1
	c.1351C>T	p.R451C	Missense	Swedish	[2]	NM_018977.3
	c.259C>T	p.R87W	Missense	Irish and Scottish	[5]	NM_020742.2
	c.759G>A	p.G99S	Missense	Portuguese	[3]	NM_020742.2
	c.1186 insT	p.D396X	Frameshift	Swedish	[2]	NM_020742.2
<i>NLGN4X</i>	c.1253delAG	p.D429X	Frameshift	French	[6]	NM_020742.2
	c.1597A>G	p.K378R	Missense	Portuguese and Greek	[3, 4]	NM_020742.2
	Not described ⁴	p.A558A	Synonymous	German	[14]	NM_020742.2
	c.2574C>T	p.R704C	Missense	Portuguese	[3]	NM_020742.2
	del exon 4	In-frame ⁵	Skipping	AGRE ⁶	[7]	NM_020742.2
	del exon 4–6	Truncated ⁷	Large deletion	Irish and English	[8]	NM_020742.2

¹NT: nucleotide.

²A-A: amino acid.

³Ref.: reference number.

⁴The number of substituted nucleotides was not mentioned in the reference.

⁵Exon 4 skipping mutation was predicted to result in an in-frame exclusion of 62 amino acids.

⁶Autism genetics resource exchange.

⁷The translated protein was predicted to be entirely truncated between exon 3 and exon 6.

In contrast, some data indicate that these two genes do not have a significant effect on ASD development. The frequency of mutation in the coding region was not high (<2%) among ASD patients [2–6]. It has been reported that mutations in these genes were not observed in ASD patients in the United Kingdom, Canada, the Autism Genetic Resource Exchange (AGRE) and the International Molecular Genetic Study of Autism Consortium (IMGSAC) [15–17].

Such contradictory results could possibly be due to genetic heterogeneity, regional characteristics, or differences in the ethnic background of ASD patients. It is, therefore, important to investigate the *NLGN3* and *NLGN4X* sequences in ASD patients from various ethnic backgrounds.

Molecular screening of *NLGN3* and *NLGN4X* in Japanese patients with ASD has not yet been reported. We, therefore, performed a mutation screening study of these genes in 62 unrelated Japanese patients with ASD and 278 control X-chromosomes using polymerase-chain-reaction (PCR) high-resolution melting (HRM) and direct sequencing analyses.

2. Materials and Methods

2.1. Patients. The Japanese ASD patients analyzed in this study were described previously [18]. All the patients were diagnosed with ASD according to the Diagnostic and Statistical Manual of Mental Disorders (Fourth Edition criteria), the Autism Diagnostic Interview-Revised, and the Childhood Autism Rating Scale by at least two trained psychiatrists. No chromosomal aberration was found in all patients. The study was approved by the ethics committee of the University of the Ryukyus, and written informed consent was obtained from all subjects.

2.2. Source of Genomic DNA. Genomic DNA in patients (51 males, 11 females) and control Japanese individuals

(30 males, 30 females) was isolated from successfully transformed B cells by Epstein-Barr virus using a standard protocol involving proteinase K digestion. In addition, control genomic DNA was isolated from blood of 198 healthy Japanese individuals (118 males, 80 females) using a QIA amp column (Qiagen, Hiden, Germany).

2.3. Mutation Screening. Initially, mutation screening was performed in the 62 ASD subjects by PCR-HRM analysis. HRM analysis is used to scan gene variations in a PCR amplicon, for example, mutations and single-nucleotide polymorphisms (SNPs), prior to sequencing by detecting differences in the thermal denaturation of double-stranded DNA [19, 20]. We set up optimal conditions for PCR-HRM analysis of the entire coding regions of *NLGN3* and *NLGN4X*. Primers for all exons were carefully designed in flanking introns of the genes taking into account the high sequence homology between neuroligins (Table 2). Exons 2, 7, and 8 of *NLGN3* and exons 2, 5, and 6 of *NLGN4X* were amplified in two or three segments to obtain a fine resolution of the HRM curves. Only an upstream part of exon 2 (exon 2.1 in Table 2) of *NLGN4X* was sequenced in all ASD patients because we failed to distinguish between the complicated SNPs located in the region.

2.4. Condition of PCR-HRM Analysis. PCR was performed under optimized conditions using Ex-Taq DNA polymerase (TAKARA Bio Inc., Japan) as follows: 20 μ L reaction mixture containing Mg²⁺-free 1 \times PCR buffer (TAKARA Bio Inc., Japan), 0.25 mM dNTPs (TAKARA Bio Inc., Japan), 0.25 units Ex-Taq DNA polymerase, 1 μ M of each primer, and the optimal concentration of MgCl₂ was loaded per well in a 96-well PCR plate (Roche, Basel, Switzerland). The amount of genomic DNA used in each reaction was 40 ng for males and 20 ng for females. Fluorescent DNA

TABLE 2: Primer sets used to screen for variants by PCR-HRM analysis.

Gene	Exon	Forward	Reverse
<i>NLGN3</i>	2.1	GCTCAGTTTTGAGGTTCAAGTC	TCACTGGGCAGTGGTACTCG
	2.2	CACAGTCAACACTCACTTTGG	GATGGTTAGAAGCATTTTCACAG
	3	GGCAGAGGCCTCCTGTTATT	CAATCCTCCCTGCAAGGCA
	4	TGGCTTGCTGGGCCACACTG	GCCAAAGACAGATGAACAGCC
	5	AGGTTGAGCAACCCCATGAGT	GGGCCAGAGGATAACACCATT
	6	CATCCCTCTGCCTTCATTGTC	TAGAAGAGAGCTGGCCGATTG
	7.1	CAGCCTCAGTGACAAAGGAAT	CAGGGTGTCTTACCCTCAG
	7.2	GTAAGGACACCCTGCGAGAG	TGGGTCTCAAAGAGGAAAA
	8.1	GTGACCCAGATTTCATGT	GGCCAGAACGTTAAGGAACA
	8.2	ATCACCCGAGGCCCAATGG	CCTCACACTCGTGGTGGGTG
8.3	GGAGGAGCTGGCAGCATTAC	CTGGAGATTGGCTGTGCTCT	
<i>NLGN4X</i>	2.1	AAAGCCCTATCTCTGCGAGG	TGAGTAGTATTTCCGGATGCCAG
	2.2	AAGAACACCGTTACCCAATGAG	GAGACATTATAAACCCCTCCTAG
	3	TTAGCATTGGTGAGTCAGTGTG	CCGTCAAACGAGAAGTGGACT
	4	CTTTTTCTATTTGGCCACCA	TTCTTGTTTCAGGGTATTTGC
	5.1	AGCTGCATTTCTGTCCTGTG	TCTCCCGCAAAGTGTCTTTC
	5.2	CCAACCTTCGTGGACAACCTT	ACCCCAACACGAAGATGAAC
	6.1	CACGTACATGTGGAAGAGT	GACGGCAATGGTGACACTTA
	6.2	TCCTCATTGAAACCAAACGA	AACATTCTGGTCTGGAGAC

dye, SYTO 9 (0.5 μ M; Life Technologies, Carlsbad, CA), was added before amplification. Genotype analysis of each sample was performed in duplicate. The melting curves were sequentially analyzed using LightCycler 480 Gene Scanning Software (Roche, Basel, Switzerland) to detect sequence variants. Evaluation of the HRM curves was sequentially confirmed by direct sequencing (Life Technologies, Carlsbad, CA).

2.5. Screening in Controls. Variants identified in ASD patients were then investigated for detecting the common variants by searching the database of Single Nucleotide Polymorphisms (dbSNP) build 132 and the Japanese Single Nucleotide Polymorphism database (JSNP). All variants found in ASD patients were scanned in genomic DNAs isolated from EBV transformed cells or blood cells in control Japanese individuals.

2.6. Assessment of the Substitutions. The effect of the substitutions was evaluated using Mutation Taster software (<http://www.mutationtaster.org/>).

3. Results

We identified four variants, a synonymous substitution and three intronic substitutions, in *NLGN3* in the 62 ASD patients. The synonymous substitution (c.1698G>A, p.K566K) was observed in a male ASD patient. We also identified four variants, comprising three synonymous substitutions and a substitution in the 5' UTR region of *NLGN4X*. One synonymous substitution, c.297C>T (p.G99G), was observed in a female patient, and two substitutions, c.516C>T (p.I172I) and c.1590C>T (p.F530F), were

observed in two different male patients. The variants identified in this study are summarized in Table 3. Nonsynonymous substitutions, including seven substitutions reported previously, were not identified in the Japanese patients by PCR-HRM analysis (Tables 1 and 3). The four synonymous and an intronic substitution were not found in the control X-chromosomes from unrelated healthy Japanese individuals, without any neuropsychiatric disorder (Table 3).

4. Discussion

By using PCR-HRM analysis we identified an intronic and four exonic substitutions only in 62 Japanese patients with ASD, two out of 62 patients (3.2%) in *NLGN3* and three out of 62 patients (4.8%) in *NLGN4X*. The exonic substitutions comprised one synonymous substitution in *NLGN3* and three synonymous substitutions in *NLGN4X* (Table 3). The PCR-HRM analysis could detect 90% of the sequence variations with 100% accuracy [19, 20]; therefore, we were able to identify almost all the changes in *NLGN3* and *NLGN4X* in these patients.

In this study, we analyzed genomic DNAs from EBV-transformed cells in patients with ASD. The source of genomic DNA, especially extracted from EBV-transformed cells, should be considerable in each experiment, because there is a possibility that unexpected substitutions occur during the transformation [21, 22]. Analyses in control genomic DNA showed that there was no sequence alteration or substitution bias in the exons of *NLGN3* and *NLGN4X* between genomic DNA isolated from blood and EBV-transformed cells in this study (Table 3).

Our mutation screening study in Japanese ASD patients failed to detect novel nonsynonymous mutations and seven

TABLE 3: Sequence variants identified in Japanese patients with autistic spectrum disorder.

Gene	Exon	SNP ID ¹	Location	Translation	ASD ²		Control ³	
					Male	Female	EBV cell line	Blood
NLGN3	4	—	c.567+22C>T	Intronic	0/51	1/11	0/90	3/278
		—	c.567+52C>T	Intronic	10/51	2/11	12/90	32/278
	5	—	c.727+47G>C	Intronic	1/51	0/11	0/90	0/278
	7	—	c.1698G>A	p.K566K	1/51	0/11	0/90	0/278
NLGN4X		rs6639602	c.-305-86T>G	5'UTR	0/51	1/11	1/90	4/278
	2	—	c.297C>T	p.G99G	0/51	1/11	0/90	0/278
	3	—	c.516C>T	p.I172I	1/51	0/11	0/90	0/278
	5	—	c.1590C>T	p.F530F	1/51	0/11	0/90	0/278

¹Reference number of the variant documented in dbSNP build 132 or JSNP. (—) indicates that the variant does not have a reference number.

²Number of ASD patients (50 males, 11 females) with the variant.

³Number of control chromosomes with the variant in EBV transformed cell line (30 males, 30 females) and blood (118 males, 80 females).

known nonsynonymous mutations that were identified in previous studies [2–5] as well as a study in a Chinese Han population [12]. Considering the low frequency of nonsynonymous substitutions in these genes seen in previous reports [15–17], a larger number of ASD patients should be sampled. Nevertheless, our results suggest that nonsynonymous substitutions in *NLGN3* and *NLGN4X* may account for only a small proportion of Japanese patients with ASD. In addition to ASD, there are some reports indicating that *NLGN3* and *NLGN4X* are also relevant to Asperger syndrome [2], X-linked mental retardation [6], and Tourette syndrome [8]. Considering the function of *NLGN3* and *NLGN4X* [1], additional mutation screening studies in such neurobehavioral disorders should be needed in near future.

Experimental evidence is increasing that synonymous substitutions could affect the protein function through transcription or translation impairment [23–25]. While the clinical and physiological importance of the four synonymous substitutions and an intronic substitution are not clear at this moment, Mutation Taster software (<http://www.mutationtaster.org/>) [26] evaluated that the substitutions might affect protein structure by altering splice site. According to previous reports of mutation screening in *NLGN3* and *NLGN4X*, two synonymous mutations, p.Y74Y in *NLGN3* and p.A558A in *NLGN4X* (Table 1), were also observed in ASD patients but not in healthy controls [13, 14]. A recent study in a Chinese Han population has indicated that a common intronic variation in *NLGN3* may influence the susceptibility of males to ASD [12]. Although further analysis is necessary to demonstrate the biological effects of synonymous and intronic substitutions on ASD, these substitutions as well as nonsynonymous substitutions should be taken into account.

5. Conclusion

We identified four substitutions, one in *NLGN3* and three in *NLGN4X*, specific to Japanese patients with ASD. They were synonymous but the possibility of the association of both synonymous and nonsynonymous substitutions with the etiology of ASD should be considered.

Conflict of Interests

The authors declare that they have no competing interests.

Acknowledgments

The authors thank all the individuals who participated in this study. They also thank Konomi Morita and Akihiro Hanafusa for expert technical assistance. This work was financially supported by a Grant-in-Aid for Scientific Research(C) from the Japan Society for the Promotion of Science (project 21590362).

References

- [1] T. C. Südhof, “Neuroligins and neuroligins link synaptic function to cognitive disease,” *Nature*, vol. 455, no. 7215, pp. 903–911, 2008.
- [2] S. Jamain, H. Quach, C. Betancur et al., “Mutations of the X-linked genes encoding neuroligins *NLGN3* and *NLGN4* are associated with autism,” *Nature Genetics*, vol. 34, no. 1, pp. 27–29, 2003.
- [3] J. Yan, G. Oliveira, A. Coutinho et al., “Analysis of the neuroigin 3 and 4 genes in autism and other neuropsychiatric patients,” *Molecular Psychiatry*, vol. 10, no. 4, pp. 329–332, 2005.
- [4] A. Pampanos, K. Volaki, E. Kanavakis et al., “A substitution involving the *NLGN4* gene associated with autistic behavior in the Greek population,” *Genetic testing and molecular biomarkers*, vol. 13, no. 5, pp. 611–615, 2009.
- [5] C. Zhang, J. M. Milunsky, S. Newton et al., “A neuroigin-4 missense mutation associated with autism impairs neuroigin-4 folding and endoplasmic reticulum export,” *Journal of Neuroscience*, vol. 29, no. 35, pp. 10843–10854, 2009.
- [6] F. Laumonier, F. Bonnet-Brihault, M. Gomot et al., “X-Linked mental retardation and autism are associated with a mutation in the *NLGN4* gene, a member of the neuroigin family,” *American Journal of Human Genetics*, vol. 74, no. 3, pp. 552–557, 2004.
- [7] Z. Talebizadeh, D. Y. Lam, M. F. Theodoro, D. C. Bittel, G. H. Lushington, and M. G. Butler, “Novel splice isoforms for *NLGN3* and *NLGN4* with possible implications in autism,” *Journal of medical genetics*, vol. 43, no. 5, p. e21, 2006.

- [8] A. Lawson-Yuen, J. S. Saldivar, S. Sommer, and J. Picker, "Familial deletion within *NLGN4* associated with autism and Tourette syndrome," *European Journal of Human Genetics*, vol. 16, no. 5, pp. 614–618, 2008.
- [9] B. Cih, S. K. Afridi, L. Clark, and P. Scheiffele, "Disorder-associated mutations lead to functional inactivation of neuroligins," *Human Molecular Genetics*, vol. 13, no. 14, pp. 1471–1477, 2004.
- [10] K. Tabuchi, J. Blundell, M. R. Etherton et al., "A neuroligin-3 mutation implicated in autism increases inhibitory synaptic transmission in mice," *Science*, vol. 318, no. 5847, pp. 71–76, 2007.
- [11] S. Jamain, K. Radyushkin, K. Hammerschmidt et al., "Reduced social interaction and ultrasonic communication in a mouse model of monogenic heritable autism," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 105, no. 5, pp. 1710–1715, 2008.
- [12] J. Yu, X. He, D. Yao, Z. Li, H. Li, and Z. Zhao, "A sex-specific association of common variants of neuroligin genes (*NLGN3* and *NLGN4X*) with autism spectrum disorders in a Chinese Han cohort," *Behavioral and Brain Functions*, vol. 7, p. 13, 2011.
- [13] T. Ylisaukko-oja, K. Rehnström, M. Auranen et al., "Analysis of four neuroligin genes as candidates for autism," *European Journal of Human Genetics*, vol. 13, no. 12, pp. 1285–1292, 2005.
- [14] A. K. Wermter, I. Kamp-Becker, K. Strauch, G. Schulte-Körne, and H. Remschmidt, "No evidence for involvement of genetic variants in the x-linked neuroligin genes *NLGN3* and *NLGN4X* in probands with autism spectrum disorder on high functioning level," *American Journal of Medical Genetics B*, vol. 147, no. 4, pp. 535–537, 2008.
- [15] J. B. Vincent, D. Kolozsvari, W. S. Roberts, P. F. Bolton, H. M. D. Gurling, and S. W. Scherer, "Mutation screening of X-chromosomal neuroligin genes: No mutations in 196 autism probands," *American Journal of Medical Genetics*, vol. 129, no. 1, pp. 82–84, 2004.
- [16] J. Gauthier, A. Bonnel, J. St-Onge et al., "*NLGN3/NLGN4* gene mutations are not responsible for autism in the Quebec population," *American Journal of Medical Genetics*, vol. 132, no. 1, pp. 74–75, 2005.
- [17] F. Blasi, E. Bacchelli, G. Pesaresi, S. Carone, A. J. Bailey, and E. Maestrini, "Absence of coding mutations in the X-linked genes neuroligin 3 and neuroligin 4 in individuals with autism from the IMGSAC collection," *American Journal of Medical Genetics*, vol. 141, no. 3, pp. 220–221, 2006.
- [18] T. Marui, O. Hashimoto, E. Nanba et al., "Association between the neurofibromatosis-1 (NF1) locus and autism in the Japanese population," *American Journal of Medical Genetics*, vol. 131, no. 1, pp. 43–47, 2004.
- [19] K. M. Ririe, R. P. Rasmussen, and C. T. Wittwer, "Product differentiation by analysis of DNA melting curves during the polymerase chain reaction," *Analytical Biochemistry*, vol. 245, no. 2, pp. 154–160, 1997.
- [20] A. D. Laurie, M. P. Smith, and P. M. George, "Detection of factor VIII gene mutations by high-resolution melting analysis," *Clinical Chemistry*, vol. 53, no. 12, pp. 2211–2214, 2007.
- [21] P. Awadalla, J. Gauthier, R. A. Myers et al., "Direct measure of the de novo mutation rate in autism and schizophrenia cohorts," *American Journal of Human Genetics*, vol. 87, no. 3, pp. 316–324, 2010.
- [22] L. Sie, S. Loong, and E. K. Tan, "Utility of lymphoblastoid cell lines," *Journal of Neuroscience Research*, vol. 87, no. 9, pp. 1953–1959, 2009.
- [23] C. Kimchi-Sarfaty, J. M. Oh, I. W. Kim et al., "A "silent" polymorphism in the *MDR1* gene changes substrate specificity," *Science*, vol. 315, no. 5811, pp. 525–528, 2007.
- [24] Z. E. Sauna, C. Kimchi-Sarfaty, S. V. Ambudkar, and M. M. Gottesman, "The sounds of silence: synonymous mutations affect functions," *Pharmacogenomics*, vol. 8, no. 6, pp. 527–532, 2007.
- [25] K. E. Knobe, E. Sjörin, and R. C. R. Ljung, "Why does the mutation G17736A/Val107Val (silent) in the *F9* gene cause mild haemophilia B in five Swedish families?" *Haemophilia*, vol. 14, no. 4, pp. 723–728, 2008.
- [26] J. M. Schwarz, C. Rödelberger, M. Schuelke, and D. Seelow, "MutationTaster evaluates disease-causing potential of sequence alterations," *Nature Methods*, vol. 7, no. 8, pp. 575–576, 2010.



Hindawi

Submit your manuscripts at <http://www.hindawi.com>

