

Short report

Ethnic differences in the adaptation rate of HIV gp120 from a vaccine trial

Marcos Pérez-Losada*¹, David Posada², Miguel Arenas², David V Jobes³, Faruk Sinangil⁴, Phillip W Berman⁵ and Keith A Crandall⁶

Address: ¹CIBIO, Centro de Investigação em Biodiversidade e Recursos Genéticos, Universidade do Porto, Campus Agrário de Vairão, 4485-661 Vairão, Portugal, ²Departamento de Bioquímica, Genética e Inmunología. Universidad de Vigo, 36310 Vigo, Spain, ³Presidio Pharmaceuticals, Inc., San Francisco, CA 94158, USA, ⁴Global Solutions for Infectious Diseases, South San Francisco, CA 94080, USA, ⁵Department of Biomolecular Engineering, University of California, Santa Cruz, CA 95064, USA and ⁶Department of Biology, Brigham Young University, Provo, UT 84602, USA

Email: Marcos Pérez-Losada* - mlosada@genoma-llc.com; David Posada - dposada@genoma-llc.com; Miguel Arenas - miguelab@uvigo.es; David V Jobes - djobes@gmail.com; Faruk Sinangil - fsinagil@gsid.org; Phillip W Berman - pwb@soe.ucs.edu; Keith A Crandall - kcrandall@genoma-llc.com

* Corresponding author

Published: 15 July 2009

Received: 3 May 2009

Retrovirology 2009, **6**:67 doi:10.1186/1742-4690-6-67

Accepted: 15 July 2009

This article is available from: <http://www.retrovirology.com/content/6/1/67>

© 2009 Pérez-Losada et al; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Differences in HIV-1 gp120 sequence variation were examined in North American volunteers who became infected during a phase III vaccine trial using the rgp120 vaccine. Molecular adaptation of the virus in vaccine and placebo recipients from different ethnic subgroups was compared by estimating the d_N/d_S ratios in viruses sampled from each individual using three different methods. ANOVA analyses detected significant differences in d_N/d_S ratios among races ($P < 0.02$). gp120 sequences from the black individuals showed higher mean d_N/d_S ratios for all estimators (1.24–1.45) than in other races (0.66–1.35), and several pairwise comparisons involving blacks remained significant ($P < 0.05$) after correction for multiple tests. In addition, black-placebo individuals showed significantly ($P < 0.02$) higher mean d_N/d_S ratios (1.3–1.66) than placebo individuals from the other races (0.65–1.56). These results suggest intrinsic differences among races in immune response and highlight the need for including multiple ethnicities in the design of future HIV-1 vaccine studies and trials.

Findings

More than 33 million people are currently infected with HIV-1, resulting in 2–3 million deaths every year. Natural immunity to the virus is virtually nonexistent; hence, the creation of a vaccine to combat this global pandemic is an international public-health priority [1,2]. In 2003, the results were released for a phase III HIV-1 vaccine efficacy trial conducted in North America and The Netherlands (VAX004) [3]. This study tested the efficacy of bivalent vaccines containing recombinant HIV-1 envelope glycoprotein 120 (rgp120) antigens, the major antigen on the

surface of the virus [4]. Overall, the vaccine candidate did not seem to reduce the incidence of HIV-1 infection, but an interesting trend was noted in the analysis of the different self-described ethnic groups [white (non-Hispanic), blacks, Hispanic, Asian, and "others"]. When only the non-white volunteers (17% of the total study population) were considered, the vaccine seemed to confer a slight benefit ($P = 0.012$). After adjustment for multiple tests, this difference was not significant ($P = 0.13$) [3]. Despite a lack of statistical support, this is not a trivial result. If this trend was to be confirmed, it would imply that non-

whites developed protective immunity to HIV-1 or, even more important, that rgp120 immunogens could protect against HIV infection under certain circumstances. Here we explored this possibility further.

HIV-1 evolution is driven, to a significant extent, by the immune response. If viruses isolated from non-white patients are in fact under a stronger selection pressure either because of genetic differences in the magnitude, specificity, or potency of the natural immune response, or because of differences in factors affecting virus replication, we should expect higher ratios of nonsynonymous (amino acid changing) to synonymous nucleotide substitutions (d_N/d_S) [5-8] than in viral samples isolated from vaccinated and placebo (non-vaccinated) white individuals.

To test whether levels of selection were significantly different between vaccinated and placebo individuals in different races, we analyzed 3 clones per individual from 345 infected North Americans from the VAX004 study (Table 1; data available at <http://www.gsid.org>). Full-length HIV-1 subtype B gp120 sequences were amplified as described in Gilbert et al. [9]. Since, as expected, viruses isolated from individuals from the same race did not form monophyletic groups [10], viral samples for each patient were analyzed separately. In each case, individual clones were aligned in MAFFT v5.7 [11], and d_N/d_S ratios were estimated using Nei and Gojobori's method [6] in SNAP [12], model M0 (one-ratio) in PAML v3.14 [13], and Fixed Effects Likelihood (FEL) with tree branch correction in HYPHY [14]. In the latter case, we took recombination into account by first detecting recombination breakpoints with GARD [15], and then estimating the d_N/d_S ratios independently for each fragment.

Mean d_N/d_S ratios across races and treatments were compared using ANOVA, linear models (*lm*) and pairwise t-tests. Because treating all non-whites as a single unit is unrealistic considering their own genetic differences [3], we tested for differences in selection pressure on a race-by-race basis. Multiple significance in the pairwise t-tests was corrected using the Benjamini and Hochberg's procedure [16].

The estimates of d_N/d_S obtained with SNAP, PAML and HYPHY were all significantly correlated among the different estimators used (correlation coefficient > 0.85; $P < 0.001$). Importantly, the mean d_N/d_S ratios varied across races (Table 1), and these differences were globally significant (ANOVA; $P < 0.02$) for SNAP and PAML estimates. Blacks (vaccinees and placebo combined) showed higher d_N/d_S ratios for all the estimators than individuals from other ethnicities (Table 1). Significant differences (pairwise t-tests; $P < 0.05$) between black and white, Hispanic and "others" viral samples were observed for all the estimators before corrections, but only the comparisons involving SNAP and PAML estimates remained significant after the Benjamini and Hochberg's adjustment (Table 2). The higher d_N/d_S ratios observed for blacks suggest that the rate of virus evolution is greater in this group than in other volunteers. Differences in immune response to HIV-1 infection have been pointed out by the rgp120 HIV Vaccine Study Group [3] as one of the potential factors to explain vaccine efficiency differences between white and non-white volunteers in the VAX004 trial. Ethnic differences in immune response have been also reported for other viruses such as the hepatitis C virus [17].

Does the greater virus adaptive variation presumed in black participants reflect genetic differences in the intrinsic (no-preconditioned) immune response to HIV-1, or is it a consequence of the conditioned immune response induced by vaccination with rgp120? Comparison of vaccine and placebo recipients showed different results based on the d_N/d_S estimators used. No significant differences were observed among vaccinees, but significant differences (ANOVA; $P = 0.025$) in SNAP d_N/d_S ratios were detected among placebo individuals. Moreover, black-placebo patients showed significantly (*lm* coefficients; $P < 0.02$) higher mean d_N/d_S ratios (1.3, 1.38 and 1.66, for SNAP, PAML and HYPHY, respectively) than the other races (0.65–1.01, 0.84–1.14 and 0.73–1.56, respectively). These results might indicate that natural differences in the immune response may have increased viral rgp120 adaptation in blacks.

In North America, blacks correspond to 42% of all newly diagnosed HIV/AIDS cases, while white (non-Hispanic) and Hispanic individuals represent approximately 40% and 17%, respectively [18]. If more data including both

Table 1: Mean d_N/d_S estimates across patients in PAML, SNAP and HYPHY.

	White (291)	Hispanic (22)	Black (12)	Asian (5)	Other (15)
SNAP	0.709	0.750	1.240	0.818	0.659
PAML	0.839	0.826	1.411	1.346	0.751
HYPHY	0.949	0.729	1.453	0.720	1.202

Individuals analyzed are indicated between parentheses.

Table 2: Statistically significant comparisons of d_N/d_S estimates among races and race-treatments.

Method	Race		Race by treatment	
	ANOVA	Corrected pairwise t-tests	ANOVA	<i>Im</i> coefficient
SNAP	(0.011)	black vs hispanic (0.020) black vs other (0.013) black vs white (0.004)	(0.025)	black placebo (0.001)
PAML	(0.019)	black vs hispanic (0.047) black vs other (0.047) black vs white (0.033)		black placebo (0.016)
HYPHY				black placebo (0.015)

Values in parentheses are *P*-values.

placebo and vaccinated recipients confirm that selection pressure differs between viruses infecting these three races, deciphering the genetic determinants of these differences should become a public-health priority. Indeed, our results highlight the need for selecting a broader representation of volunteers, based on ethnicity, in the design of future HIV-1 vaccine studies and trials [19].

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

MPL, DP, and KC developed the genetic and statistical strategies implemented in this work. MPL, DP, and MA performed the genetic and statistical analyses. DVJ, FS, and PWB carried out the molecular genetic studies and immunoassays. All authors participated in the design of the study and helped to draft the manuscript. All authors read and approved the final manuscript.

Acknowledgements

This study was supported by a Bill & Melinda Gates Foundation grant to Global Solutions for Infectious Diseases. It was also supported by the Spanish Ministry of Science and Education [grant number BIO2007-61411 to DP, FPI fellowship BES-2005-9151 to MA]. We also want to thank the reviewers for their excellent suggestions.

References

- Esparza J, Bhamarapravati N: **Accelerating the development and future availability of HIV-1 vaccines: why, when, where, and how?** *Lancet* 2000, **355**:2061-2066.
- Spearman P: **HIV vaccine development: lessons from the past and promise for the future.** *Curr HIV Res* 2003, **1**:101-120.
- rgp120 HIV Vaccine Study Group: **Placebo controlled phase 3 trial of a recombinant glycoprotein 120 vaccine to prevent HIV-1 infection.** *rgp120 HIV Vaccine Study Group.* *J Infect Dis* 2005, **191**:654-665.
- Eggink D, Melchers M, Sanders RW: **Antibodies to HIV-1: aiming at the right target.** *Trends Microbiol* 2007, **15**:291-294.
- Sharp PM: **In search of molecular Darwinism.** *Nature* 1997, **385**:111-112.
- Nei M, Gojobori T: **Simple methods for estimating the numbers of synonymous and nonsynonymous nucleotide substitutions.** *Mol Biol Evol* 1986, **3**:418-426.
- Ross HA, Rodrigo AG: **Immune-mediated positive selection drives human immunodeficiency virus type I molecular var-**

- iation and predicts disease duration.** *J Virol* 2002, **76**:11715-11720.
- Williamson S: **Adaptation in the env gene of HIV-1 and evolutionary theories of disease progression.** *Mol Biol Evol* 2003, **20**:1318-1325.
- Gilbert PB, Peterson ML, Follmann D, Hudgens MG, Francis DP, Gurwith M, Heyward WL, Jobes DV, Popovic V, Self SG, et al.: **Correlation between immunologic responses to a recombinant glycoprotein 120 vaccine and incidence of HIV-1 infection in a phase 3 HIV-1 preventive vaccine trial.** *J Infect Dis* 2005, **191**:666-677.
- Pérez-Losada M, Jobes DV, Sinangil F, Crandall KA, Posada D, Berman PW: **Phylogenetics of gp120 sequences from a Phase 3 HIV-1 vaccine trial in North America.** *Mol Biol Evol* 2009. in review
- Katoh K, Kuma K, Toh H, Miyata T: **MAFFT version 5: improvement in accuracy of multiple sequence alignment.** *Nucleic Acids Res* 2005, **33**:511-518.
- Korber B: **HIV signature and sequence variation analysis.** In *Computational and evolutionary analysis of HIV molecular sequences Volume 4*. Edited by: Rodrigo AG, Learn GH. Dordrecht, Netherlands: Kluwer Academic Publishers; 2000:55-72.
- Yang Z: **PAML: Phylogenetic Analysis by Maximum Likelihood.** 3.1th edition. London: University College London; 2001.
- Kosakovskiy Pond SL, Frost SDW, Muse SV: **HyPhy: hypothesis testing using phylogenies.** *Bioinformatics* 2005, **21**:676-679.
- Kosakovskiy Pond SL, Posada D, Gravenor MB, Woelk CH, Frost SD: **GARD: a genetic algorithm for recombination detection.** *Bioinformatics* 2006, **22**:3096-3098.
- Benjamini Y, Hochberg Y: **Controlling the false discovery rate: a practical and powerful approach to multiple testing.** *J R Statist Soc B* 1995, **57**:289-300.
- Sugimoto K, Stadanlick J, Ikeda F, Brensinger C, Furth EE, Alter HJ, Chang KM: **Influence of ethnicity in the outcome of hepatitis C virus infection and cellular immune response.** *Hepatology* 2003, **37**:590-599.
- CDC: **HIV/AIDS Surveillance Report.** Volume 19. Atlanta: Centers for Disease Control and Prevention (CDC); 2007.
- Graham BS, Mascola JR: **Lessons from failure – preparing for future HIV-1 vaccine efficacy trials.** *J Infect Dis* 2005, **191**:647-649.