

Nagoya Math. J. **201** (2011), 23–43
DOI 10.1215/00277630-2010-016

DISTRIBUTION OF POLYNOMIALS WITH CYCLES OF A GIVEN MULTIPLIER

GIOVANNI BASSANELLI AND FRANÇOIS BERTELOOT

Abstract. In the space of degree d polynomials, the hypersurfaces defined by the existence of a cycle of period n and multiplier $e^{i\theta}$ are known to be contained in the bifurcation locus. We prove that these hypersurfaces equidistribute the bifurcation current. This is a new result, even for the space of quadratic polynomials.

§1. Introduction

In a holomorphic family $(f_\lambda)_{\lambda \in M}$ of rational maps, the sets $\text{Per}_n(w)$ of parameters for which f_λ has a cycle of exact period n and multiplier w turn out to be hypersurfaces. One knows, since the fundamental work of Mañé, Sad, and Sullivan [15], that the closure of the union of the hypersurfaces $\text{Per}_n(e^{i\theta})$ coincides with the bifurcation locus $\mathcal{Bif}(M)$, that is, the set of parameters λ_0 for which the dynamics of f_{λ_0} drastically change under small perturbation. Our aim here is to describe precisely, from a measure-theoretic point of view, the asymptotic behavior of $\text{Per}_n(e^{i\theta})$ as the period n grows.

Our main tool is the *bifurcation current* T_{bif} introduced by DeMarco [8]. It is a positive, closed $(1, 1)$ -current supported by $\mathcal{Bif}(M)$ which admits both the Lyapunov function $L(\lambda)$ and the sum of values of the Green function on critical points as potentials (see [9] or [1]):

$$T_{\text{bif}} = dd^c L(\lambda) = dd^c \sum G_\lambda(c_\lambda).$$

The bifurcation current and its powers T_{bif}^k ($k \leq \dim_{\mathbb{C}} M$) have been used in several recent works (see [1], [12], [11], [6], [2]) devoted to the study of measurable or complex analytic properties of the bifurcation locus. In particular, Dujardin and Favre ([12, Theorems 1 and 4.2]) have used the Green-like potentials to get equidistribution results concerning the hypersurfaces of M defined by the preperiodicity of a critical point.

Received May 13, 2009. Accepted March 25, 2010.

2000 *Mathematics Subject Classification.* Primary 37F45; Secondary 32U15.

© 2011 by The Editorial Board of the *Nagoya Mathematical Journal*

In order to study the asymptotic distribution of the hypersurfaces $\text{Per}_n(w)$, it is more convenient to use the Lyapunov function since $L(\lambda)$ is well related to the multipliers of n -periodic repelling cycles (see Theorem 2.3). This dynamical property allows us to compare $L(\lambda)$ with $d^{-n} \ln |p_n(\lambda, w)|$, where the functions $p_n(\cdot, w)$ canonically define the hypersurfaces $\text{Per}_n(w)$. Setting $[\text{Per}_n(w)] := dd^c \ln |p_n(\lambda, w)|$ and using basic potential-theoretic tools, we get the following general equidistribution statements.

THEOREM 1.1. *Let T_{bif} be the bifurcation current of some holomorphic family of rational maps $(f_\lambda)_{\lambda \in M}$. Then*

$$\begin{aligned} d^{-n}[\text{Per}_n(w)] &\rightarrow T_{\text{bif}}, \quad \text{when } |w| < 1, \\ \frac{d^{-n}}{2\pi} \int_0^{2\pi} [\text{Per}_n(re^{i\theta})] d\theta &\rightarrow T_{\text{bif}}, \quad \text{when } r \geq 0, \text{ and} \\ d^{-n} dd^c_{(\lambda,w)} \ln |p_n(\lambda, w)| &\rightarrow dd^c L(\lambda), \end{aligned}$$

where the convergence is weak, occurs in M and, for the last statement, in $M \times \mathbf{C}$.

The first two assertions are quite easily obtained and were actually essentially given in [2] (the case $w = 0$ is also covered by Dujardin and Favre’s result).

When $|w| = 1$, the convergence of $d^{-n}[n(w)]$ would easily follow from our arguments if the density of hyperbolic parameters were known. Our basic observation is that the same conclusion occurs when M is a Riemann surface, in which the set of nonhyperbolic parameters is compact and the bifurcation locus is contained in the closure of hyperbolic parameters (see Proposition 3.4). This already covers the case of the family of quadratic polynomials. To go further, our strategy is therefore to slice M with hypersurfaces which are chosen for the good repartition of hyperbolic parameters. The existence of such slices is intimately related to the behavior at infinity of the bifurcation locus, and for this reason we work in the family of degree d polynomials. In this family, the control we need actually follows from the work of Branner and Hubbard [5] on the compactness of the connectedness locus (see Theorem 4.2). Our main result is the following.

THEOREM 1.2. *Let $d \geq 2$, and let $\{P_{c,a}\}_{(c,a) \in \mathbf{C}^{d-1}}$ be the holomorphic family of degree d polynomials parameterized by defining $P_{c,a}$ as the polynomial of degree d whose critical points are $(0, c_1, \dots, c_{d-2})$ and such that $P_{c,a}(0) = a^d$.*

Let T_{bif} be the bifurcation current of this family. Then $\lim_n d^{-n} [\text{Per}_n(w)] = T_{\text{bif}}$ for any w such that $|w| \leq 1$.

§2. Some tools

2.1. Hypersurfaces $\text{Per}_n(w)$

For any holomorphic family of rational maps, the following result describes precisely the set of maps having a cycle of given period and multiplier.

THEOREM 2.1. *Let $f : M \times \mathbf{P}^1 \rightarrow \mathbf{P}^1$ be a holomorphic family of degree $d \geq 2$ rational maps. Then for every integer $n \in \mathbf{N}^*$, there exists a holomorphic function p_n on $M \times \mathbf{C}$ which is polynomial on \mathbf{C} and such that*

- (1) *for any $w \in \mathbf{C} \setminus \{1\}$, the function $p_n(\lambda, w)$ vanishes if and only if f_λ has a cycle of exact period n and multiplier w ;*
- (2) *$p_n(\lambda, 1) = 0$ if and only if f_λ has a cycle of exact period n and multiplier 1 or a cycle of exact period m whose multiplier is a primitive r th root of unity with $r \geq 2$ and $n = mr$;*
- (3) *for every $\lambda \in M$, the degree $N_d(n)$ of $p_n(\lambda, \cdot)$ satisfies $d^{-n} N_d(n) \sim 1/n$.*

This leads to the following definitions. For any integer n and any $w \in \mathbf{C}$, the subset $\text{Per}_n(w)$ of M is the hypersurface given by

$$\text{Per}_n(w) := \{ \lambda \in M / p_n(\lambda, w) = 0 \}$$

and, taking into account the possible multiplicities, we consider the following integration currents:

$$[\text{Per}_n(w)] := dd^c_\lambda \ln |p_n(\lambda, w)|.$$

Let us briefly recall the construction of the functions p_n . (For more details, we refer the reader to Milnor [16] or Silverman [17, Chapter 4].)

One first constructs the dynatomic polynomials $\Phi_{\varphi, n}^*$ associated to a rational map φ of degree $d \geq 2$. Let us denote by $F^n = (F_1^n, F_2^n)$ the iterates of some lift F of φ to \mathbf{C}^2 , and let us define homogeneous polynomials $\Phi_{\varphi, n}$ on \mathbf{C}^2 by setting

$$\Phi_{\varphi, n}(X, Y) := YF_1^n(X, Y) - XF_2^n(X, Y).$$

The divisor $\text{Div}(\Phi_{\varphi, n})$ induced by $\Phi_{\varphi, n}(X, Y)$ on \mathbf{P}^1 is precisely the set of periodic points of φ with exact period dividing n . Denoting μ the classical Möbius function, one then sets

$$\Phi_{\varphi, n}^*(X, Y) := \prod_{k|n} (\Phi_{\varphi, n}(X, Y))^{\mu(n/k)}.$$

Using the fact that the sum $\sum_{k|n} \mu(k/n)$ vanishes if $n > 1$ and that it is equal to 1 if $n = 1$, one may show that $\Phi_{\varphi,n}^*$ is a polynomial whose degree $\nu_d(n)$ depends only on n and d . The divisor $\text{Div}(\Phi_{\varphi,n}^*)$ induced by $\Phi_{\varphi,n}^*(X, Y)$ on \mathbf{P}^1 clearly contains the periodic points of φ with exact period equal to n . The other points contained in $\text{Div}(\Phi_{\varphi,n}^*)$ are precisely the periodic points of φ whose exact period m divides n ($m = nr, r \geq 2$) and whose multiplier is a primitive r th root of unity (see [17, Theorem 4.5, p. 151]).

If $z \in \text{Div}(\Phi_{\varphi,n}^*)$ has exact period m with $n = mr$, then we will denote by $w_n(z)$ the r th power of the multiplier of z (i.e., $(\varphi^n)'(z)$ in good coordinates). One sees in particular that the following fact occurs: a point z is periodic of exact period n and $w_n(z) \neq 1$ if and only if $z \in \text{Div}(\Phi_{\varphi,n}^*)$ and $w_n(z) \neq 1$.

Let us now consider the sets

$$\Lambda_n^*(\varphi) := \{w_n(z); z \in \text{Div}(\Phi_{\varphi,n}^*)\},$$

where the points in $\text{Div}(\Phi_{\varphi,n}^*)$ are counted with multiplicity, and let us denote by $\sigma_i^{*(n)}(\varphi)$, $1 \leq i \leq \nu_d(n)$ the associated symmetric functions. We define the polynomials $p_n(\varphi, w)$ by

$$(p_n(\varphi, w))^n := \prod_{i=0}^{\nu_d(n)} \sigma_i^{*(n)}(\varphi)(-w)^{\nu_d(n)-i},$$

and therefore $p_n(\varphi, w) = 0$ if and only if $w \in \Lambda_n^*(\varphi)$. The properties of p_n follow easily from this construction. The degree $N_d(n)$ of $p_n(\lambda, \cdot)$ is equal to $(1/n)\nu_d(n) = (1/n)\sum_{k|n} \mu(n/k)d^k$. In particular, $d^{-n}N_d(n) \sim 1/n$.

2.2. Lyapunov exponent and bifurcation current

Every rational map of degree $d \geq 2$ on the Riemann sphere admits a maximal entropy measure μ_f . The Lyapunov exponent of f with respect to the measure μ_f is given by $L(f) = \int_{\mathbf{P}^1} \log |f'| \mu_f$ (see [10] for a general exposition in any dimension).

When $f : M \times \mathbf{P}^1 \rightarrow \mathbf{P}^1$ is a holomorphic family of degree d rational maps, the Lyapunov function L on the parameter space M is defined by

$$L(\lambda) = \int_{\mathbf{P}^1} \log |f'_\lambda| \mu_\lambda,$$

where μ_λ is the maximal entropy measure of f_λ . The function L is plurisubharmonic (PSH) on M , and the bifurcation current T_{bif} of the family is a closed, positive $(1, 1)$ -current on M which may be defined by

$$T_{\text{bif}} := dd^c L(\lambda).$$

As it has been shown by DeMarco [9], the support of T_{bif} coincides with the bifurcation locus of the family in the sense of Mañé, Sad, and Sullivan [15] (see also [1, Theorem 5.2]).

Let us recall that Mañé, Sad, and Sullivan have shown that the complement of the bifurcation locus is a dense open subset of the parameter space M whose connected components are called *stable components*. They have also shown that any neutral cycle is persistent on the stable components. In the language of Theorem 2.1, this property may be expressed as follows.

REMARK 2.2. For $|w_0| = 1$, a function $p_n(\lambda, w_0)$ either does not vanish on any stable component or vanishes identically on M .

In our study, we combine classical potential-theoretic methods with the following dynamical property (see [3] or [4], where this has been proved for endomorphisms of \mathbf{P}^k).

THEOREM 2.3. *Let $f : \mathbf{P}^1 \rightarrow \mathbf{P}^1$ be a degree $d \geq 2$ rational map, let μ be its maximal entropy measure, and let L be the Lyapunov exponent of f with respect to μ . Then*

$$L = \lim_n \frac{d^{-n}}{n} \sum_{p \in R_n^*} \ln |(f^n)'(p)|,$$

where $R_n^* := \{p \in \mathbf{P}^1 / p \text{ has exact period } n \text{ and } |(f^n)'(p)| > 1\}$.

The continuity of the Lyapunov function will also play a crucial role. This was proved by Mañé [14], but a simple argument based on DeMarco’s formula shows that this function is actually Hölder-continuous (see [1, Corollary 3.4]).

THEOREM 2.4. *Let $f : M \times \mathbf{P}^1 \rightarrow \mathbf{P}^1$ be a holomorphic family of degree $d \geq 2$ rational maps. Let $L(\lambda)$ be the Lyapunov exponent of $(\mathbf{P}^1, f_\lambda, \mu_\lambda)$, where μ_λ is the maximal entropy measure of f_λ . Then the function $L(\lambda)$ is Hölder-continuous on M .*

We end this section by recalling a well-known compactness principle for subharmonic functions which will be used frequently in the paper.

THEOREM 2.5. *Let (φ_j) be a sequence of subharmonic functions which is locally uniformly bounded from above on some domain $\Omega \subset \mathbf{R}^n$. If (φ_j) does not converge to $-\infty$, then a subsequence (φ_{j_k}) converges in $L^1_{\text{loc}}(\Omega)$ to some subharmonic function φ . In particular, (φ_j) converges in $L^1_{\text{loc}}(\Omega)$ to some subharmonic function φ if it converges pointwise to φ .*

§3. Distribution of $\text{Per}_n(w)$ in general families

In this section, we consider an arbitrary holomorphic family $f : M \times \mathbf{P}^1 \rightarrow \mathbf{P}^1$ of degree $d \geq 2$ rational maps. We investigate the convergence of the currents $(1/d^n)[\text{Per}_n(w)]$ towards the bifurcation current T_{bif} by considering the sequences of their potentials, and therefore compare the Lyapunov function L with the limits of $(1/d^n) \ln |p_n(\lambda, w)|$, where $p_n(\lambda, w)$ are the polynomials given by Theorem 2.1.

This leads us to consider the following sequences of PSH functions

$$L_n^r(\lambda) := \frac{d^{-n}}{2\pi} \int_0^{2\pi} \ln |p_n(\lambda, re^{i\theta})| d\theta,$$

$$L_n^+(\lambda, w) := d^{-n} \sum_{j=1}^{N_d(n)} \ln^+ |w - w_{n,j}(\lambda)|,$$

$$L_n(\lambda, w) := d^{-n} \ln |p_n(\lambda, w)|,$$

where $p_n(\lambda, w) =: \prod_{j=1}^{N_d(n)} (w - w_{n,j}(\lambda))$ are the polynomials associated to the family f by Theorem 2.1.

The pointwise convergence of $L_n(\lambda, w)$ to L for $|w| < 1$ is quite a straightforward consequence of Theorem 2.3 and immediately implies that $d^{-n}[\text{Per}_n(w)]$ converges to T_{bif} when $|w| < 1$. However, when $|w| \geq 1$ and λ is a nonhyperbolic parameter, the control of $L_n(\lambda, w) = d^{-n} \sum \ln |w - w_{n,j}(\lambda)|$ is very delicate because f_λ may have many cycles whose multipliers are close to w . This is why we introduce the PSH functions L_n^+ , which both coincide with L_n on the hyperbolic components and are quite easily seen to converge nicely.

3.1. A basic result

We present here what is obtained by combining the dynamical Theorem 2.3 with basic potential-theoretic facts. Our main result is the following.

THEOREM 3.1. *Let $f : M \times \mathbf{P}^1 \rightarrow \mathbf{P}^1$ be a holomorphic family of degree $d \geq 2$ rational maps. Let $L(\lambda)$ be the Lyapunov exponent of $(\mathbf{P}^1, f_\lambda, \mu_\lambda)$, where μ_λ is the maximal entropy measure of f_λ . Let $(L_n)_n$, $(L_n^r)_n$, and $(L_n^+)_n$ be the sequences of PSH functions defined as above. Then*

- (1) *the sequence L_n converges pointwise to L on $M \times \Delta$ and, for any $w \in \Delta$, the sequence $L_n(\cdot, w)$ converges in L_{loc}^1 to L on M ;*

- (2) the sequence L_n^r converges pointwise and in L_{loc}^1 to L on M for $r \geq 0$;
- (3) the sequence L_n^+ converges pointwise and in L_{loc}^1 to L on $M \times \mathbf{C}$ and, for every $w \in \mathbf{C}$, the sequence $L_n^+(\cdot, w)$ converges in L_{loc}^1 to L on M ;
- (4) the sequence L_n converges in L_{loc}^1 to L on $M \times \mathbf{C}$.

Let us stress that Theorem 1.1 follows immediately from the first, second, and last statements of Theorem 3.1 by taking dd^c .

Proof. All the statements are local, and therefore, taking charts, we may assume that $M = \mathbf{C}^k$. We write the polynomials p_n as follows:

$$p_n(\lambda, w) =: \prod_{i=1}^{N_d(n)} (w - w_{n,i}(\lambda)).$$

Throughout the proof, we shall use the fact that $d^{-n}N_d(n) \sim 1/n$ (see Theorem 2.1). In particular, this implies that the sequences L_n and L_n^+ are locally uniformly bounded from above.

• We first establish the convergence of $L_n(\lambda, w)$ when $|w| < 1$. According to Theorem 2.1, the set $\{w_{n,j}(\lambda)/w_{n,j}(\lambda) \neq 1\}$ coincides with the set of multipliers of cycles of exact period n (counted with multiplicity) from which the cycles of multiplier 1 are deleted. Using the notation $R_n^*(\lambda) := \{p \in \mathbf{P}^1/p \text{ has exact period } n \text{ and } |(f_\lambda^n)'(p)| > 1\}$, we thus have

$$(3.1) \quad \sum_{j=1}^{N_d(n)} \ln^+ |w_{n,j}(\lambda)| = \frac{1}{n} \sum_{p \in R_n^*(\lambda)} \ln |(f^n)'(p)|.$$

Since f_λ has a finite number of nonrepelling cycles (Fatou’s theorem), one sees that there exists $n(\lambda) \in \mathbf{N}$ such that

$$(3.2) \quad n \geq n(\lambda) \Rightarrow |w_{n,j}(\lambda)| > 1, \quad \text{for any } 1 \leq j \leq N_d(n).$$

By (3.1) and (3.2), one gets

$$\begin{aligned} L_n(\lambda, 0) &= d^{-n} \sum_{j=1}^{N_d(n)} \ln |w_{n,j}(\lambda)| \\ &= d^{-n} \sum_{j=1}^{N_d(n)} \ln^+ |w_{n,j}(\lambda)| = \frac{d^{-n}}{n} \sum_{R_n^*(\lambda)} \ln |(f^n)'(p)| \end{aligned}$$

for $n \geq n(\lambda)$ which, by Theorem 2.3, yields

$$(3.3) \quad \lim_n L_n(\lambda, 0) = L(\lambda), \quad \forall \lambda \in M.$$

Let us now pick $w \in \Delta$. By (3.2), we have $L_n(\lambda, w) - L_n(\lambda, 0) = d^{-n} \times \sum_j \ln(|w_{n,j}(\lambda) - w|/|w_{n,j}(\lambda)|)$ and $\ln(1 - |w|) \leq \ln(|w_{n,j}(\lambda) - w|/|w_{n,j}(\lambda)|) \leq \ln(1 + |w|)$ for $1 \leq j \leq N_d(n)$ and $n \geq n(\lambda)$. We thus get

$$d^{-n} N_d(n) \ln(1 - |w|) \leq |L_n(\lambda, w) - L_n(\lambda, 0)| \leq d^{-n} N_d(n) \ln(1 + |w|)$$

for $n \geq n(\lambda)$ and, using (3.3), we get $\lim_n L_n(\lambda, w) = L(\lambda)$ for any $(\lambda, w) \in M \times \Delta$.

The L^1_{loc} convergence of $L_n(\cdot, w)$ now follows immediately from Theorem 2.5.

• Let us show that the convergence of $L_n(\lambda, 0) = L_n^0$ implies the convergence of L_n^r for any $r > 0$. We essentially show that $\lim_n |L_n^r(\lambda) - L_n(\lambda, 0)| = 0$ by using the formula $\ln \max(|a|, r) = (1/2\pi) \int_0^{2\pi} \ln |a - re^{i\theta}| d\theta$. Indeed, this formula yields

$$L_n^r(\lambda) = \frac{1}{2\pi d^n} \int_0^{2\pi} \ln \prod_j |re^{i\theta} - w_{n,j}(\lambda)| d\theta = d^{-n} \sum_j \ln \max(|w_{n,j}(\lambda)|, r).$$

Since $|w_{n,j}(\lambda)| \geq 1$ for $n \geq n(\lambda)$ (see (3.2)), we deduce from the above identity that

$$\begin{aligned} L_n^r(\lambda) &= d^{-n} \sum_j \ln |w_{n,j}(\lambda)| + d^{-n} \sum_{1 \leq |w_{n,j}(\lambda)| < r} \ln \frac{r}{|w_{n,j}(\lambda)|} \\ &= L_n(\lambda, 0) + d^{-n} \sum_{1 \leq |w_{n,j}(\lambda)| < r} \ln \frac{r}{|w_{n,j}(\lambda)|}, \end{aligned}$$

and thus

$$0 \leq L_n^r(\lambda) - L_n(\lambda, 0) = d^{-n} \sum_{1 \leq |w_{n,j}(\lambda)| < r} \ln \frac{r}{|w_{n,j}(\lambda)|} \leq d^{-n} N_d(n) \ln^+ r.$$

By (3.3), this implies that L_n^r converges pointwise to L . It also shows that $(L_n^r)_n$ is locally uniformly bounded from above which, by Theorem 2.5, implies that $(L_n^r)_n$ converges to L in $L^1_{\text{loc}}(M)$.

• Let us now deal with the convergence of L_n^+ . We will show that $L_n^+(\cdot, w)$ converges pointwise to L on M for every $w \in \mathbf{C}$. As $(L_n^+)_n$ is locally uniformly bounded, this implies the convergence of $L_n^+(\cdot, w)$ in $L^1_{\text{loc}}(M)$ (see

Theorem 2.5), and the convergence of L_n^+ in $L^1_{\text{loc}}(M \times \mathbf{C})$ then follows by Lebesgue’s theorem.

We have to estimate $L_n^+(\lambda, w) - L_n(\lambda, 0) =: \epsilon_n(\lambda, w)$ on M . Let us fix $\lambda \in M$, $w \in \mathbf{C}$ and pick $R > |w|$. We may assume that $n \geq n(\lambda)$ so that $|w_{n,j}(\lambda)| \geq 1$ for all $1 \leq j \leq N_d(n)$ (see (3.2)), and we then decompose $\epsilon_n(\lambda, w)$ in the following way:

$$\begin{aligned} \epsilon_n(\lambda, w) &= d^{-n} \sum_{1 \leq |w_{n,j}(\lambda)| < R+1} \ln^+ |w_{n,j}(\lambda) - w| \\ &\quad + d^{-n} \sum_{|w_{n,j}(\lambda)| \geq R+1} \ln \frac{|w_{n,j}(\lambda) - w|}{|w_{n,j}(\lambda)|} \\ &\quad - d^{-n} \sum_{1 \leq |w_{n,j}(\lambda)| < R+1} \ln |w_{n,j}(\lambda)|. \end{aligned}$$

We may write this decomposition as $\epsilon_n(\lambda, w) =: \epsilon_{n,1}(\lambda, w) + \epsilon_{n,2}(\lambda, w) - \epsilon_{n,1}(\lambda, 0)$. As $L_n(\lambda, 0)$ converges to L , we simply have to check that $\epsilon_{n,1}(\lambda, w)$ and $\epsilon_{n,2}(\lambda, w)$ tend to 0 when n tends to ∞ . One clearly has $0 \leq \epsilon_{n,1}(\lambda, w) \leq d^{-n} N_d(n) \ln(2R + 1)$, and thus $\lim_n \epsilon_{n,1}(\lambda, w) = 0$. Similarly, $\lim_n \epsilon_{n,2}(\lambda, w) = 0$ follows from the fact that, for $|w_{n,j}(\lambda)| > R + 1 > |w| + 1$, one has

$$\begin{aligned} \ln\left(1 - \frac{R}{R+1}\right) &\leq \ln \frac{|w_{n,j}(\lambda)| - R}{|w_{n,j}(\lambda)|} \leq \ln \frac{|w_{n,j}(\lambda) - w|}{|w_{n,j}(\lambda)|} \\ &\leq \ln \frac{|w_{n,j}(\lambda)| + R}{|w_{n,j}(\lambda)|} \leq \ln\left(1 + \frac{R}{R+1}\right). \end{aligned}$$

- We are finally ready to prove the L^1_{loc} convergence of $(L_n)_n$. As the functions L_n are PSH and the sequence $(L_n)_n$ is locally uniformly bounded from above, we shall again use the compactness properties of PSH functions given by Theorem 2.5. Since $L_n(\lambda, 0)$ converges to $L(\lambda)$, the sequence $(L_n)_n$ does not converge to $-\infty$, and it therefore suffices to show that, among PSH functions on $M \times \mathbf{C}$, the function L is the only possible limit for $(L_n)_n$ in $L^1_{\text{loc}}(M \times \mathbf{C})$.

Let φ be a PSH function on $M \times \mathbf{C}$, and let $(L_{n_j})_j$ be a subsequence of $(L_n)_n$ which converges to φ in $L^1_{\text{loc}}(M \times \mathbf{C})$. Pick $(\lambda_0, w_0) \in M \times \mathbf{C}$. We have to prove that $\varphi(\lambda_0, w_0) = L(\lambda_0)$.

Let us first observe that $\varphi(\lambda_0, w_0) \leq L(\lambda_0)$. Take a ball B_ϵ of radius ϵ and centered at $(\lambda_0, w_0) \in M \times \mathbf{C}$. By the submean value property and the

L^1_{loc} -convergence of L_n^+ , we have

$$\begin{aligned} \varphi(\lambda_0, w_0) &\leq \frac{1}{|B_\epsilon|} \int_{B_\epsilon} \varphi \, dm = \lim_j \frac{1}{|B_\epsilon|} \int_{B_\epsilon} L_{n_j} \, dm \\ &\leq \lim_j \frac{1}{|B_\epsilon|} \int_{B_\epsilon} L_{n_j}^+ \, dm = \frac{1}{|B_\epsilon|} \int_{B_\epsilon} L \, dm, \end{aligned}$$

and then, making $\epsilon \rightarrow 0$, one obtains $\varphi(\lambda_0, w_0) \leq L(\lambda_0)$.

Let us now check that $\limsup_j L_{n_j}(\lambda_0, w_0 e^{i\theta}) = L(\lambda_0)$ for almost all $\theta \in [0, 2\pi]$. Let $r_0 := |w_0|$. As we saw, L_n^+ converges pointwise to L , and therefore

$$\limsup_j L_{n_j}(\lambda_0, w_0 e^{i\theta}) \leq \limsup_j L_{n_j}^+(\lambda_0, w_0 e^{i\theta}) = L(\lambda_0).$$

On the other hand, by pointwise convergence of $L_n^{r_0}$ to L and Fatou's lemma, we have

$$\begin{aligned} L(\lambda_0) &= \lim_n L_n^{r_0}(\lambda_0) = \limsup_j \frac{1}{2\pi} \int_0^{2\pi} L_{n_j}(\lambda_0, r_0 e^{i\theta}) \, d\theta \\ &\leq \frac{1}{2\pi} \int_0^{2\pi} \limsup_j L_{n_j}(\lambda_0, r_0 e^{i\theta}) \, d\theta, \end{aligned}$$

and the desired property follows immediately.

To end the proof, we argue by contradiction and assume that $\varphi(\lambda_0, w_0) < L(\lambda_0)$. As φ is upper semicontinuous and L -continuous (see Theorem 2.4), there exists a neighborhood V_0 of (λ_0, w_0) and $\epsilon > 0$ such that

$$\varphi - L \leq -\epsilon \text{ on } V_0.$$

Pick a small ball B_{λ_0} centered at λ_0 and a small disc Δ_{w_0} centered at w_0 such that $B_0 := B_{\lambda_0} \times \Delta_{w_0}$ is relatively compact in V_0 . Then, according to Hartogs's lemma (see [13, page 94]), we have

$$\limsup_j \left(\sup_{B_0} (L_{n_j} - L) \right) \leq \sup_{B_0} (\varphi - L) \leq -\epsilon.$$

This is impossible since, as we have seen before, we may find $(\lambda_0, r_0 e^{i\theta_0}) \in B_0$ such that $\limsup_j (L_{n_j}(\lambda_0, r_0 e^{i\theta_0}) - L(\lambda_0)) = 0$. \square

REMARK 3.2. Using standard techniques, one may deduce from the fourth assertion of Theorem 3.1 that the set of multipliers w for which the bifurcation current T_{bif} is not a limit of the sequence $d^{-n}[\text{Per}_n(w)]$ is contained in a polar subset of the complex plane.

3.2. Further results

The fact that the functions L_n^+ and L_n coincide on hyperbolic components would easily yield the convergence of $d^{-n}[\text{Per}_n(w)]$ towards T_{bif} for any $w \in \mathbf{C}$ if the density of hyperbolic parameters in M were known.

One may, however, overcome this difficulty when the hyperbolic parameters are sufficiently nicely distributed. Here we establish a few facts of this nature which we use in our study of polynomial families in Section 4.

The following proposition summarizes some useful remarks.

PROPOSITION 3.3. *Let us make the same assumptions and adopt the same notations as in Theorem 3.1. Let $w_0 \in \mathbf{C}$. Then the following hold.*

- (1) *Any PSH limit value of $L_n(\lambda, w_0)$ in $L_{\text{loc}}^1(M)$ is smaller than L .*
- (2) *If a subsequence $L_{n_k}(\lambda, w_0)$ converges pointwise to L on the stable set, then it also converges to L in $L_{\text{loc}}^1(M)$.*
- (3) *Assume that $|w_0| = 1$ and that the family has no persistent neutral cycle. If a subsequence $L_{n_k}(\lambda, w_0)$ converges to φ in $L_{\text{loc}}^1(M)$, then φ is pluriharmonic on any stable component Ω and the convergence is locally uniform on Ω .*
- (4) *For any hyperbolic component $\Omega \subset M$, the sequence $L_n(\lambda, w_0)$ converges locally uniformly to L on Ω .*

Proof. (1) Let us set $\varphi_n(\lambda) := L_n(\lambda, w_0)$ and assume that a subsequence φ_{n_j} converges in $L_{\text{loc}}^1(M)$ to some PSH function φ . Since $L_n^+(\lambda, w_0)$ converges to L in $L_{\text{loc}}^1(M)$ and since $\varphi_{n_j}(\lambda) \leq L_{n_j}^+(\lambda, w_0)$, we get $\varphi(\lambda_0) \leq (1/|B_\epsilon|) \int_{B_\epsilon} \varphi \, dm \leq (1/|B_\epsilon|) \int_{B_\epsilon} L \, dm$ for any small ball B_ϵ centered at λ_0 . The desired inequality then follows by making $\epsilon \rightarrow 0$.

(2) Recall that the stable set is an open dense subset of M . Let φ be any PSH limit of $L_{n_k}(\lambda, w_0)$ in $L_{\text{loc}}^1(M)$. We have to show that $\varphi = L$. By the first assertion, $\varphi \leq L$. As $\varphi = L$ on a dense subset, the semicontinuity of φ and the continuity of L (see Theorem 2.4) imply that $\varphi \geq L$.

(3) Using Remark 2.2, one sees that the functions $L_{n_k}(\lambda, w_0)$ are pluriharmonic on Ω . This implies that φ itself is pluriharmonic on Ω and that $L_{n_k}(\lambda, w_0)$ converges actually locally uniformly on Ω to φ .

(4) If λ is a hyperbolic parameter, then f_λ has only attracting or repelling cycles and is expansive on its Julia set. Thus, as f_λ has at most a finite number of attracting cycles, one sees that $|w_{n,j}(\lambda)| \geq |w_0| + 1$ for all $1 \leq j \leq N_d(n)$ and n big enough. In other words, $L_n(\lambda, w_0) = L_n^+(\lambda, w_0)$ for n big enough, and therefore, according to Theorem 3.1, $L_n(\lambda, w_0)$ converges

to $L(\lambda)$. By Theorem 2.5, $L_n(\lambda, w_0)$ converges to L in $L^1_{\text{loc}}(\Omega)$. The local uniform convergence then follows from the previous assertion. \square

In the last section of the paper, we will focus on the case $|w_0| = 1$ and work with polynomial families. Slicing the parameter space in different ways, we will obtain 1-dimensional holomorphic families where the hyperbolic parameters are well distributed. Here is a typical example which, in particular, covers the case of the quadratic polynomial family.

PROPOSITION 3.4. *Let M be a Riemann surface, and let $f : M \times \mathbf{P}^1 \rightarrow \mathbf{P}^1$ be a holomorphic family of degree $d \geq 2$ rational maps which satisfies the following two conditions:*

- (1) *the bifurcation locus is contained in the closure of hyperbolic parameters;*
- (2) *the set of nonhyperbolic parameters is compact in M .*

Let $L(\lambda)$ and $L_n(\lambda, w)$ be the subharmonic functions defined in Theorem 3.1. Then, if $|w_0| = 1$, the sequence $L_n(\lambda, w_0)$ converges to L in $L^1_{\text{loc}}(M)$.

The proof of this proposition is based on the following technical lemma. This lemma actually deals with more general situations which we will encounter in Section 4.

LEMMA 3.5. *Let M be a Riemann surface, and let $f : M \times \mathbf{P}^1 \rightarrow \mathbf{P}^1$ be a holomorphic family of degree $d \geq 2$ rational maps. Let $L(\lambda)$ and $L_n(\lambda, w)$ be the subharmonic functions defined in Theorem 3.1. Let $w_0 \in \mathbf{C}$ with $|w_0| = 1$, and let φ be a subharmonic limit value of $L_n(\lambda, w_0)$ in $L^1_{\text{loc}}(M)$ such that*

- (1) *the bifurcation locus is contained in the closure of the set of parameters where $\varphi = L$;*
- (2) *$\varphi = L$ on the stable component which is not relatively compact in M .*

Then $\varphi = L$.

Proof. We start by proving the lemma. On several occasions we shall use the fact that the function L is continuous (see Theorem 2.4). Assume that $\varphi_{n_j} := L_n(\cdot, w_0)$ converges to φ . Then the holomorphic functions $p_{n_j}(\lambda, w_0)$ cannot vanish identically for j big enough. According to Remark 2.2, this implies that the functions φ_{n_j} are harmonic on all stable components of M . This leads to the simple but crucial observation that φ is harmonic on any stable component, or in other words, that the Laplacian $\Delta\varphi$ is supported in the bifurcation locus.

According to the first assertion of Proposition 3.3, we have $\varphi \leq L$. We may now see that $\varphi = L$ on the bifurcation locus. Indeed, if λ_0 belongs to

the bifurcation locus, then, by assumption, there exists a sequence λ_k which converges to λ_0 such that $\varphi(\lambda_k) = L(\lambda_k)$. Then, using the upper semicontinuity of φ and the continuity of L , we get $\varphi(\lambda_0) = \limsup_{\lambda \rightarrow \lambda_0} \varphi(\lambda) \geq \limsup \varphi(\lambda_k) = \lim L(\lambda_k) = L(\lambda_0)$.

By the first observation and the fact that L is continuous, we see that φ is continuous on the support of its Laplacian. According to some well-known continuity principle, this implies that φ is continuous on M . We may now prove that $\varphi \equiv L$. If this were not the case, then $\varphi(\lambda_0) < L(\lambda_0)$ for some $\lambda_0 \in M$. As L and φ coincide on the bifurcation locus and (by assumption) on nonrelatively compact stable components, λ_0 would belong to some stable component Ω which is relatively compact in M . This contradicts the maximum principle since $(\varphi - L)$ is continuous on $\overline{\Omega}$, harmonic on Ω , and vanishes on $\partial\Omega$.

Let us now establish the proposition. By the fourth assertion of Proposition 3.3, the sequence $L_n(\lambda, w_0)$ does not converge to $-\infty$. According to Theorem 2.5, it thus suffices to show that any subharmonic limit value φ of $L_n(\lambda, w_0)$ in $L^1_{\text{loc}}(M)$ coincides with L . This follows immediately from Lemma 3.5 since, once again by the fourth assertion of Proposition 3.3, $\varphi = L$ on the nonrelatively compact stable components. □

REMARK 3.6. The first assumption of Proposition 3.4 is a well-known open question in the space of polynomials of degree $d \geq 3$. This explains why a slicing argument will be used in the proof of Theorem 1.2.

§4. Distribution of $\text{Per}_n(w)$ in polynomial families

4.1. The space of degree d polynomials

Let \mathcal{P}_d be the space of polynomials of degree $d \geq 2$ with $d - 1$ marked critical points up to conjugacy by affine transformations. Although this space has a natural structure of affine variety of dimension $d - 1$, we shall actually work with a specific parameterization of \mathcal{P}_d which was introduced by Dujardin and Favre in [12]. We refer to their paper and to the seminal paper of Branner and Hubbard [5] for a better description of \mathcal{P}_d .

For every $(c, a) := (c_1, c_2, \dots, c_{d-2}, a) \in \mathbf{C}^{d-1}$, we denote by $P_{c,a}$ the polynomial of degree d whose critical points are $(0, c_1, \dots, c_{d-2})$ and such that $P_{c,a}(0) = a^d$. This polynomial is explicitly given by

$$P_{c,a} := \frac{1}{d}z^d + \sum_2^{d-1} \frac{(-1)^{d-j}}{j} \sigma_{d-j}(c)z^j + a^d,$$

where $\sigma_i(c)$ is the symmetric polynomial of degree i in (c_1, \dots, c_{d-2}) . For convenience, we shall set $c_0 := 0$.

We shall thus work within the holomorphic family $(P_{c,a})_{(c,a) \in M}$, where the parameter space M is simply \mathbf{C}^{d-1} . As explained by Milnor [16], it is convenient to consider the projective compactification \mathbf{P}^{d-1} of $\mathbf{C}^{d-1} = M$ and see the sets $\text{Per}_n(w)$ as algebraic hypersurfaces of \mathbf{P}^{d-1} . We shall denote the projective space at infinity $\{[c : a : 0]; (c, a) \in \mathbf{C}^{d-1} \setminus \{0\}\}$ by \mathbf{P}_∞ .

4.2. The behavior of the bifurcation locus at infinity

We aim to show that the bifurcation locus of the family $\{P_{c,a}\}_{(c,a) \in \mathbf{C}^{d-1}}$ can only cluster on certain hypersurfaces of \mathbf{P}_∞ . The ideas here are essentially those used by Branner and Hubbard for proving the compactness of the connectedness locus (see [5, Chapter 1, Section 3]), but we also borrow from Dujardin and Favre [12].

For every $0 \leq i \leq d-2$, we will denote by α_i the homogeneous polynomial defined by

$$\alpha_i(c, a) := P_{c,a}(c_i) = \frac{1}{d}c_i^d + \sum_{j=2}^{d-1} \frac{(-1)^{d-j}}{j} \sigma_{d-j}(c) c_i^j + a^d,$$

and we will consider the hypersurface Γ_i of \mathbf{P}_∞ defined by

$$\Gamma_i := \{[c : a : 0] / \alpha_i(c, a) = 0\}.$$

By a simple degree argument, one sees that $P_{c,a}(0) = P_{c,a}(c_1) = \dots = P_{c,a}(c_{d-2}) = 0$ implies that $c_1 = \dots = c_{d-2} = a = 0$. This observation and Bezout’s theorem lead to the following.

LEMMA 4.1. *The intersection $\Gamma_0 \cap \Gamma_1 \cap \dots \cap \Gamma_{d-2}$ is empty, and $\Gamma_{i_1} \cap \dots \cap \Gamma_{i_k}$ has codimension k in \mathbf{P}_∞ if $0 \leq i_1 < \dots < i_k \leq d-2$.*

We shall denote by \mathcal{P}_i the set of parameters (c, a) for which the critical point c_i of $P_{c,a}$ has a bounded forward orbit (recall that $c_0 = 0$). The announced result can now be stated as follows.

THEOREM 4.2. *For every $0 \leq i \leq d-2$, the cluster set of \mathcal{P}_i in \mathbf{P}_∞ is contained in Γ_i and, in particular, the connectedness locus is compact in \mathbf{C}^{d-1} .*

Since any cycle of attracting basins captures a critical orbit, the above theorem implies that the intersection of \mathbf{P}_∞ with an algebraic subset of the

form $\text{Per}_{m_1}(\eta_1) \cap \dots \cap \text{Per}_{m_k}(\eta_k)$ is contained in some $\Gamma_{i_1} \cap \dots \cap \Gamma_{i_k}$ as soon as the m_i are mutually distinct and the $|\eta_i|$ strictly smaller than 1. Then, using Bezout’s theorem, one gets the following.

COROLLARY 4.3. *If $1 \leq k \leq d-1$, $m_1 < m_2 < \dots < m_k$, and $\sup_{1 \leq i \leq k} |\eta_i| < 1$, then $\text{Per}_{m_1}(\eta_1) \cap \dots \cap \text{Per}_{m_k}(\eta_k)$ is an algebraic subset of codimension k whose intersection with \mathbf{C}^{d-1} is not empty.*

The proof of Theorem 4.2 relies on estimates on the Green function and, more precisely, on the following result which is proved in [12, Section 6.1].

PROPOSITION 4.4. *Let $g_{c,a}(z) := \lim_n d^{-n} \ln^+ |P_{c,a}^n(z)|$ be the Green function of $P_{c,a}$, and let G be the function defined on \mathbf{C}^{d-1} by: $G(c, a) := \max\{g_{c,a}(c_k); 0 \leq k \leq d-2\}$. Let $\delta := (\sum_{k=0}^{d-2} c_k)/(d-1)$. Then the following estimates occur:*

- (1) $\max\{g_{c,a}(z), G(c, a)\} \geq \ln |z - \delta| - \ln 4$;
- (2) $G(c, a) = \ln^+ \max\{|a|, |c_k|\} + O(1)$.

Proof of Theorem 4.2. Let $\|(c, a)\| := \max(|a|, |c_k|)$. We simply have to check that $\alpha_i((c, a)/\|(c, a)\|)$ tends to 0 when $\|(c, a)\|$ tends to $+\infty$ and that $g_{c,a}(c_i)$ stays equal to 0. As $P_{c,a}(c_i) = \alpha_i(c, a)$ and $g_{c,a}(c_i) = 0$, the estimates given by Proposition 4.4 yield

$$\begin{aligned} \ln^+ \|(c, a)\| + O(1) &= \max(dg_{c,a}(c_i), G(c, a)) = \max(g_{c,a} \circ P_{c,a}(c_i), G(c, a)) \\ &\geq \ln \frac{1}{4} |\alpha_i(c, a) - \delta|. \end{aligned}$$

Since α_i is d -homogeneous, we then get for $\|(c, a)\| > 1$

$$(1-d) \ln \|(c, a)\| + O(1) \geq \ln \frac{1}{4} \left| \alpha_i \left(\frac{(c, a)}{\|(c, a)\|} \right) - \frac{\delta}{\|(c, a)\|^d} \right|,$$

and the conclusion follows since $\delta/\|(c, a)\|^d$ tends to 0 when $\|(c, a)\|$ tends to $+\infty$. □

4.3. Proof of the main result

We shall denote by λ the parameter in \mathbf{C}^{d-1} (i.e., $\lambda := (c, a)$), and we will then set

$$\varphi_n(\lambda) := d^{-n} \ln |p_n(\lambda, w)|,$$

where the polynomials $p_n(\lambda, w)$ are those given by Theorem 2.1. We have to show that the sequence $(\varphi_n)_n$ converges to L in L^1_{loc} . When $|w| < 1$, this

has been shown to be true for any holomorphic family of rational maps (see the first assertion of Theorem 3.1), so we assume that $|w| = 1$.

As it has been previously observed, the case $d = 2$ is covered by Proposition 3.4. To give a flavor of the proof when $d \geq 2$, we will first sketch it for $d = 3$.

Sketch of proof for degree 3 polynomials. Let us first treat the problem on a curve $\text{Per}_{k_0}(\eta_0)$ for $|\eta_0| < 1$. We will show that the sequence $\varphi_n(\lambda)$ converges uniformly to L near any stable (in $\text{Per}_{k_0}(\eta_0)$) parameter λ_0 . For this purpose, one desingularizes an irreducible component of $\text{Per}_{k_0}(\eta_0)$ containing λ_0 and thus obtains a 1-dimensional holomorphic family $(P_{\pi(u)})_{u \in M}$. Keeping in mind that the elements of this family are degree 3 polynomials which do admit an attracting basin of period k_0 and using the fact that the connectedness locus in \mathbf{C}^2 is compact, one sees that the family $(P_{\pi(u)})_{u \in M}$ satisfies the assumptions of Proposition 3.4. The associated sequence $L_n(u, w) = \varphi_n(\pi(u))$ converges therefore in L^1_{loc} to L , and this convergence is locally uniform on stable components by Proposition 3.3.

Let us now consider the problem on the full parameter space \mathbf{C}^2 . Since the family $\{P_{c,a}\}_{(c,a) \in \mathbf{C}^2}$ contains hyperbolic parameters, the fourth assertion of Proposition 3.3 shows that the sequence $\varphi_n(\lambda)$ does not converge to $-\infty$. According to Theorem 2.5, it thus suffices to show that any PSH limit value φ of $\varphi_n(\lambda)$ in $L^1_{\text{loc}}(\mathbf{C}^2)$ coincides with L . Let us therefore assume that φ_{n_k} tends to φ in $L^1_{\text{loc}}(\mathbf{C}^2)$.

We first show that $\varphi = L$ on any open subset of the type

$$A_{k_0} := \bigcup_{|\eta| < 1} \text{Per}_{k_0}(\eta).$$

According to the second assertion of Proposition 3.3, it suffices to show that $\varphi = L$ on any stable component Ω of A_{k_0} . By the third assertion of Proposition 3.3, φ_{n_k} actually converges pointwise to φ on Ω . As (by the previous step) $\varphi_n(\lambda)$ converges locally uniformly on the stable components of $\text{Per}_{k_0}(\eta)$, one thus obtains that $\varphi = L$ on Ω .

According to Theorem 4.2, the set of nonhyperbolic parameters in \mathbf{C}^2 can only cluster on a finite subset of \mathbf{P}_∞ . We may therefore foliate \mathbf{C}^2 by parallel complex lines $(T_t)_{t \in \mathbf{C}}$ whose intersection with the set of nonhyperbolic parameters is compact. After taking a subsequence, we may assume that φ_{n_k} converges to φ in $L^1_{\text{loc}}(T_t)$ for almost every $t \in \mathbf{C}$. To conclude, it remains to

see that $\varphi|_{T_t} \equiv L|_{T_t}$ for these t . For this, one uses Lemma 3.5. The assumptions of the lemma are satisfied since, by construction, the unbounded stable component of T_t is hyperbolic, and the bifurcation locus in T_t is accumulated by sets of the form $T_t \cap A_{k_0}$ where, as we have previously shown, $\varphi = L$. \square

Proof of Theorem 1.2. For $1 \leq q \leq d - 2$, the notation W_q will refer to any irreducible component of a q -codimensional analytic subspace of \mathbf{C}^{d-1} of the form $\text{Per}_{n_1}(\eta_1) \cap \dots \cap \text{Per}_{n_q}(\eta_q)$, where $(\eta_1, \dots, \eta_q) \in \Delta^q$ and where the integers $n_j \geq 2$ are mutually distinct (by Corollary 4.3, such sets do exist). Let us stress that if $\lambda \in W_q$, then the polynomial P_λ admits q distinct attracting basins besides the basin at infinity. Analogously, we shall set $W_0 := \mathbf{C}^{d-1}$. By W_q^{reg} we shall denote the regular part of W_q . The proof will consist in showing by decreasing induction on $0 \leq q \leq d - 2$ that

$(*_q)$: the sequence $\varphi_n|_{W_q}$ tends to L in $L^1_{\text{loc}}(W_q^{\text{reg}})$ for any W_q .

Let us first establish $(*_{d-2})$. The analytic set W_{d-2} is a curve in \mathbf{C}^{d-1} . Desingularizing, we get a proper holomorphic map $\pi : M \rightarrow W_{d-2}$, where M is a Riemann surface. We claim that the 1-dimensional holomorphic family $(P_{\pi(u)})_{u \in M}$ satisfies the assumptions of Proposition 3.4. To see this, we observe that there exists at most one critical point of the polynomial $P_{\pi(u)}$ whose orbit is not captured by one of the $d - 2$ distinct attracting basins of $P_{\pi(u)}$. Let us denote by $c(u)$ this critical point. Assume that u_0 lies in the bifurcation locus of $(P_{\pi(u)})_{u \in M}$. Since all critical points of $P_{\pi(u)}$, except a priori $c(u)$, stay in some attracting basin for u close to u_0 , the orbit of $c(u)$ cannot be uniformly bounded on a small neighborhood of u_0 . This implies that $c(u)$ must belong to the basin of infinity for a convenient small perturbation of u_0 , and shows that $P_{\pi(u_0)}$ becomes hyperbolic after a convenient small perturbation. In other words, the bifurcation locus of $(P_{\pi(u)})_{u \in M}$ is accumulated by hyperbolic parameters.

The above argument also shows that if $P_{\pi(u)}$ is nonhyperbolic, then $c(u)$ cannot belong to the basin at infinity, and therefore $P_{\pi(u)}$ belongs to the connectedness locus. Using the compactness of the connectedness locus and the properness of the map π , one sees that the set of nonhyperbolic parameters of M is compact.

By Proposition 3.4, $\varphi_n(\pi(u))$ converges in $L^1_{\text{loc}}(M)$ to $L \circ \pi$. By the third assertion of Proposition 3.3, the convergence is actually pointwise on the stable components of M , and thus φ_n converges pointwise to L on the stable

set of W_q^{reg} . By the second assertion of Proposition 3.3, $\varphi_n|_{W_q}$ converges to L in $L^1_{\text{loc}}(W_q^{\text{reg}})$. We have proved $(*_{d-2})$. \square

Assuming now that $(*_{q+1})$ is satisfied, we shall prove that $(*_q)$ is true. Let us fix an irreducible q -codimensional analytic set $W_q \subset \text{Per}_{n_1}(\eta_1) \cap \dots \cap \text{Per}_{n_q}(\eta_q)$.

One easily deduces from Corollary 4.3 that W_q contains hyperbolic parameters and that this fact prevents $\varphi_n|_{W_q^{\text{reg}}}$ from converging to $-\infty$ (see Proposition 3.3). According to Theorem 2.5, we thus have to show that for any subsequence $\varphi_{n_k}|_{W_q^{\text{reg}}}$ which converges to some PSH function φ in $L^1_{\text{loc}}(W_q^{\text{reg}})$, one actually has $\varphi = L|_{W_q^{\text{reg}}}$.

We shall use the following two facts which will be proved later.

FACT 1. Let A_m be an open subset of \mathbf{C}^{d-1} defined by

$$A_m := \bigcup_{|\eta| < 1} \text{Per}_m(\eta),$$

where $m > \max(n_1, \dots, n_q)$. If $W_q^{\text{reg}} \cap A_m$ is not empty, then $\varphi = L$ on $W_q^{\text{reg}} \cap A_m$.

FACT 2. There exists a foliation $\bigcup_{t \in A} T_t$ of \mathbf{C}^{d-1} by $(q + 1)$ -dimensional parallel affine subspaces such that, for almost every $t \in A$, the slices $T_t \cap W_q$ are curves on which the set of nonhyperbolic parameters is relatively compact.

Let us consider the curves $T_t \cap W_q$ which are given by Fact 2. By standard arguments, φ_{n_k} converges to φ in L^1_{loc} on almost all these curves and it thus remains to show that $\varphi = L$ on them. For this purpose, we consider an irreducible component Γ of $T_t \cap W_q$ and desingularize it. This yields a proper holomorphic map $\pi : M \rightarrow \Gamma$, where M is a Riemann surface. We shall reach the conclusion by applying Lemma 3.5 to the family $(P_{\pi(u)})_{u \in M}$.

By the properness of π and Fact 2, the set of nonhyperbolic parameters in M is compact, and therefore any nonrelatively compact stable component in M is hyperbolic. Then, by the fourth assertion of Proposition 3.3, $\varphi \circ \pi = L \circ \pi$ on such components.

Using Fact 1, we shall now prove that the bifurcation locus of $(P_{\pi(u)})_{u \in M}$ is accumulated by parameters, where $\varphi \circ \pi = L \circ \pi$. Let u_0 be a point in the bifurcation locus. We may assume that π is locally biholomorphic at u_0 , and it thus suffices to accumulate $\pi(u_0)$ by points where $\varphi = L$. As it is well known, u_0 is accumulated by parameters u_k such that $P_{\pi(u_k)} \in$

$\text{Per}_{m_k}(0)$ and $m_k \rightarrow +\infty$ (this follows also from the general fact that $T_{\text{bif}} = \lim_m d^{-m}[\text{Per}_m(0)]$). This implies that $\pi(u_0)$ is accumulated by open sets of the form $W_q \cap A_{m_k}$. It then follows from Fact 1 that $\pi(u_0)$ is accumulated by points λ_k for which $\varphi(\lambda_k) = L(\lambda_k)$. This ends the proof. \square

Let us finally establish the facts.

Fact 1. Let Ω be a stable component of $W_q^{\text{reg}} \cap A_m$. According to the first and third assertions of Proposition 3.3, the sequence $\varphi_{n_k} - L$ converges locally uniformly to the pluriharmonic negative function $\varphi - L$ on Ω (as previously observed, W_q contains hyperbolic parameters and therefore has no persistent neutral cycles). For all but a finite number of $\eta \in \Delta$, the analytic set $W_q \cap \text{Per}_m(\eta)$ is of codimension $q + 1$ (otherwise W_q would be contained in infinitely many hypersurfaces $\text{Per}_m(\eta)$ and P_λ would have an infinite number of attracting basins when $\lambda \in W_q$). Let us thus pick $\eta_0 \in \Delta$ and $\lambda_0 \in \Omega \cap \text{Per}_m(\eta_0)$ such that $W_q \cap \text{Per}_m(\eta_0)$ has codimension $q + 1$ and is regular at λ_0 . Let us denote by W_{q+1} the irreducible component of $W_q \cap \text{Per}_m(\eta_0)$ to which belongs λ_0 . Then by construction, λ_0 belongs to some stable component ω of W_{q+1}^{reg} . Combining the induction assumption ($*_{q+1}$) with the third assertion of Proposition 3.3, one sees that $\varphi - L = 0$ on ω . In particular, $\varphi(\lambda_0) - L(\lambda_0) = 0$ and, by the maximum principle, $\varphi - L = 0$ on Ω .

It now follows from the second assertion of Proposition 3.3 that $\varphi = L$ on $W_q^{\text{reg}} \cap A_m$. Fact 1 is proved. \square

Fact 2. Let \widetilde{W}_q be the algebraic subset of \mathbf{P}^{d-1} such that $\widetilde{W}_q \cap \mathbf{C}^{d-1} = W_q$. When $q > 0$ and $\lambda \in W_q$, then P_λ has q distinct attracting basins, and therefore at least q of its critical points have a bounded orbit. According to Theorem 4.2, we thus have

$$\widetilde{W}_q \cap \mathbf{P}_\infty \subset \bigcup_{0 \leq i_1 < \dots < i_q \leq d-2} \Gamma_{i_1} \cap \dots \cap \Gamma_{i_q},$$

and moreover, $\bigcup_{0 \leq i_1 < \dots < i_{q+1} \leq d-2} \Gamma_{i_1} \cap \dots \cap \Gamma_{i_{q+1}}$ is a $(d - 3 - q)$ -dimensional algebraic subset of \mathbf{P}_∞ . Thus, as it is classical (see [7, subchapter 7.3]), we may pick a q -dimensional complex plane C_∞ in \mathbf{P}_∞ (a point when $q = 0$) such that

$$C_\infty \cap \left(\bigcup_{0 \leq i_1 < \dots < i_{q+1} \leq d-2} \Gamma_{i_1} \cap \dots \cap \Gamma_{i_{q+1}} \right) = \emptyset.$$

We now slice \mathbf{C}^{d-1} by $(q+1)$ -dimensional parallel affine subspaces T_t which cluster on C_∞ in \mathbf{P}^{d-1} , and we write $\mathbf{C}^{d-1} = \bigcup_{t \in A} T_t$, where A is a $(d-q-2)$ -dimensional complex plane which is transverse to the foliation.

If $\lambda \in W_q$, then at least q of the critical points of P_λ belong to some attracting basin. This implies that the set of nonhyperbolic parameters in $W_q \cap T_t$ may only cluster on the intersection of C_∞ with $\bigcup_{0 \leq i_1 < \dots < i_{q+1} \leq d-2} \Gamma_{i_1} \cap \dots \cap \Gamma_{i_{q+1}}$. The choice of C_∞ guarantees therefore that, for all $t \in A$, the set of nonhyperbolic parameters in $W_q \cap T_t$ is compact.

It remains to show that, for almost all $t \in A$, the analytic set $W_q \cap T_t$ is a curve. For this purpose, let us denote by $\sigma : W_q \rightarrow A$ the canonical projection from W_q onto A . The fiber of $\sigma^{-1}(a)$ has dimension greater than $(d-1) - \dim A - q = 1$. Then, the set of points $a \in A$ for which the fiber $\sigma^{-1}\{a\}$ is of dimension strictly greater than 1 is contained in a countable union of analytic subsets of A whose dimensions are smaller than $\dim W_q - 2 = (d-1) - q - 2 = \dim A - 1$ (see [7, subchapter 3.8]). It is therefore Lebesgue negligible. In other words, $W_q \cap T_t$ is a curve for almost all t , and Fact 2 is proved. \square

REFERENCES

- [1] G. Bassanelli and F. Berteloot, *Bifurcation currents in holomorphic dynamics on \mathbf{P}^k* , *J. Reine Angew. Math.* **608** (2007), 201–235.
- [2] ———, *Lyapunov exponents, bifurcation currents, and laminations in bifurcation loci*, *Math. Ann.* **345** (2009), 1–23.
- [3] F. Berteloot, *Lyapunov exponent of a rational map and multipliers of repelling cycles*, to appear in *Riv. Mat. Univ. Parma*.
- [4] F. Berteloot, C. Dupont, and L. Molino, *Normalization of random families of holomorphic contractions and applications to dynamics*, *Ann. Inst. Fourier (Grenoble)* **58** (2008), 2137–2168.
- [5] B. Branner and J. H. Hubbard, *The iteration of cubic polynomials, I: The global topology of parameter space*, *Acta Math.* **160** (1988), 143–206.
- [6] X. Buff and A. Epstein, “Bifurcation measure and postcritically finite rational maps” in *Complex Dynamics: Families and Friends*, A. K. Peters, Wellesley, Mass., 2009, 491–512.
- [7] E. M. Chirka, *Complex Analytic Sets*, Kluwer Academic Publishers, 1989.
- [8] L. DeMarco, *Dynamics of rational maps: A current on the bifurcation locus*, *Math. Res. Lett.* **8** (2001), 57–66.
- [9] ———, *Dynamics of rational maps: Lyapunov exponents, bifurcations, and capacity*, *Math. Ann.* **326** (2003), 43–73.
- [10] T. C. Dinh and N. Sibony, *Dynamics in several complex variables: Endomorphisms of projective spaces and polynomial-like mappings*, preprint, arXiv:0810.0811
- [11] R. Dujardin, “Cubic polynomials: A measurable view on the parameter space” in *Complex Dynamics: Families and Friends*, A. K. Peters, Wellesley, Mass., 2009, 451–489.

- [12] R. Dujardin and C. Favre, *Distribution of rational maps with a preperiodic critical point*, Amer. J. Math. **130** (2008), 979–1032.
- [13] L. Hörmander, *The Analysis of Linear Partial Differential Operators, I*, Springer, New York, 1983.
- [14] R. Mañé, “The Hausdorff dimension of invariant probabilities of rational maps” in *Dynamical Systems (Valparaiso, Chile, 1986)*, Lect. Notes in Math. **1331** (1988), 86–117.
- [15] R. Mañé, P. Sad, and D. Sullivan, *On the dynamics of rational maps*, Ann. Sci. Éc. Norm. Supér. (4) (1983), 193–217.
- [16] J. W. Milnor, *Geometry and dynamics of quadratic rational maps*, Experiment. Math. **2** (1993), 37–83.
- [17] J. H. Silverman, *The Arithmetic of Dynamical Systems*, Grad. Texts in Math. **241**, Springer, New York, 2007.

Giovanni Bassanelli
(deceased)

François Berteloot
Université Paul Sabatier MIG
Institut de Mathématiques de Toulouse
31062 Toulouse Cedex 9
France
`berteloo@picard.ups-tlse.fr`