The contribution of interactions to arthropod community structure depends on the place and time: Why community ecologists should think locally and regionally


Matthew K. Lau, Laura E. Hagenauer and Thomas G. Whitham

Department of Biological Sciences Northern Arizona University

• Disintegrating the community (Ricklefs 2008)


- Disintegrating the community (Ricklefs 2008)
 - Brooks et al. 2009

- Disintegrating the community (Ricklefs 2008)
 - Brooks et al. 2009



Regional Scale

- Disintegrating the community (Ricklefs 2008)
 - Brooks et al. 2009







Local Scale

Regional Scale

- Disintegrating the community (Ricklefs 2008)
 - Brooks et al. 2009
- Local community structure can be affected by local interactions, context dependent

- Disintegrating the community (Ricklefs 2008)
 - Brooks et al. 2009
- Local community structure can be affected by local interactions, context dependent
- What does this mean for the study of communities?








aphid-exclusion ramets percent aphid galls attacked on the second days and the second days with inquilines 15-10inquilines fungi


Dickson and Whitham 1996


















How important are interactions of canopy arthropod species?









Three stands of cottonwood



- Three stands of cottonwood
 - Populus fremontii




- Three stands of cottonwood
 - Populus fremontii
 - Planting dates (2000, 2002, 2005)



- Three stands of cottonwood
 - Populus fremontii
 - Planting dates (2000, 2002, 2005)
 - ~650 trees





- Three stands of cottonwood
 - Populus fremontii
 - Planting dates (2000, 2002, 2005)
 - ~650 trees






- Three stands of cottonwood
 - Populus fremontii
 - Planting dates (2000, 2002, 2005)
 - ~650 trees









- Three stands of cottonwood
 - Populus fremontii
 - Planting dates (2000, 2002, 2005)
 - ~650 trees









- Three stands of cottonwood
 - Populus fremontii
 - Planting dates (2000, 2002, 2005)
 - ~650 trees
- Sampling (n = 15)








- Three stands of cottonwood
 - Populus fremontii
 - Planting dates (2000, 2002, 2005)
 - ~650 trees
- Sampling (n = 15)
 - Sampled monthly in 2006












- Three stands of cottonwood
 - Populus fremontii
 - Planting dates (2000, 2002, 2005)
 - ~650 trees
- Sampling (n = 15)
 - Sampled monthly in 2006
 - Timed-observation sampling












- Three stands of cottonwood
 - Populus fremontii
 - Planting dates (2000, 2002, 2005)
 - ~650 trees
- Sampling (n = 15)
 - Sampled monthly in 2006
 - Timed-observation sampling
 - Juvenile foliage







Occurrence Matrix

Species	Obs I	Obs2	Obs3
Spl	I	I	0
Sp2	0	0	I
Sp3	0		I

Occurrence Matrix

Species	Obs I	Obs2	Obs3
Spl	I	I	0
Sp2	0	0	I
Sp3	0	I	I

Occurrence Matrix

Species	Obs I	Obs2	Obs3
SpI	I	П	0
Sp2	0	0	I
Sp3	0		




Observed
Co-occurrence
Index

Occurrence Matrix

Species	Obs I	Obs2	Obs3
SpI	I	I	0
Sp2	0	0	I
Sp3	0		I



Observed
Co-occurrence
Index




Occurrence Matrix

Species	Obs I	Obs2	Obs3
SpI	I	I	0
Sp2	0	0	I
Sp3	0		I





Observed
Co-occurrence
Index




Occurrence Matrix


Species	ObsI	Obs2	Obs3
Spl	I	I	0
Sp2	0	0	I
Sp3	0	I	I








Occurrence Matrix



































• C-score (Stone and Roberts 1990)

- C-score (Stone and Roberts 1990)
 - Measures average degree of NON-co-occurrence

- C-score (Stone and Roberts 1990)
 - Measures average degree of NON-co-occurrence
- Fixed-equiprobable algorithm (Gotelli 2000)

- C-score (Stone and Roberts 1990)
 - Measures average degree of NON-co-occurrence
- Fixed-equiprobable algorithm (Gotelli 2000)
- Standardized effect size (SES)


- C-score (Stone and Roberts 1990)
 - Measures average degree of NON-co-occurrence
- Fixed-equiprobable algorithm (Gotelli 2000)
- Standardized effect size (SES)

SES


- C-score (Stone and Roberts 1990)
 - Measures average degree of NON-co-occurrence
- Fixed-equiprobable algorithm (Gotelli 2000)
- Standardized effect size (SES)


SES 0 Random


- C-score (Stone and Roberts 1990)
 - Measures average degree of NON-co-occurrence
- Fixed-equiprobable algorithm (Gotelli 2000)
- Standardized effect size (SES)





- C-score (Stone and Roberts 1990)
 - Measures average degree of NON-co-occurrence
- Fixed-equiprobable algorithm (Gotelli 2000)
- Standardized effect size (SES)





























Results Month 3 June Standardized Effect Size (SES) August 2 October * 0 * -2 *** -3 *** 4 years I year 6 years STAND AGE P-value: * < 0.10, *** < 0.01







 Richness and diversity consistent from stand to stand and month to month

- Richness and diversity consistent from stand to stand and month to month
- Community composition varied by month and stand


- Richness and diversity consistent from stand to stand and month to month
- Community composition varied by month and stand

PerMANOVA: month*stand P < 0.001 ***

October


Stand Age

- I year
- 4 years
- 6 years




Axis I

PerMANOVA: month*stand P < 0.001 ***



PerMANOVA: month*stand P < 0.001 ***



2005 (I year)

PerMANOVA: month*stand P < 0.001 ***

Month

- June
- August
- October

2005 (I year) 2002 (4 years)

Month

June


August

October


PerMANOVA: month*stand P < 0.001 ***







PerMANOVA: month*stand P < 0.001 ***



Month

PerMANOVA: month*stand P < 0.001 ***

• Local interactions can play a role in producing local community structure


- Local interactions can play a role in producing local community structure
- Forces vary by month and by stand


- Local interactions can play a role in producing local community structure
- Forces vary by month and by stand
 - Could result from stand-environmental or compositional differences

• Understanding local processes is important


- Understanding local processes is important
- Regional process are also important

- Understanding local processes is important
- Regional process are also important
- Focus on the interplay of ecological and evolutionary process across spatial and temporal scales







Local Scale



Local Scale





Local Scale




Local Scale








Local Scale



"Pattern is neither a property of the system alone nor of the observer, but an interaction between them."

Simon Levin 1989

Local Scale



<u>Acknowledgements</u>

Todd Wojtowicz*

Jamie Lamit



NAU IGERT Program





Analytical Software

EcoSim (Aquired Intelligence and Kesey-Bear, Inc.)







statistical computing language

