

RESEARCH

Open Access

Limit theorems for delayed sums of random sequence

Ding Fang-qing¹ and Wang Zhong-zhi^{2*}* Correspondence: wzz30@ahut.edu.cn²Faculty of Mathematics & Physics, Anhui University of Technology, Ma'anshan 243002, P. R. China
Full list of author information is available at the end of the article**Abstract**

For a sequence of arbitrarily dependent random variables $(X_n)_{n \in \mathbf{N}}$ and Borel sets $(B_n)_{n \in \mathbf{N}}$, on real line the strong limit theorems, represented by inequalities, i.e. the strong deviation theorems of the delayed average $S_{n,k_n}(\omega)$ are investigated by using the notion of asymptotic delayed log-likelihood ratio. The results obtained popularizes the methods proposed by Liu.

Mathematics Subject Classification 2000: Primary, 60F15.**Keywords:** strong deviation theorem, likelihood ratio, delayed sums**1. Introduction**

Let $(a_n)_{n \in \mathbf{N}}$ be a sequence of real numbers and let $(k_n)_{n \in \mathbf{N}}$ be a sequence of positive integers. The numbers

$$\rho_{n,k_n} = \left\{ \sum_{j=1}^{k_n} a_{n+j-1} \right\} / k_n$$

are called the (forward) delayed first arithmetic means (See [1]). In [2], using the limiting behavior of delayed average, Chow found necessary and sufficient conditions for the Borel summability of i.i.d. random variables and also obtained very simple proofs of a number of well-known results such as the Hsu-Robbins-Spitzer-Katz theorem. In [3], Lai studied the analogues of the law of the iterated logarithm for delayed sums of independent random variables. Recently, Chen [4] has presented an accurate description the limiting behavior of delayed sums under a non-identically distribution setup, and has deduced Chover-type laws of the iterated logarithm for them.

Our aim in this article is to establish strong deviation theorems (limit theorem expressed by inequalities, see [5]) of delayed average for the dependent absolutely continuous random variables. By using the notion of asymptotic delayed log-likelihood ratio, we extend the analytic technique proposed by Liu [5] to the case of delayed sums. The crucial part of the proof is to construct a delayed likelihood ratio depending on a parameter, and then applies the Borel-Cantelli lemma.

Throughout, let $(X_n)_{n \in \mathbf{N}}$ be a sequence of absolutely continuous random variables on a fixed probability space $\{\Omega, \mathcal{F}, P\}$ with the joint density function $g^{1, n}(x_1, \dots, x_n)$, $n \in \mathbf{N}$, and $f_j(x)$, $j = 1, 2, \dots$ be the the marginal density function of random variable X_j . (k_n)

$n \in \mathbf{N}$ be a subsequence of positive integers, such that, for every $\varepsilon > 0$, $\sum_{n=1}^{\infty} \exp(-k_n \varepsilon) < \infty$.

Definition 1. The delayed likelihood ratio is defined by

$$\mathcal{L}_n(\omega) = \begin{cases} \frac{\prod_{j=n}^{n+k_n-1} f_j(X_j)}{g^{n,n+k_n-1}(x_n, \dots, x_{n+k_n-1})}, & \text{if denominator} > 0 \\ 0, & \text{otherwise} \end{cases} \quad (1.1)$$

Let

$$\mathcal{L}(\omega) = -\liminf_n \frac{1}{k_n} \log \mathcal{L}_n(\omega) \quad (1.2)$$

$\mathcal{L}(\omega)$ is called asymptotic delayed log-likelihood ratio, where $g^{n,n+k_n-1}(x_n, \dots, x_{n+k_n-1})$ denotes the joint density function of random vector $(X_n, \dots, X_{n+k_n-1})$, ω is a sample point (with $\log 0 = -\infty$).

It will be shown in Lemma 1 that $\mathcal{L}(\omega) \geq 0$ a.e. in any case.

Remark 1. It will be seen below that $\mathcal{L}(\omega)$ has the analogous properties of the likelihood ratio in [5], Although $\mathcal{L}(\omega)$ is not a proper metric among probability measures, we nevertheless consider it as a measure of “discrimination” between the dependence (their joint distribution) and independence (the product of their marginals). Obviously, $\mathcal{L}_n(\omega) = 1$, a.e. $n \in \mathbf{N}$ if $(X_n)_{n \in \mathbf{N}}$ is independent. In view of the above discussion of the asymptotic logarithmic delayed likelihood ratio, it is natural for us to think of $\mathcal{L}(\omega)$ as a measure how far (the random deviation) of $(X_n)_{n \in \mathbf{N}}$ is from being independent and how dependent they are. The closer $\mathcal{L}(\omega)$ approaches to 0, the smaller the deviation is.

Lemma 1. Let $\mathcal{L}_n(\omega)$ be define as above, then

$$\limsup_n \frac{1}{k_n} \log \mathcal{L}_n(\omega) \leq 0, \text{ a.e.} \quad (1.3)$$

Proof. Let $B = \{(x_n, \dots, x_{n+k_n-1}) : g^{n,n+k_n-1}(x_n, \dots, x_{n+k_n-1}) > 0\}$. Since

$$\begin{aligned} & E[\mathcal{L}_n(\omega)] \\ &= \int \cdots \int_{(x_n, \dots, x_{n+k_n-1}) \in B} \frac{\prod_{j=n}^{n+k_n-1} f_j(x_j)}{g^{n,n+k_n-1}(x_n, \dots, x_{n+k_n-1})} \\ & \quad \cdot g^{n,n+k_n-1}(x_n, \dots, x_{n+k_n-1}) dx_n \cdots dx_{n+k_n-1} \\ &= \int \cdots \int_{(x_n, \dots, x_{n+k_n-1}) \in B} \prod_{j=n}^{n+k_n-1} f_j(x_j) dx_n \cdots dx_{n+k_n-1} \\ &\leq \int \cdots \int_{(x_n, \dots, x_{n+k_n-1}) \in \mathbf{R}^{k_n}} \prod_{j=n}^{n+k_n-1} f_j(x_j) dx_n \cdots dx_{n+k_n-1} = 1. \end{aligned}$$

From Markov inequality, for every $\varepsilon > 0$, we have

$$P\left[\frac{1}{k_n} \log \mathcal{L}_n(\omega) \geq \varepsilon\right] = P[\mathcal{L}_n(\omega) \geq \exp(k_n \varepsilon)] \leq 1 \cdot \exp(-k_n \varepsilon).$$

Hence

$$\sum_{n=1}^{\infty} P \left[\frac{1}{k_n} \log \mathcal{L}_n(\omega) \geq \varepsilon \right] \leq \sum_{n=1}^{\infty} \exp(-k_n \varepsilon) < \infty.$$

By Borel-Cantelli lemma, we have

$$P \left[\limsup_n \frac{1}{k_n} \log \mathcal{L}_n(\omega) \geq 2\varepsilon \right] = 0,$$

for any $\varepsilon > 0$, (1.3) follows immediately. \square

2. Main results and proofs

Theorem 1. Let $(X_n)_{n \in \mathbb{N}}$, $\mathcal{L}_n(\omega)$, $\mathcal{L}(\omega)$ be defined as above, $(B_n)_{n \in \mathbb{N}}$ be a sequence of Borel sets of the real line. Let $S_{n,k_n}(\omega) = \frac{1}{k_n} \sum_{j=n}^{n+k_n-1} \mathbf{1}_{B_j}(X_j)$, and assume

$$c = \limsup_n \frac{1}{k_n} \sum_{j=n}^{n+k_n-1} P(X_j \in B_j), \tag{2.1}$$

then

$$\limsup_n S_{n,k_n}(\omega) \leq (\sqrt{\mathcal{L}(\omega)} + \sqrt{c})^2, \text{ a.e.} \tag{2.2}$$

where $\mathbf{1}_{B_n}(\cdot)$ be the indicator function of B_n .

Proof. Assume $s > 0$ to be a constant, and let

$$h_j(x_j) = \frac{s^{\mathbf{1}_{B_j}(x_j)} f_j(x_j)}{1 + (s-1) \int_{B_j} f_j(x_j) dx_j}, \quad j = 1, 2, \dots \tag{2.3}$$

It is not difficult to see that $\int h_j(x_j) dx_j = 1$, $j = 1, 2, \dots$. Let

$$\Lambda_n(s, \omega) = \begin{cases} \frac{\prod_{j=n}^{n+k_n-1} h_j(X_j)}{s^{n, n+k_n-1}(X_n, \dots, X_{n+k_n-1})}, & \text{if denominator} > 0 \\ 0, & \text{otherwise} \end{cases} \tag{2.4}$$

From Lemma 1, there exists $A(s) \in \mathcal{F}$, $P(A(s)) = 1$, such that

$$\limsup_n \frac{1}{k_n} \log \Lambda_n(s, \omega) \leq 0, \quad \omega \in A(s) \tag{2.5}$$

Since $\int_{B_j} f_j(x_j) dx_j = P(X_j \in B_j)$, by (2.3) we have

$$\begin{aligned} & \prod_{j=n}^{n+k_n-1} h_j(x_j) \\ &= \prod_{j=n}^{n+k_n-1} \frac{s^{\mathbf{1}_{B_j}(x_j)} f_j(x_j)}{1 + (s-1) \int_{B_j} f_j(x_j) dx_j} \\ &= s^{\sum_{j=n}^{n+k_n-1} \mathbf{1}_{B_j}(x_j)} \prod_{j=n}^{n+k_n-1} \frac{f_j(x_j)}{1 + (s-1) P(X_j \in B_j)} \end{aligned} \tag{2.6}$$

It follows from (1.1), (2.4) and (2.6) that

$$\log \Lambda_n(s, \omega) = \sum_{j=n}^{n+k_n-1} \mathbf{1}_{B_j}(X_j) \log s - \sum_{j=n}^{n+k_n-1} \log [1 + (s-1)P(X_j \in B_j)] + \log \mathcal{L}_n(\omega) \quad (2.7)$$

(2.5) and (2.7) yield

$$\limsup_n \frac{1}{k_n} \left(\log s \sum_{j=n}^{n+k_n-1} \mathbf{1}_{B_j}(X_j) - \sum_{j=n}^{n+k_n-1} \log [1 + (s-1)P(X_j \in B_j)] + \log \mathcal{L}_n(\omega) \right) \leq 0, \quad \omega \in A(s) \quad (2.8)$$

Let $s > 1$, dividing the two sides of (2.8) by $\log s$, we have

$$\limsup_n \frac{1}{k_n} \left(\sum_{j=n}^{n+k_n-1} \mathbf{1}_{B_j}(X_j) - \sum_{j=n}^{n+k_n-1} \frac{\log [1 + (s-1)P(X_j \in B_j)]}{\log s} + \frac{\log \mathcal{L}_n(\omega)}{\log s} \right) \leq 0, \quad \omega \in A(s) \quad (2.9)$$

By (1.2), (2.9) and the property $\limsup_n(a_n - b_n) \leq d \Rightarrow \limsup_n(a_n - c_n) \leq \limsup_n(b_n - c_n) + d$, one gets

$$\limsup_n \frac{1}{k_n} \left(\sum_{j=n}^{n+k_n-1} \mathbf{1}_{B_j}(X_j) - \sum_{j=n}^{n+k_n-1} \frac{\log [1 + (s-1)P(X_j \in B_j)]}{\log s} \right) \leq \frac{\mathcal{L}(\omega)}{\log s}, \quad \omega \in A(s) \quad (2.10)$$

By (2.10) and the property of the superior above and the inequality $0 < \log(1+x) \leq x$ ($x > 0$), we obtain

$$\begin{aligned} & \limsup_n S_{n,k_n}(\omega) \\ & \leq \limsup_n \frac{1}{k_n} \left(\sum_{j=n}^{n+k_n-1} \frac{\log [1 + (s-1)P(X_j \in B_j)]}{\log s} \right) + \frac{\mathcal{L}(\omega)}{\log s} \\ & \leq \limsup_n \frac{1}{k_n} \left(\sum_{j=n}^{n+k_n-1} \frac{(s-1)P(X_j \in B_j)}{\log s} \right) + \frac{\mathcal{L}(\omega)}{\log s} \\ & \leq c \left(\frac{s-1}{\log s} \right) + \frac{\mathcal{L}(\omega)}{\log s}, \quad \omega \in A(s) \end{aligned} \quad (2.11)$$

(2.11) and the inequality $1 - \frac{1}{s} < \log s$ ($s > 1$) imply

$$\limsup_n S_{n,k_n}(\omega) \leq c \cdot s + \frac{s\mathcal{L}(\omega)}{s-1}, \quad \omega \in A(s) \quad (2.12)$$

Let D be a set of countable real numbers dense in the interval $(1, +\infty)$, and let $A^* = \bigcap_{s \in D} A(s)$, $g(s, x) = cs + sx/(s-1)$, then we have by (2.12)

$$\limsup_n S_{n,k_n}(\omega) \leq g(s, \mathcal{L}(\omega)), \quad \omega \in A^*, \quad s \in D \quad (2.13)$$

Let $c > 0$, it easy to see that if $\mathcal{L}(\omega) > 0$, *a.e.*, then, for fixed ω , $g(s, \mathcal{L}(\omega))$ as a function of s attains its smallest value $g\left(1 + \sqrt{\mathcal{L}(\omega)/c}, \mathcal{L}(\omega)\right) = 2\sqrt{c\mathcal{L}(\omega)} + \mathcal{L}(\omega) + c$ on the interval $(1, +\infty)$, and $g(s, 0)$ is increasing on the interval $(1, +\infty)$ and $\lim_{s \rightarrow 1^+} g(s, 0) = 0$. For each $\omega \in A^* \cap A(1)$, if $\mathcal{L}(\omega) \neq \infty$, take $\kappa_n(\omega) \in D$, $n = 1, 2, \dots$, such that $\kappa_n(\omega) \rightarrow 1 + \sqrt{\mathcal{L}(\omega)/c}$. We have by the continuity of $g(s, \mathcal{L}(\omega))$ with respect to s ,

$$\lim_{n \rightarrow +\infty} g(\kappa_n(\omega), \mathcal{L}(\omega)) = (\sqrt{\mathcal{L}(\omega)} + \sqrt{c})^2, \quad (2.14)$$

By (2.13), we obtain

$$\limsup_n S_{n,k_n} \leq g(\kappa_n(\omega), \mathcal{L}(\omega)), \quad n = 1, 2, \dots \quad (2.15)$$

(2.14) and (2.15) imply

$$\limsup_n S_{n,k_n}(\omega) \leq (\sqrt{\mathcal{L}(\omega)} + \sqrt{c})^2, \quad \omega \in A^* \cap A(1) \quad (2.16)$$

If $\mathcal{L}(\omega) = \infty$, (2.16) holds trivially. Since $P(A^* \cap A(1)) = 1$, (2.2) holds by (2.16), when $c > 0$.

When $c = 0$, we have by letting $s = e$ in (2.11),

$$\limsup_n S_{n,k_n}(\omega) \leq \mathcal{L}(\omega), \quad \omega \in A(e) \quad (2.17)$$

since $P(A(e)) = 1$, (2.2) also holds by (2.17) when $c = 0$. \square

Theorem 2. Let $(X_n)_{n \in \mathbb{N}}$, $\mathcal{L}_n(\omega)$, $\mathcal{L}(\omega)$, $(B_n)_{n \in \mathbb{N}}$, $S_{n,k_n}(\omega)$ be defined as in Theorem 1 and assume

$$c' = \liminf_n \frac{1}{k_n} \sum_{j=n}^{n+k_n-1} P(X_j \in B_j), \quad (2.18)$$

then, if $0 \leq \mathcal{L}(\omega) \leq c'$ a.e., then

$$\liminf_n S_{n,k_n}(\omega) \geq c' - 2\sqrt{c'\mathcal{L}(\omega)}, \quad a.e. \quad (2.19)$$

Proof. Let $0 < s < 1$, dividing the two sides of (2.8) by $\log s$, we have

$$\liminf_n \frac{1}{k_n} \left(\sum_{j=n}^{n+k_n-1} \mathbf{1}_{B_j}(X_j) - \sum_{j=n}^{n+k_n-1} \frac{\log[1 + (s-1)P(X_j \in B_j)]}{\log s} + \frac{\log \mathcal{L}_n(\omega)}{\log s} \right) \geq 0, \quad \omega \in A(s) \quad (2.20)$$

By (1.2), (2.20) and the property $\liminf_n (a_n - b_n) \geq d \Rightarrow \liminf_n (a_n - c_n) \geq \liminf_n (b_n - c_n) + d$, one gets

$$\liminf_n \frac{1}{k_n} \left(\sum_{j=n}^{n+k_n-1} \mathbf{1}_{B_j}(X_j) - \sum_{j=n}^{n+k_n-1} \frac{\log[1 + (s-1)P(X_j \in B_j)]}{\log s} \right) \geq \frac{\mathcal{L}(\omega)}{\log s}, \quad \omega \in A(s) \quad (2.21)$$

By (2.21) and the property of the inferior above and the inequality $\log(1+x) \leq x(-1 < x \leq 0)$, we obtain

$$\begin{aligned} & \liminf_n S_{n,k_n}(\omega) \\ & \geq \liminf_n \frac{1}{k_n} \left(\sum_{j=n}^{n+k_n-1} \frac{\log[1 + (s-1)P(X_j \in B_j)]}{\log s} \right) + \frac{\mathcal{L}(\omega)}{\log s} \\ & \geq \liminf_n \frac{1}{k_n} \left(\sum_{j=n}^{n+k_n-1} \frac{(s-1)P(X_j \in B_j)}{\log s} \right) + \frac{\mathcal{L}(\omega)}{\log s} \\ & \geq c' \left(\frac{s-1}{\log s} \right) + \frac{\mathcal{L}(\omega)}{\log s}, \quad \omega \in A(s) \end{aligned} \quad (2.22)$$

(2.22) and the inequality $1 - \frac{1}{s} < \log s$ and $\log s < s - 1$ ($0 < s < 1$) imply

$$\liminf_n S_{n,k_n}(\omega) \geq c' \cdot s + \frac{\mathcal{L}(\omega)}{s-1}, \quad \omega \in A(s) \cap A(1) \quad (2.23)$$

Let D' be a set of countable real numbers dense in the interval $(0, 1)$, and let $A_* = \bigcap_{s \in D'} A(s)$, $h(s, x) = c's + x/(s - 1)$, then we have by (2.23)

$$\liminf_n S_{n,k_n}(\omega) \geq h(s, \mathcal{L}(\omega)), \quad \omega \in A_*, \quad s \in D' \quad (2.24)$$

Let $c' > 0$, it easy to see that if $0 < \mathcal{L}(\omega) < c'$, *a.e.*, then, for fixed ω , $h(s, \mathcal{L}(\omega))$ as a function of s attains its maximum value $h\left(1 - \sqrt{\mathcal{L}(\omega)/c'}, \mathcal{L}(\omega)\right) = c' - 2\sqrt{c'\mathcal{L}(\omega)}$, on the interval $(0, 1)$, and $h(s, 0)$ is increasing on the interval $(0, 1)$ and $\lim_{s \rightarrow 1^+} h(s, 0) = c'$. For each $\omega \in A_* \cap A(1)$, if $\mathcal{L}(\omega) \neq \infty$, take $l_n(\omega) \in D'$, $n = 1, 2, \dots$, such that $l_n(\omega) \rightarrow 1 - \sqrt{\mathcal{L}(\omega)/c'}$. We have by the continuity of $h(s, \mathcal{L}(\omega))$ with respect to s ,

$$\lim_{n \rightarrow +\infty} h(l_n(\omega), \mathcal{L}(\omega)) = c' - 2\sqrt{c'\mathcal{L}(\omega)}, \quad (2.25)$$

By (2.24), we obtain

$$\liminf_n S_{n,k_n} \geq h(l_n(\omega), \mathcal{L}(\omega)), \quad n = 1, 2, \dots \quad (2.26)$$

(2.25) and (2.26) imply

$$\liminf_n S_{n,k_n}(\omega) \geq c' - 2\sqrt{c'\mathcal{L}(\omega)}, \quad \omega \in A_* \cap A(1) \quad (2.27)$$

If $\mathcal{L}(\omega) = \infty$, (2.27) holds trivially. Since $P(A_* \cap A(1)) = 1$, (2.19) holds by (2.27), when $c' > 0$. (2.19) also holds trivially when $c' = 0$. \square

Remark 2. In case $\mathcal{L}(\omega) > c' \geq 0$, *a.e.*, we cannot get a better lower bound of $\liminf_n S_{n,k_n}(\omega)$. This motivates the following problem: under the conditions of Theorem 2, how to get a better lower bound of $\liminf_n S_{n,k_n}(\omega)$ in case of $\mathcal{L}(\omega) > c' \geq 0$, *a.e.*?

Definition 2. (Generalized empirical distribution function) Let $(X_n)_{n \in \mathbf{N}}$ be identically distribution with common distribution function F , for each $m, n \in \mathbf{N}$, let

$$F_{m,n}(x) = \frac{1}{n} \sum_{k=m}^{m+n-1} \mathbf{1}_{(X_k \leq x)}.$$

$F_{m,n}$ is the observed frequency of values that are $\leq x$ from time m to $m + n - 1$. The $F_{1,n}$ is the usual empirical distribution function, hence the name given above.

In particular, let $B = (-\infty, x]$, $x \in R$ in Theorems 1 and 2, we can get a strong limit theorem for the generalized empirical distribution function.

Corollary 1. Let $(X_n)_{n \in \mathbf{N}}$ be *i.i.d.* random variables with common distribution function F , let $B_n = (-\infty, x]$, $n = 1, 2, \dots$, then

$$\lim_n F_{n,n+k_n-1}(x) = F(x), \quad \textit{a.e.}$$

Corollary 2. Let $(X_n)_{n \in \mathbf{N}}$ be independent random variables and $(B_n)_{n \in \mathbf{N}}$ be as Theorem 1, then

$$\lim_n \frac{1}{k_n} \sum_{j=n}^{n+k_n-1} [\mathbf{1}_{B_j}(X_j) - P(X_j \in B_j)] = 0, \text{ a.e.} \quad (2.28)$$

Proof. Note that $P(X_j \in B_j) \leq 1, j = 1, 2, \dots$ and in this case, $0 \leq c, c' \leq 1, \mathcal{L}(\omega) = 0$ a.e., we have by (2.11)

$$\limsup_n \frac{1}{k_n} \left(\sum_{j=n}^{n+k_n-1} \mathbf{1}_{B_j}(X_j) - \sum_{j=n}^{n+k_n-1} \frac{\log[1 + (s-1)P(X_j \in B_j)]}{\log s} \right) \leq 0, \quad \omega \in A(s) \quad (2.29)$$

by (2.29) and the property of the superior above and the inequality $0 \leq \log(1+x) \leq x$ ($x > 0$), we obtain

$$\begin{aligned} & \limsup_n \frac{1}{k_n} \sum_{j=n}^{n+k_n-1} [\mathbf{1}_{B_j}(X_j) - P(X_j \in B_j)] \\ & \leq \limsup_n \frac{1}{k_n} \left(\sum_{j=n}^{n+k_n-1} \frac{\log[1 + (s-1)P(X_j \in B_j)]}{\log s} - P(X_j \in B_j) \right) \\ & \leq \limsup_n \frac{1}{k_n} \left(\sum_{j=n}^{n+k_n-1} \frac{(s-1)P(X_j \in B_j)}{\log s} - P(X_j \in B_j) \right) \\ & \leq \left(\frac{s-1}{\log s} - 1 \right), \quad \omega \in A(s) \end{aligned} \quad (2.30)$$

Analogously as in the proof of Theorem 1, we obtain

$$\limsup_n \frac{1}{k_n} \sum_{j=n}^{n+k_n-1} [\mathbf{1}_{B_j}(X_j) - P(X_j \in B_j)] \leq 0, \text{ a.e.} \quad (2.31)$$

Similarly, we have $\liminf_n \frac{1}{k_n} \sum_{j=n}^{n+k_n-1} [\mathbf{1}_{B_j}(X_j) - P(X_j \in B_j)] \geq 0, \text{ a.e.}$ hence (2.28) follows immediately. \square

Remark 3. Let $B_n = B$, Corollary 2 implies that $\frac{\lim S_{n,k_n}}{k_n} = P(X_1 \in B)$ which gives the strong law of large numbers for the delayed arithmetic means.

Acknowledgements

This work is supported by The National Natural Science Foundation of China (Grant No. 11071104) and the An Hui University of Technology research grant: D2011025. The authors would like to thank two referees for their insightful comments which resulted in improving Theorems 1, 2 and Corollary 2 significantly.

Author details

¹Department of Mathematics & Physics, HeFei University, HeFei 230601, P. R. China ²Faculty of Mathematics & Physics, AnHui University of Technology, Ma'anshan 243002, P. R. China

Authors' contributions

WZ and DF carried out the design of the study and performed the analysis. DF drafted the manuscript. All authors read and approved the final manuscript.

Competing interests

The authors declare that they have no competing interests.

Received: 29 October 2011 Accepted: 31 May 2012 Published: 31 May 2012

References

1. Zygmund, A: *Trigonometric Series* 1. Cambridge University Press, Cambridge (1959)

2. Chow, YS: Delayed sums and Borel summability for independent, identically distributed random variables. *Bull Inst Math Academia Sinica.* **1**, 207–220 (1972)
3. Lai, TL: Limit theorems for delayed sums. *Ann Probab.* **2**(3), 432–440 (1974). doi:10.1214/aop/1176996658
4. Chen, PY: Limiting behavior of delayed sums under a non-identically distribution setup. *Ann Braz Acad Sci.* **80**(4), 617–625 (2008)
5. Liu, W: Strong deviation theorems and analytical method. Academic press, Beijing (2003) (in Chinese)

doi:10.1186/1029-242X-2012-124

Cite this article as: Fang-qing and Zhong-zhi: Limit theorems for delayed sums of random sequence. *Journal of Inequalities and Applications* 2012 **2012**:124.

Submit your manuscript to a SpringerOpen[®] journal and benefit from:

- ▶ Convenient online submission
- ▶ Rigorous peer review
- ▶ Immediate publication on acceptance
- ▶ Open access: articles freely available online
- ▶ High visibility within the field
- ▶ Retaining the copyright to your article

Submit your next manuscript at ▶ springeropen.com
