

Hindawi Publishing Corporation
Evidence-Based Complementary and Alternative Medicine
Volume 2014, Article ID 721341, 2 pages
<http://dx.doi.org/10.1155/2014/721341>



Erratum

Erratum to “PXR-Mediated Upregulation of CYP3A Expression by Herb Compound Praeruptorin C from *Peucedanum praeruptorum* Dunn”

Ling Huang,^{1,2} Qian Wu,² Yu-Hua Li,¹ Yi-Tao Wang,³ and Hui-Chang Bi¹

¹ *Laboratory of Drug Metabolism and Pharmacokinetics, School of Pharmaceutical Sciences, Sun Yat-sen University, 132 Waihuan Dong Road, Guangzhou 510006, China*

² *School of Pharmaceutical Sciences, Hainan Medical University, Haikou 571010, China*

³ *Institute of Chinese Medical Sciences, Macau University, Macau 999078, China*

Correspondence should be addressed to Hui-Chang Bi; bihchang@mail.sysu.edu.cn

Received 25 June 2014; Accepted 20 July 2014; Published 28 October 2014

Copyright © 2014 Ling Huang et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Figure 3(a) was incorrectly published in the original paper titled “PXR-mediated upregulation of CYP3A expression by herb compound praeruptorin C from *Peucedanum praeruptorum* Dunn.” The corrected figure is attached below. The *hPXR* mRNA expression in LS174T cells was remarkably increased to 675-fold after transfection with pSG5-*hPXR* expression plasmid.







FIGURE 3: Effects of PC on the expression of *CYP3A4* mRNA in LS174T cells. (a) LS174T cells were transfected with hPXR expression plasmids for 6 h. Total RNA of LS174T cells was isolated, and *hPXR* mRNA levels were analyzed by real-time PCR. The effect of herbal compounds on *hPXR* mRNA levels is presented as fold mRNA expression to control vehicle treated cells. (b) LS174T cells were transfected with hPXR expression plasmids for 6 h. The cells were treated with vehicle control (0.1% DMSO); 10 μ M CITCO; and 2.5, 10, and 40 μ M PC for 48 h, respectively. The *CYP3A4* mRNA levels were analyzed by real-time PCR. The effect of herbal compounds on *CYP3A4* mRNA levels is presented as fold mRNA expression to control vehicle treated cells. * $P < 0.05$, ** $P < 0.01$ for comparison with the control groups. Values are expressed as mean \pm S.E.M ($n = 3$).



Hindawi
Submit your manuscripts at
<http://www.hindawi.com>

