

POSTER PRESENTATION

Open Access

Integrating multi-scale data for a network model of macaque visual cortex

Maximilian Schmidt^{1*}, Sacha van Albada¹, Rembrandt Bakker^{1,2}, Markus Diesmann^{1,3,4}*From* Twenty Second Annual Computational Neuroscience Meeting: CNS*2013
Paris, France. 13-18 July 2013

Models of cortical dynamics usually either cover small cortical circuits in detail, or represent large patches in a highly simplified manner, for instance using a few differential equations for each area. This is due partly to limited computational resources, and partly to sparsity of large-scale structural connectivity data. These two barriers are now being lifted. Recent developments in the simulation technology NEST [1] and the availability of supercomputers like JUQUEEN in Jülich and the K computer in Japan make it possible to represent substantial parts of primate cortical tissue at cellular and synaptic resolution. At the same time, gaps in experimental knowledge on large-scale connectivity are gradually being closed [2].

Making use of these developments, we construct a model comprising the 32 areas of the macaque cortex associated with visual processing, where the individual areas are based on our recent layered cortical microcircuit model [3]. While this model accounts for the distribution of spike rates across layers and populations, it is heavily underconstrained because only about half the synapses onto the cells in the microcircuit have a local origin. The extension to multiple areas makes it possible to account for a large majority of the synapses and thereby enhances the self-consistency of the model. We compile a realistic structural connectivity map by incorporating multiple datasets from both electro-physiological [4] and anatomical studies [2,5] on different species. The map is adapted to the macaque using realistic area- and layer-specific neuron densities and in-degrees. Both local and inter-area connectivity are layer-specific. The CoCo-Mac database [5] provides the basis for the inter-area connectivity, and is supplemented with quantitative information from further tracing studies [2]. We complete

these data sets by applying empirical connectivity rules, for instance exploiting the dependence of connection strengths on inter-area distances, and of laminar connection patterns on architectural type differences [6].

Combining a simple neuron model with complex connectivity enables us to study the influence of the structural connectivity itself on cortical dynamics. We present preliminary results on basic dynamical features like area- and population-specific firing rates and degrees of irregularity. In addition, we examine the influence of the cortico-cortical connections by comparing dynamical features like power spectra of the isolated primary visual cortex (V1) with Poisson input to its dynamics when embedded in the global network. This model bridges the gap between abstract large-scale models and small-scale microcircuit models. It therefore provides a link between global activity patterns and local spiking activity.

Acknowledgements

We are grateful for the on-going discussion with Claus Hilgetag on cortical architecture. This work was partly supported by EU FP7 Grant 269921 (BrainScaleS), the Helmholtz Alliance on Systems Biology, the Helmholtz Association in the Portfolio theme "Supercomputing and Modeling for the Human Brain", the Jülich Aachen Research Alliance (JARA), the VSR computation time grant JINB33 on the JUGENE supercomputer in Jülich, and the Next-Generation Supercomputer Project of MEXT. All network simulations carried out with NEST (<http://www.nest-initiative.org>).

Author details

¹Institute of Neuroscience and Medicine (INM-6) and Institute for Advanced Simulation (IAS-6), Jülich Research Centre and JARA, Jülich, Germany.

²Donders Institute for Brain, Cognition and Behavior, Radboud University Nijmegen, the Netherlands. ³Faculty of Medicine, RWTH Aachen University, Aachen, Germany. ⁴RIKEN Brain Science Institute, Wako-Shi, Japan.

Published: 8 July 2013

References

1. Gewaltig M-O, Diesmann M: NEST (Neural Simulation Tool). *Scholarpedia* 2007, **2**:1430.

* Correspondence: max.schmidt@fz-juelich.de

¹Institute of Neuroscience and Medicine (INM-6) and Institute for Advanced Simulation (IAS-6), Jülich Research Centre and JARA, Jülich, Germany
Full list of author information is available at the end of the article

2. Markov NT, Misery P, Falchier A, Lamy C, Vezoli J, et al: **Weight consistency specifies regularities of macaque cortical networks.** *Cerebral Cortex* 2011, **21**:1254-1272.
3. Potjans T, Diesmann M: **The cell-type specific cortical microcircuit: relating structure and activity in a full-scale spiking network model.** *Cerebral Cortex* 2012, doi:10.1093/cercor/bhs358.
4. Binzegger T, Douglas RJ, Martin KAC: **A quantitative map of the circuit of cat primary visual cortex.** *J Neurosci* 2004, **39**(24):8441-8453.
5. Stephan KE, Kamper L, Bozkurt A, Burns GAPC, Young MP, Kötter R: **Advanced database methodology for the collation of connectivity data on the macaque brain (CoCoMac).** *Phil Trans R Soc Lond B* 2001, **356**:1159-1186.
6. Hilgetag CC, Burns GAPC, O'Neill MA, Scannell JW, Young MP: **Anatomical connectivity defines the organization of clusters of cortical areas in the macaque monkey and the cat.** *Philos Trans R Soc Lond B Biol Sci* 2000, **355**:91-110.

doi:10.1186/1471-2202-14-S1-P111

Cite this article as: Schmidt et al.: Integrating multi-scale data for a network model of macaque visual cortex. *BMC Neuroscience* 2013 **14**(Suppl 1):P111.

**Submit your next manuscript to BioMed Central
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

