

Poster presentation

Design and synthesis of the first NO- and haem-independent sGC activator BAY 58-2667 for the treatment of acute decompensated heart failure

Michael G Hahn*, Cristina Alonso-Alija, Friederike Stoll, Markus Heil, Joachim Mittendorf, Karl-Heinz Schlemmer, Frank Wunder and Johannes-Peter Stasch

Address: Pharma Research Center, Bayer HealthCare AG, Aprather Weg 18a, 42096 Wuppertal, Germany

Email: Michael G Hahn* - michael.hahn1@bayerhealthcare.com

* Corresponding author

from 3rd International Conference on cGMP Generators, Effectors and Therapeutic Implications
Dresden, Germany. 15–17 June 2007

Published: 25 July 2007

BMC Pharmacology 2007, 7(Suppl 1):P25 doi:10.1186/1471-2210-7-S1-P25

This abstract is available from: <http://www.biomedcentral.com/1471-2210/7/S1/P25>

© 2007 Hahn et al; licensee BioMed Central Ltd.

Soluble guanylate cyclase (sGC) is a signal-transduction enzyme activated by nitric oxide (NO) and plays a key role in a variety of physiological processes such as vasodilatation, antiaggregation, antiproliferation and neuronal signaling as well as in a variety of disorders of these functions. Current therapies that involve the use of organic nitrates and other NO donors have limitations, including non-specific interactions of NO with various

biomolecules and lack of response and the development of tolerance [1,2]. Compounds that activate sGC in an NO-independent manner might therefore provide considerable therapeutic advantages.

High throughput screening led to the identification of a series of dicarboxylic acids as potent sGC activators. Lead structure 1 showed promising *in vitro* potency (sGC over-




Figure 1

Discovery of sGC activators via ultra-high throughput screening using sGC-overexpressing CHO cell lines.

expressing CHO-cells and relaxation of precontracted rabbit arteria saphena rings) and haemodynamic effects in anaesthetized rats (Fig. 1). Extensive chemical derivatization resulted in conclusive structure activity relationships, which will be presented. Eventually, optimization led to the discovery of BAY 58-2667, a highly potent sGC activator which is currently in clinical trials in patients for the treatment of acute decompensated chronic heart failure.

References

1. Stasch JP, Schmidt P, Alonso-Alija C, Apeler H, Dembowski K, Haerter M, Heil M, Minuth T, Perzborn E, Pleiss U, Schramm M, Schroeder W, Schröder H, Stahl E, Steinke W, Wunder F: **NO- and haem-independent activation of soluble guanylyl cyclase: molecular basis and cardiovascular implications of a new pharmacological principle.** *Br J Pharmacol* 2002, **136**:773-783.
2. Evgenov OV, Pacher P, Schmidt PM, Haskó G, Schmidt HHHW, Stasch JP: **NO-independent stimulators and activators of soluble guanylate cyclase: discovery and therapeutic potential.** *Nature Reviews Drug Discovery* 2006, **5**:755-768.

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp

