

Hindawi Publishing Corporation  
Discrete Dynamics in Nature and Society  
Volume 2012, Article ID 619197, 10 pages  
doi:10.1155/2012/619197

## Research Article

# Some Identities on Laguerre Polynomials in Connection with Bernoulli and Euler Numbers

Dae San Kim,<sup>1</sup> Taekyun Kim,<sup>2</sup> and Dmitry V. Dolgy<sup>3</sup>

<sup>1</sup> Department of Mathematics, Sogang University, Seoul 121-742, Republic of Korea

<sup>2</sup> Department of Mathematics, Kwangwoon University, Seoul 139-701, Republic of Korea

<sup>3</sup> Hanrimwon, Kwangwoon University, Seoul 139-701, Republic of Korea

Correspondence should be addressed to Dae San Kim, [dskim@sogang.ac.kr](mailto:dskim@sogang.ac.kr)

Received 15 May 2012; Accepted 28 June 2012

Academic Editor: Lee Chae Jang

Copyright © 2012 Dae San Kim et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

We study some interesting identities and properties of Laguerre polynomials in connection with Bernoulli and Euler numbers. These identities are derived from the orthogonality of Laguerre polynomials with respect to inner product  $\langle f, g \rangle = \int_0^\infty e^{-x^2} f(x)g(x)dx$ .

## 1. Introduction/Preliminaries

As is well known, Laguerre polynomials are defined by the generating function as

$$\frac{\exp(-xt/(1-t))}{1-t} = \sum_{n=0}^{\infty} L_n(x)t^n \quad (1.1)$$

(see [1, 2]). By (1.1), we get

$$\begin{aligned} \sum_{n=0}^{\infty} L_n(x)t^n &= \frac{\exp(-xt/(1-t))}{1-t} = \sum_{r=0}^{\infty} \frac{(-1)^r x^r t^r}{r!} (1-t)^{-(r+1)} \\ &= \sum_{n=0}^{\infty} \sum_{r=0}^{\infty} \frac{(-1)^r x^r (r+s)!}{r!r!s!} t^{r+s} = \sum_{n=0}^{\infty} \left( \sum_{r=0}^n \frac{(-1)^r \binom{n}{r}}{r!} x^r \right) t^n. \end{aligned} \quad (1.2)$$

Thus, from (1.2), we have

$$L_n(x) = \sum_{r=0}^n \frac{(-1)^r \binom{n}{r}}{r!} x^r. \quad (1.3)$$

By (1.3), we see that  $L_n(x)$  is a polynomial of degree  $n$  with rational coefficients and the leading coefficient  $(-1)^n/n!$ . It is well known that Rodrigues' formula is given by

$$L_n(x) = \frac{1}{n!} e^x \left( \frac{d^n}{dx^n} e^{-x} x^n \right) \quad (1.4)$$

(see [1–27]). From (1.1), we can derive the following of Laguerre polynomials:

$$\begin{aligned} L_0(x) &= 1, & L_1(x) &= -x + 1, \\ (n+1)L_{n+1}(x) &= (2n+1-x)L_n(x) - nL_{n-1}(x), & (n \geq 1), \end{aligned} \quad (1.5)$$

$$L'_n(x) = L'_{n-1}(x) - L_{n-1}(x) = 0, \quad (n \geq 1), \quad (1.6)$$

$$xL'_n(x) = nL_n(x) - nL_{n-1}(x) = 0, \quad (n \geq 1). \quad (1.7)$$

By (1.7), we easily see that  $u = L_n(x)$  is a solution of the following differential equation of order 2:

$$xu''(x) + (1-x)u'(x) + nu(x) = 0. \quad (1.8)$$

The Bernoulli numbers,  $B_n$ , are defined by the generating function as

$$\frac{t}{e^t - 1} = e^{Bt} = \sum_{n=0}^{\infty} B_n \frac{t^n}{n!} \quad (1.9)$$

(see [1–28, 28]), with the usual convention about replacing  $B^n$  by  $B_n$ .

It is well known that Bernoulli polynomials of degree  $n$  are given by

$$B_n(x) = (B+x)^n = \sum_{l=0}^n \binom{n}{l} B_{n-l} x^l \quad (1.10)$$

(see [2, 26]). Thus, from (1.10), we have

$$B'_n(x) = \frac{dB_n(x)}{dx} = nB_{n-1}(x) \quad (1.11)$$

(see [3–12]). From (1.9) and (1.10), we can derive the following recurrence relation:

$$B_0 = 1, \quad (B+1)^n - B_n = \delta_{1,n} \quad (1.12)$$

where  $\delta_{n,k}$  is Kronecker's symbol.

The Euler polynomials  $E_n(x)$  are also defined by the generating function as

$$\frac{2}{e^t + 1} e^{xt} = e^{E(x)t} = \sum_{n=0}^{\infty} E_n(x) \frac{t^n}{n!} \quad (1.13)$$

(see [27, 28]), with the usual convention about replacing  $E^n(x)$  by  $E_n(x)$ .

In this special case,  $x = 0$ ,  $E_n(0) = E_n$  are called the  $n$ th Euler numbers. From (1.13), we note that the recurrence formula of  $E_n$  is given by

$$E_0 = 1, \quad (E + 1)^n + E_n = 2\delta_{0,n} \quad (1.14)$$

(see [24]). Finally, we introduce Hermite polynomials, which are defined by

$$e^{2xt-t^2} = e^{H(x)t} = \sum_{n=0}^{\infty} H_n(x) \frac{t^n}{n!} \quad (1.15)$$

(see [29]). In the special case,  $x = 0$ ,  $H_n(0) = H_n$  is called the  $n$ -th Hermite number. By (1.15), we get

$$H_n(x) = (H + 2x)^n = \sum_{l=0}^n \binom{n}{l} H_{n-l} 2^l x^l \quad (1.16)$$

(see [29]). It is not difficult to show that

$$\int_0^{\infty} e^{-x} L_m(x) L_n(x) dx = \delta_{m,n}, \quad (m, n \in \mathbb{Z}_+ = \mathbb{N} \cup \{0\}). \quad (1.17)$$

In the present paper, we investigate some interesting identities and properties of Laguerre polynomials in connection with Bernoulli, Euler, and Hermite polynomials. These identities and properties are derived from (1.17).

## 2. Some Formulae on Laguerre Polynomials in Connection with Bernoulli, Euler, and Hermite Polynomials

Let

$$\mathbf{P}_n = \{p(x) \in \mathbb{Q}[x] \mid \deg p(x) \leq n\}. \quad (2.1)$$

Then  $\mathbf{P}_n$  is an inner product space with the inner product

$$\langle p_1(x), p_2(x) \rangle = \int_0^{\infty} e^{-x} p_1(x) p_2(x) dx, \quad (p_1(x), p_2(x) \in \mathbf{P}_n). \quad (2.2)$$

By (1.17), (2.1), and (2.2), we see that  $L_0(x), L_1(x), \dots, L_n(x)$  are orthogonal basis for  $\mathbf{P}_n$ .

For  $p(x) \in \mathbf{P}_n$ , it is given by

$$p(x) = \sum_{k=0}^n C_k L_k(x), \quad (2.3)$$

where

$$C_k = \langle p(x), L_k(x) \rangle = \int_0^\infty e^{-x} L_k(x) p(x) dx = \frac{1}{k!} \int_0^\infty \left( \frac{d^k}{dx^k} e^{-x} x^k \right) p(x) dx. \quad (2.4)$$

Let us take  $p(x) = x^n \in \mathbf{P}_n$ . From (2.3) and (2.4), we note that

$$\begin{aligned} C_k &= \frac{1}{k!} \int_0^\infty \left( \frac{d^k}{dx^k} e^{-x} x^k \right) x^n dx = \frac{-n}{k!} \int_0^\infty \left( \frac{d^{k-1}}{dx^{k-1}} e^{-x} x^k \right) x^{n-1} dx \\ &= \frac{(-n)(-(n-1))}{k!} \int_0^\infty \left( \frac{d^{k-2}}{dx^{k-2}} e^{-x} x^k \right) x^{n-2} dx \\ &= \dots \\ &= (-1)^k \frac{n(n-1) \cdots (n-k+1)}{k!} \int_0^\infty e^{-x} x^n dx = (-1)^k \binom{n}{k} n!. \end{aligned} \quad (2.5)$$

Therefore, by (2.3), (2.4), and (2.5), we obtain the following theorem.

**Theorem 2.1.** For  $n \in \mathbb{Z}_+$ , one has

$$x^n = n! \sum_{k=0}^n (-1)^k \binom{n}{k} L_k(x). \quad (2.6)$$

Let us consider  $p(x) = B_n(x) \in \mathbf{P}_n$ . Then, by (2.3) and (2.4), we get

$$\begin{aligned} C_k &= \frac{1}{k!} \int_0^\infty \left( \frac{d^k}{dx^k} e^{-x} x^k \right) B_n(x) dx = \frac{-n}{k!} \int_0^\infty \left( \frac{d^{k-1}}{dx^{k-1}} e^{-x} x^k \right) B_{n-1}(x) dx \\ &= \frac{(-n)(-(n-1))}{k!} \int_0^\infty \left( \frac{d^{k-2}}{dx^{k-2}} e^{-x} x^k \right) B_{n-2}(x) dx \\ &= \dots \\ &= (-1)^k \frac{n(n-1) \cdots (n-k+1)}{k!} \sum_{l=0}^{n-k} \binom{n-k}{l} B_{n-k-l} \int_0^\infty e^{-x} x^{k+l} dx \\ &= (-1)^k \binom{n}{k} \sum_{l=0}^{n-k} \binom{n-k}{l} B_{n-k-l} (k+l)! = n! (-1)^k \sum_{l=0}^{n-k} \binom{k+l}{k} \frac{B_{n-k-l}}{(n-k-l)!}. \end{aligned} \quad (2.7)$$

Therefore, by (2.3), (2.4), and (2.7), we obtain the following theorem.

**Theorem 2.2.** For  $n \in \mathbb{Z}_+$ , one has

$$B_n(x) = n! \sum_{k=0}^n \sum_{l=0}^{n-k} (-1)^k \binom{k+l}{k} \frac{B_{n-k-l}}{(n-k-l)!} L_k(x). \quad (2.8)$$

Let us take  $p(x) = E_n(x) \in \mathbf{P}_n$ . By the same method, we easily see that

$$E_n(x) = n! \sum_{k=0}^n \sum_{l=0}^{n-k} (-1)^k \binom{k+l}{k} \frac{E_{n-k-l}}{(n-k-l)!} L_k(x). \quad (2.9)$$

For  $p(x) = H_n(x) \in \mathbf{P}_n$ , we have

$$H_n(x) = \sum_{k=0}^n C_k L_k(x), \quad (2.10)$$

where

$$\begin{aligned} C_k &= \frac{1}{k!} \int_0^\infty \left( \frac{d^k}{dx^k} e^{-x} x^k \right) H_n(x) dx = \frac{-2n}{k!} \int_0^\infty \left( \frac{d^{k-1}}{dx^{k-1}} e^{-x} x^k \right) H_{n-1}(x) dx \\ &= \frac{(-2n)(-2(n-1))}{k!} \int_0^\infty \left( \frac{d^{k-2}}{dx^{k-2}} e^{-x} x^k \right) H_{n-2}(x) dx \\ &= \dots \\ &= \frac{(-2n)(-2(n-1)) \cdots (-2(n-k+1))}{k!} \int_0^\infty e^{-x} x^k H_{n-k}(x) dx \\ &= \frac{(-1)^k 2^k n(n-1) \cdots (n-k+1)}{k!} \sum_{l=0}^{n-k} \binom{n-k}{l} H_{n-k-l} 2^l \int_0^\infty e^{-x} x^{k+l} dx \\ &= (-1)^k \binom{n}{k} \sum_{l=0}^{n-k} \binom{n-k}{l} 2^{k+l} H_{n-k-l} (k+l)! = n! (-1)^k \sum_{l=0}^{n-k} 2^{k+l} \binom{k+l}{k} \frac{H_{n-k-l}}{(n-k-l)!}. \end{aligned} \quad (2.11)$$

Therefore, by (2.10) and (2.11), we obtain the following theorem.

**Theorem 2.3.** For  $n \in \mathbb{Z}_+$ , one has

$$H_n(x) = n! \sum_{k=0}^n \sum_{l=0}^{n-k} (-1)^k 2^{k+l} \binom{k+l}{k} \frac{H_{n-k-l}}{(n-k-l)!} L_k(x). \quad (2.12)$$

Let  $p(x) = \sum_{k=0}^n B_k(x) B_{n-k}(x) \in \mathbf{P}_n$ . Then we have

$$p(x) = \sum_{k=0}^n B_k(x) B_{n-k}(x) = \sum_{k=0}^n C_k L_k(x), \quad (2.13)$$

where

$$C_k = \frac{1}{k!} \int_0^\infty \left( \frac{d^k}{dx^k} e^{-x} x^k \right) p(x) dx. \quad (2.14)$$

In [15], it is known that

$$\sum_{k=0}^n B_k(x) B_{n-k}(x) = \frac{2}{n+2} \sum_{l=0}^{n-2} \binom{n+2}{l} B_{n-l} B_l(x) + (n+1) B_n(x). \quad (2.15)$$

By (2.14) and (2.15), we get

$$\begin{aligned} C_k = \frac{1}{k!} \left\{ \frac{2}{n+2} \sum_{l=0}^{n-2} \binom{n+2}{l} B_{n-l} \int_0^\infty \left( \frac{d^k}{dx^k} e^{-x} x^k \right) B_l(x) dx \right. \\ \left. + (n+1) \int_0^\infty \left( \frac{d^k}{dx^k} e^{-x} x^k \right) B_n(x) dx \right\}. \end{aligned} \quad (2.16)$$

From (2.16), we can derive the following equations ((2.17)-(2.18)):

$$C_n = (-1)^n (n+1)!, \quad C_{n-1} = n(n+1)!(-1)^{n-1} - \frac{1}{2}(-1)^{n-1}(n+1)!. \quad (2.17)$$

For  $0 \leq k \leq n-2$ , we have

$$\begin{aligned} C_k = \frac{2}{n+2} \sum_{l=k}^{n-2} \sum_{m=0}^{n-2-l-k} \binom{n+2}{l} \binom{k+m}{k} l! (-1)^k B_{n-l} \frac{B_{l-k-m}}{(l-k-m)!} \\ + (-1)^k (n+1)! \sum_{m=0}^{n-k} \binom{k+m}{k} \frac{B_{n-k-m}}{(n-k-m)!}. \end{aligned} \quad (2.18)$$

Therefore, by (2.13), (2.17), and (2.18), we obtain the following theorem.

**Theorem 2.4.** For  $n \in \mathbb{Z}_+$ , one has

$$\begin{aligned} \sum_{k=0}^n B_k(x) B_{n-k}(x) = \sum_{k=0}^{n-2} \left\{ \frac{2}{n+2} \sum_{l=k}^{n-2} \sum_{m=0}^{n-2-l-k} (-1)^k l! \binom{n+2}{l} \binom{k+m}{k} B_{n-l} \frac{B_{l-k-m}}{(l-k-m)!} \right. \\ \left. + (-1)^k (n+1)! \sum_{m=0}^{n-k} \binom{k+m}{k} \frac{B_{n-k-m}}{(n-k-m)!} \right\} L_k(x) \\ + \left( n(n+1)!(-1)^{n-1} - \frac{1}{2}(-1)^{n-1}(n+1)! \right) L_{n-1}(x) + (-1)^n (n+1)! L_n(x). \end{aligned} \quad (2.19)$$

Let us take  $p(x) = \sum_{k=0}^n E_k(x)E_{n-k}(x) \in \mathbf{P}_n$ . By (2.3) and (2.4), we get

$$p(x) = \sum_{k=0}^n E_k(x)E_{n-k}(x) = \sum_{k=0}^n C_k L_k(x), \quad (2.20)$$

where

$$C_k = \frac{1}{k!} \int_0^\infty \left( \frac{d^k}{dx^k} e^{-x} x^k \right) p(x) dx. \quad (2.21)$$

It is known (see [15]) that

$$\sum_{k=0}^n E_k(x)E_{n-k}(x) = \sum_{k=0}^{n-1} \frac{(n+1)\binom{n}{k}}{n-k+1} \left( \sum_{l=k}^n E_{l-k}E_{n-l} - 2E_{n-k} \right) E_k(x) + 2(n+1)E_n(x). \quad (2.22)$$

From (2.20), (2.21), and (2.22), we can derive the following equations ((2.23)-(2.24)):

$$C_n = \frac{(-1)^n}{n!} 2(n+1)(n!)^2 = 2(-1)^n (n+1)!. \quad (2.23)$$

For  $0 \leq k \leq n-1$ , we have

$$\begin{aligned} C_k &= \sum_{l=k}^{n-1} \frac{(n+1)\binom{n}{l}}{(n-l+1)!} \left( \sum_{m=l}^n E_{m-l}E_{n-m} - 2E_{n-l} \right) \sum_{p=0}^{l-k} (-1)^k l! \binom{k+p}{k} \frac{E_{l-k-p}}{(l-k-p)!} \\ &\quad + 2(n+1)! (-1)^k \sum_{p=0}^{n-k} \binom{k+p}{k} \frac{E_{n-k-p}}{(n-k-p)!}. \end{aligned} \quad (2.24)$$

Therefore, by (2.20) and (2.24), we obtain the following theorem.

**Theorem 2.5.** For  $n \in \mathbb{Z}_+$ , one has

$$\begin{aligned} \sum_{k=0}^n E_k(x)E_{n-k}(x) &= \sum_{k=0}^{n-1} \left\{ \sum_{l=k}^{n-1} \frac{(n+1)\binom{n}{l}}{(n-l+1)!} \left( \sum_{m=l}^n E_{m-l}E_{n-m} - 2E_{n-l} \right) \sum_{p=0}^{l-k} (-1)^k l! \binom{k+p}{k} \right. \\ &\quad \left. \times \frac{E_{l-k-p}}{(l-k-p)!} + 2(n+1)! (-1)^k \sum_{p=0}^{n-k} \binom{k+p}{k} \frac{E_{n-k-p}}{(n-k-p)!} \right\} L_k(x) \\ &\quad + 2(-1)^n (n+1)! L_n(x). \end{aligned} \quad (2.25)$$

It is known that

$$\sum_{k=0}^n E_k(x)E_{n-k}(x) = p(x) = -\frac{4}{n+2} \sum_{k=0}^n \binom{n+2}{k} E_{n-k+1} B_k(x) \quad (2.26)$$

(see [15]). From (2.20), (2.21), and (2.23), we have

$$\begin{aligned} C_k &= \frac{1}{k!} \int_0^\infty \left( \frac{d^k}{dx^k} e^{-x} x^k \right) p(x) dx \\ &= -\frac{4}{n+2} \sum_{l=k}^n \binom{n+2}{l} E_{n-l+1} \frac{1}{k!} \int_0^\infty \left( \frac{d^k}{dx^k} e^{-x} x^k \right) B_l(x) dx \\ &= -\frac{4}{n+2} \sum_{l=k}^n \sum_{m=0}^{l-k} \binom{n+2}{l} (-1)^k E_{n-l+1} \frac{B_{l-k-m}}{(l-k-m)!} l! \binom{m+k}{k}. \end{aligned} \quad (2.27)$$

Therefore, by (2.20) and (2.27), we obtain the following theorem.

**Theorem 2.6.** For  $n \in \mathbb{Z}_+$ , one has

$$\begin{aligned} &\sum_{k=0}^n E_k(x)E_{n-k}(x) \\ &= -\frac{4}{n+2} \sum_{k=0}^n \sum_{l=k}^n \sum_{m=0}^{l-k} \binom{n+2}{l} (-1)^k E_{n-l+1} \frac{B_{l-k-m}}{(l-k-m)!} l! \binom{m+k}{k} L_k(x). \end{aligned} \quad (2.28)$$

*Remark 2.7.* Laguerre's differential equation

$$ty'' + (1-t)y' + ny = 0 \quad (2.29)$$

is known to possess polynomial solutions when  $n$  is a nonnegative integer. These solutions are naturally called Laguerre polynomials and are denoted by  $L_n(t)$ . That is,  $y = L_n(t)$  are solutions of (2.29) which are given by

$$y = L_n(t) = \sum_{r=0}^n \frac{\binom{n}{r} (-1)^r}{r!} t^r, \quad L_0(1) = 1. \quad (2.30)$$

From (2.30), we note that Laplace transform of  $y = L_n(t)$  is given by

$$\mathcal{L}(y) = \mathcal{L}(L_n(t)) = \frac{1}{s} \sum_{r=0}^n \binom{n}{r} (-1)^r \left( \frac{1}{s} \right)^r = \frac{(s-1)^n}{s^{n+1}}. \quad (2.31)$$



It is not difficult to show that

$$\mathcal{L}\left(\frac{e^t}{n!}\left(\frac{d^n}{dt^n}e^{-t}t^n\right)\right) = \mathcal{L}(y) = \frac{(s-1)^n}{s^{n+1}}. \quad (2.32)$$


Thus, we conclude that

$$L_n(t) = \frac{e^t}{n!}\left(\frac{d^n}{dt^n}e^{-t}t^n\right), \quad \text{for } n \in \mathbb{Z}_+. \quad (2.33)$$

## References

- [1] M. A. Özarslan and C. Kaanoğlu, "Multilateral generating functions for classes of polynomials involving multivariable Laguerre polynomials," *Journal of Computational Analysis and Applications*, vol. 13, no. 4, pp. 683–691, 2011.
- [2] A. K. Shukla and S. K. Meher, "Group-theoretic origin of some generating functions for Laguerre polynomials of two variables," *Applied Mathematical Sciences*, vol. 5, no. 13–16, pp. 775–784, 2011.
- [3] S. Araci, "Novel identities for  $q$ -genocchi numbers and polynomials," *Journal of Functions Spaces and Applications*. In press.
- [4] S. Araci, D. Erdal, and J. J. Seo, "A study on the fermionic  $p$ -adic  $q$ -integral representation on  $\mathbb{Z}_p$  associated with weighted  $q$ -Bernstein and  $q$ -Genocchi polynomials," *Abstract and Applied Analysis*, vol. 2011, Article ID 649248, 10 pages, 2011.
- [5] A. Bayad, "Arithmetical properties of elliptic Bernoulli and Euler numbers," *International Journal of Algebra*, vol. 4, no. 5–8, pp. 353–372, 2010.
- [6] A. Bayad, "Modular properties of elliptic Bernoulli and Euler functions," *Advanced Studies in Contemporary Mathematics*, vol. 20, no. 3, pp. 389–401, 2010.
- [7] A. Bayad, "Fourier expansions for Apostol-Bernoulli, Apostol-Euler and Apostol-Genocchi polynomials," *Mathematics of Computation*, vol. 80, no. 276, pp. 2219–2221, 2011.
- [8] A. Bayad and T. Kim, "Identities involving values of Bernstein,  $q$ -Bernoulli, and  $q$ -Euler polynomials," *Russian Journal of Mathematical Physics*, vol. 18, no. 2, pp. 133–143, 2011.
- [9] I. N. Cangul, V. Kurt, H. Ozden, and Y. Simsek, "On the higher-order  $w$ - $q$ -Genocchi numbers," *Advanced Studies in Contemporary Mathematics*, vol. 19, no. 1, pp. 39–57, 2009.
- [10] M. Cenkci, "The  $p$ -adic generalized twisted  $(h, q)$ -Euler- $l$ -function and its applications," *Advanced Studies in Contemporary Mathematics*, vol. 15, no. 1, pp. 37–47, 2007.
- [11] M. W. Coffey, "On finite sums of Laguerre polynomials," *Rocky Mountain Journal of Mathematics*, vol. 41, no. 1, pp. 79–93, 2011.
- [12] M. Gülsu, B. Gürbüz, Y. Öztürk, and M. Sezer, "Laguerre polynomial approach for solving linear delay difference equations," *Applied Mathematics and Computation*, vol. 217, no. 15, pp. 6765–6776, 2011.
- [13] N. S. Jung, H. Y. Lee, and C. S. Ryoo, "Some relations between twisted  $(h, q)$ -Euler numbers with weight  $\alpha$  and  $q$ -Bernstein polynomials with weight  $\alpha$ ," *Discrete Dynamics in Nature and Society*, vol. 2011, Article ID 176296, 11 pages, 2011.
- [14] D. S. Kim, D. V. Dolgy, T. Kim, and S.-H. Rim, "Some formulae for the product of two bernoulli and euler polynomials," *Abstract and Applied Analysis*, vol. 2012, Article ID 784307, 15 pages, 2012.
- [15] D. S. Kim, D. V. Dolgy, H. M. Kim, S. H. Lee, and T. Kim, "Integral formulae of bernoulli polynomials," *Discrete Dynamics in Nature and Society*, vol. 2012, Article ID 269847, 15 pages, 2012.
- [16] M.-S. Kim, "A note on sums of products of Bernoulli numbers," *Applied Mathematics Letters*, vol. 24, no. 1, pp. 55–61, 2011.
- [17] T. Kim, "On the weighted  $q$ -Bernoulli numbers and polynomials," *Advanced Studies in Contemporary Mathematics*, vol. 21, no. 2, pp. 207–215, 2011.
- [18] T. Kim, "Some identities on the  $q$ -Euler polynomials of higher order and  $q$ -Stirling numbers by the fermionic  $p$ -adic integral on  $\mathbb{Z}_p$ ," *Russian Journal of Mathematical Physics*, vol. 16, no. 4, pp. 484–491, 2009.
- [19] T. Kim, "Symmetry of power sum polynomials and multivariate fermionic  $p$ -adic invariant integral on  $\mathbb{Z}_p$ ," *Russian Journal of Mathematical Physics*, vol. 16, no. 1, pp. 93–96, 2009.

- [20] H. Y. Lee, N. S. Jung, and C. S. Ryoo, "A note on the  $q$ -Euler numbers and polynomials with weak weight  $\alpha$ ," *Journal of Applied Mathematics*, Article ID 497409, 14 pages, 2011.
- [21] H. Ozden, Y. Simsek, and I. N. Cangul, "Euler polynomials associated with  $p$ -adic  $q$ -Euler measure," *General Mathematics*, vol. 15, no. 2, pp. 24–37, 2007.
- [22] H. Ozden, I. N. Cangul, and Y. Simsek, "Multivariate interpolation functions of higher-order  $q$ -Euler numbers and their applications," *Abstract and Applied Analysis*, vol. 2088, Article ID 390857, 16 pages, 2008.
- [23] C. S. Ryoo, "Some identities of the twisted  $q$ -Euler numbers and polynomials associated with  $q$ -Bernstein polynomials," *Proceedings of the Jangjeon Mathematical Society*, vol. 14, no. 2, pp. 239–248, 2011.
- [24] C. S. Ryoo, "Some relations between twisted  $q$ -Euler numbers and Bernstein polynomials," *Advanced Studies in Contemporary Mathematics*, vol. 21, no. 2, pp. 217–223, 2011.
- [25] P. Rusev, "Laguerre's polynomials and the nonreal zeros of Riemann's  $\zeta$ -function," *Comptes Rendus de l'Académie Bulgare des Sciences*, vol. 63, no. 11, pp. 1547–1550, 2010.
- [26] D. Zill and M. R. Cullen, *Advanced Engineering Mathematics*, Jones and Bartlett, 2005.
- [27] Y. Simsek, "Construction a new generating function of Bernstein type polynomials," *Applied Mathematics and Computation*, vol. 218, no. 3, pp. 1072–1076, 2011.
- [28] Y. Simsek and M. Acikgoz, "A new generating function of ( $q$ -) Bernstein-type polynomials and their interpolation function," *Abstract and Applied Analysis*, vol. 2010, Article ID 769095, 12 pages, 2010.
- [29] R. Koekoek, P. A. Lesky, and R. F. Swarttouw, *Hypergeometric Orthogonal Polynomials and Their  $q$ -Analogues*, Springer, Berlin, Germany, 2010.



# Hindawi

Submit your manuscripts at  
<http://www.hindawi.com>

