

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

---

USDA National Wildlife Research Center - Staff  
Publications

U.S. Department of Agriculture: Animal and  
Plant Health Inspection Service

---

January 2001

## IMPACT OF BLACKBIRD DAMAGE TO SUNFLOWER BIOENERGETIC AND ECONOMIC MODELS

Brian D. Peer  
*North Dakota State University, Fargo*

H. Jeffrey Homan  
*USDA/National Wildlife Research Center/Great Plains Field Station*

George M. Linz  
*USDA/APHIS/WS National Wildlife Research Center, george\_m\_linz@yahoo.com*

William J. Bleier  
*North Dakota State University, Fargo*

Follow this and additional works at: [https://digitalcommons.unl.edu/icwdm\\_usdanwrc](https://digitalcommons.unl.edu/icwdm_usdanwrc)



Part of the [Environmental Sciences Commons](#)

---

Peer, Brian D.; Homan, H. Jeffrey ; Linz, George M.; and Bleier, William J., "IMPACT OF BLACKBIRD DAMAGE TO SUNFLOWER BIOENERGETIC AND ECONOMIC MODELS" (2001). *USDA National Wildlife Research Center - Staff Publications*. 560.  
[https://digitalcommons.unl.edu/icwdm\\_usdanwrc/560](https://digitalcommons.unl.edu/icwdm_usdanwrc/560)

This Article is brought to you for free and open access by the U.S. Department of Agriculture: Animal and Plant Health Inspection Service at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in USDA National Wildlife Research Center - Staff Publications by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

Published in *Proceedings of the 23rd Sunflower Research Workshop* (January 17–18, 2001, Fargo, ND). National Sunflower Association, Bismarck, North Dakota, USA.

Research articles from subsequent workshops (2002– ) are available online at :

<http://www.sunflowernsa.com/research/default.asp?contentID=70>

The National Sunflower Association (NSA) is a non-profit commodity organization working on problems and opportunities for the improvement of all members. Membership in the NSA includes growers and the support industry.

NSA Contact Information:

**National Sunflower Association**  
**4023 State Street**  
**Bismarck, ND 58503-0690**

<http://www.sunflowernsa.com/>

(701) 328-5100  
fax (701) 328-5101  
(888) 718-7033



## IMPACT OF BLACKBIRD DAMAGE TO SUNFLOWER: BIOENERGETIC AND ECONOMIC MODELS

BRIAN D. PEER<sup>1</sup>, H. JEFFREY HOMAN<sup>2</sup>, GEORGE M. LINZ<sup>2</sup>, and WILLIAM J. BLEIER<sup>1</sup>

<sup>1</sup>Department of Zoology, North Dakota State University, Fargo, ND 58105

<sup>2</sup>USDA/National Wildlife Research Center/Great Plains Field Station,  
2110 Miriam Circle, Bismarck, ND 58501

We constructed bioenergetic models to assess the economic impact on sunflower production from a regional population of blackbirds (Icteridae) consisting of three species: red-winged blackbirds (*Agelaius phoeniceus*), common grackles (*Quiscalus quiscula*), and yellow-headed blackbirds (*Xanthocephalus xanthocephalus*). Values generated from the bioenergetic models were used to perform a cost-benefit analysis to determine efficacy of a proposed avicide baiting program to manage red-winged blackbirds residing in the Central Lowlands and Northern Great Plains regions of North America. The bioenergetic model included metabolic rates, caloric value and moisture content of achenes, and percentage of sunflower in the diets of males and females of each species. The following estimates of annual sunflower consumption are per male and female, respectively: red-winged blackbird 265 g (9.3 oz) and 159 g (5.6 oz); common grackle, 257 g (9.1 oz) and 220 g (7.8 oz); and yellow-headed blackbird, 237 g (8.4 oz) and 132 g (4.6 oz). Based on the average price of sunflower from 1993-1997, male red-winged blackbirds damaged \$0.09 and \$0.18 (respective annual and lifetime damage per bird), females \$0.05 and \$0.21; male common grackles \$0.08 and \$0.12, females \$0.07 and \$0.22; and male yellow-headed blackbirds \$0.08 and \$0.15, females \$0.04 and \$0.14. The population of all three species combined impacted production by \$5.2 million, annually. Lifetime damage was \$13.3 million. The annual cost of baiting 2 million red-winged blackbirds with avicide would be \$65,000. A baiting program would yield savings of \$73,000 based on annual damage estimates and \$325,000 based on lifetime damage estimates.